Supplemental Table 1. Immunosuppressive medications in patients who received hepatitis B booster

vaccination

Number of	Immunosuppressive medications received
patients	
7	Corticosteroid only
6	Tacrolimus only
5	Corticosteroid and mycophenolate mofetil (MMF)
2	MMF only
2	Corticosteroid and tacrolimus
1	Azathioprine only
1	Plasmapheresis only
1	Rituximab only
1	Tacrolimus and MMF
1	Corticosteroid and Azathioprine
1	Corticosteroid and sirolimus
1	Corticosteroid, MMF, and sirolimus
1	Corticosteroid, MMF, and tacrolimus
1	Corticosteroid, cyclophosphamide, and plasmapheresis
1	Corticosteroid, MMF, and cyclosporine
1	Carboplatin, cyclophosphamide, etoposide, and doxorubicin

Medical Center	1	2	3	4	5	6	7	8	9	10	11	12
Single dose 10 mcg (n)	0	4	0	0	4	0	0	2	6	1	1	Unknown
Single dose 20 mcg (n)	1	9	14	51	0	0	20	0	0	0	0	Unknown
Single dose 40 mcg (n)	17	5	1	0	2	7	0	0	1	0	0	Unknown
Multi-dose 10 mcg (n)	0	7	0	0	3	0	0	7	3	1	0	Unknown
Multi-dose 20 mcg (n)	0	5	0	5	2	0	10	0	0	2	3	Unknown
Multi-dose 40 mcg (n)	2	9	0	0	0	0	0	0	0	0	0	Unknown
% given as multi-dose booster series	10%	53%	0%	9%	45%	0%	33%	70%	30%	75%	75%	100%
% given to patients receiving immuno- suppressive drugs	20%	28%	0%	11%	0%	29%	7%	25%	11%	20%	50%	25%
% given as repeat booster*	19%	21%	25%	28%	10%	40%	0%	0%	0%	0%	0%	0%
Mean total hepatitis B vaccine dose (mcg)**	43	24	21	22	27	40	33	23	16	40	36	Unknown

Supplemental Table 2. Hepatitis B vaccination practice at each center

\*Proportion of hepatitis B booster vaccinations at each center given to patients who had previously received hepatitis B booster vaccination during the study period.

\*\* Total vaccine dose is defined as the cumulative dose for a multi-dose booster series or single dose booster. For example, a 3 dose series of 20 mcg/dose would equal a total vaccine dose of 60 mcg, while a single-dose booster of 20 mcg would equal a total vaccine dose of 20 mcg.