
Supplemental table 1. Univariate and multivariate COX analysis predicting fatal and non-fatal cardiovascular events (a composite of 103 fatal and non-fatal events) in patients without previous history of cardiovascular disease.

	Univariate COX			Multivariate COX		
	HR	**95% CI**	**P**	**HR**	**95% CI**	**P**
FMD (%)	0.66	0.45-0.78	**<0.001**			
IL-10 tertiles (pg/ml)	2.41	1.81-3.21	**<0.001**	2.23	1.65-3.02	**<0.001**
IL-6 tertiles (pg/ml)	1.29	1.01-1.65	**0.04**			
IL6/IL10 ratio	0.11	0.05-0.32	**<0.001**			
Pentraxin 3 (ng/ml)	1.02	1.01-1.04	**<0.001**	1.01	1.01-1.03	**0.04**
hsCRP(mg/l)	1.03	1.01-1.05	**0.02**			
Serum albumin (g/dl)	0.62	0.34-1.12	**0.11**			
eGFR (ml/min)	0.99	0.98-0.99	**<0.001**			
Diabetes (Yes/no)	5.11	3.62-7.23	**<0.001**	4.26	2.87-6.31	**<0.001**
Hypertension (Yes/no)	2.56	1.66-3.94	**<0.001**			
Smoking (Yes/no)	1.75	1.18-2.58	**0.005**	1.82	1.22-2.72	**0.003**
24 hour proteinuria (mg/day)	1.00	1.00-1.00	0.79			
SBP (mmHg)	1.03	1.02-1.04	**<0.001**			
LDL Cholesterol (mg/dl)	1.01	0.99-1.01	0.90			
HOMA	1.37	1.17-1.61	**<0.001**			
Calcium (mg/dl)	0.90	0.64-1.27	0.35			
Phosphate (mg/dl)	1.08	0.95-1.23	**0.22**			
iPTH (pg/ml)	1.01	1.00-1.01	**<0.001**			
Age (years)	1.01	0.99-1.02	0.43			
Gender (M/F)	1.31	0.91-1.87	0.14			

Supplemental figure 1. Kaplan-Meier survival curves (a composite of 103 fatal and nonfatal events) according to lower median and higher IL-10 levels) in patients without previous history of cardiovascular disease.



Supplemental figure 2. Kaplan-Meier survival curves (a composite of 103 fatal and nonfatal events) according to lower and higher median IL-6/ IL-10 ratio) in patients without previous history of cardiovascular disease.

