

Authors: Anna C. Porter MD,¹ James P. Lash MD,¹ Dawei Xie PhD,^{2,3} Qiang Pan MA,³ Jennifer DeLuca,⁴ Radhika Kanthety MD,⁵ John W. Kusek PhD,⁶ Claudia M. Lora MD,¹ Lisa Nessel MSS, MLSP³ Ana C. Ricardo MD MPH,¹ Julie Wright Nunes MD,⁷ Michael J. Fischer MD MSPH^{1,8} and the CRIC Study Investigators*

¹Medicine, Section of Nephrology, University of Illinois Hospital and Health Sciences System and Jesse Brown VA Medical Center, Chicago, IL

²Renal, Electrolyte and Hypertension Division, Perelman School of Medicine at the University of Pennsylvania. Philadelphia, PA

³Department of Biostatistics and Epidemiology, University of Pennsylvania. Philadelphia, PA.

⁴St. John's Health System, Detroit, MI

⁵Division of Nephrology and Hypertension, University Hospitals, Case Medical Center, Case Western Reserve University School of Medicine, Cleveland, OH

⁶National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD

⁷Division of Nephrology, University of Michigan at Ann Arbor, Ann Arbor, MI

⁸Center of Innovation for Complex Chronic Healthcare, Edward Hines Jr., VA Hospital, Hines, IL

⁹Departments of Medicine and Epidemiology Johns Hopkins University, Baltimore, MD

¹⁰Division of Research, Kaiser Permanente Northern California; Oakland, CA

¹¹Departments of Epidemiology and Medicine, Tulane University School of Public Health and Tropical Medicine, New Orleans, Louisiana

* CRIC Study Investigators include: Lawrence J. Appel, MD, MPH;⁹ Harold I. Feldman, MD, MSCE;² Alan S. Go, MD;¹⁰ Jiang He, MD, PhD;¹¹ John W. Kusek, PhD;⁶ James P. Lash, MD;¹ Akinlolu Ojo, MD, PhD;⁷ Mahboob Rahman, MD;⁵ and Raymond R. Townsend, MD.²