

Anke Doyon¹, Bettina Schmiedchen², Anja Sander³, Aysun Bayazit⁴, Ali Duzova⁵, Nur Canpolat⁶, Daniela Thurn⁷, Karolis Azukaitis⁸, Ali Anarat⁴, Justine Bacchetta⁸, Sevgi Mir⁹, Rukshana Shroff¹⁰, Ebru Yilmaz¹¹, Cengiz Candan¹², Markus Kemper¹³, Michel Fischbach¹⁴, Gerard Cortina¹⁵, Günter Klaus¹⁶, Matthias Wuttke¹⁷, Anna Köttgen¹⁷, Anette Melk⁷, Uwe Querfeld¹⁸, Franz Schaefer¹, for the 4C Study Consortium

¹ Pediatric Nephrology Division, Center for Pediatrics and Adolescent Medicine, Heidelberg, Germany

² Department of Physiology and Pathophysiology, Institute of Nutritional Science, University of Potsdam

³ Institute of Medical Biometry and Informatics, University of Heidelberg, Heidelberg, Germany

⁴ Division of Pediatric Nephrology, Adana University, Adana, Turkey

⁵ Division of Pediatric Nephrology, Hacettepe University Faculty of Medicine, Ankara, Turkey

⁶ Division of Pediatric Nephrology, Istanbul University Cerrahpasa Medical Faculty, Istanbul, Turkey

⁷ Department of Pediatric Nephrology, Hannover Medical School, Hannover, Germany

⁸ Center for Pediatrics, Vilnius University, Vilnius, Lithuania

⁸ Service de Néphrologie pédiatrique, Hôpital femme, mère, enfant, Lyon, France

⁹ Division of Pediatric Nephrology, Ege University Medical Faculty, Izmir, Turkey

¹⁰ Paediatric Nephrology and Urology Unit, Great Ormond Street Hospital, London, UK

¹¹ Pediatric Nephrology, Sanliurfa Children's Hospital, Sanliurfa, Turkey

¹² Pediatric Nephrology, Goztepe Educational and Research Hospital, Istanbul, Turkey

¹³ Pediatric Nephrology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

¹⁴ Hôpital de Hautepierre, Service de Pédiatrie 3, Strasbourg, France

¹⁵ Department of Pediatrics, Medical University of Innsbruck, Austria

¹⁶ KfH Children's Kidney Center, Marburg, Germany

¹⁷ Division of Nephrology, Medical Center - University of Freiburg, Germany

¹⁸ Clinic of Pediatric Nephrology, Charite Children's Hospital, Berlin, Germany