

Spectrum of Steroid-Resistant and Congenital Nephrotic Syndrome in Children: The PodoNet Registry Cohort

Agnes Trautmann, Monica Bodria, Fatih Ozaltin, Alaleh Gheisari, Anette Melk, Marta Azocar, Ali Anarat, Salim Caliskan, Francesco Emma, Jutta Gellermann, Jun Oh, Esra Baskin, Joanna Ksiazek, Giuseppe Remuzzi, Ozlem Erdogan, Sema Akman, Jiri Dusek, Tinatin Davitaia, Ozan Özkaya, Fotios Papachristou, Agnieszka Firszt-Adamczyk, Tomasz Urasinski, Sara Testa, Rafael T. Krmar, Lidia Hyla-Klekot, Andrea Pasini, Z. Birsin Özcahar, Peter Sallay, Nilgun Cakar, Monica Galanti, Joelle Terzic, Bilal Aoun, Alberto Caldas Afonso, Hanna Szymanik-Grzelak, Beata S. Lipska, Sven Schnaidt, and Franz Schaefer for the PodoNet Consortium

Abstract

Background and objectives Steroid-resistant nephrotic syndrome is a rare kidney disease involving either immune-mediated or genetic alterations of podocyte structure and function. The rare nature, heterogeneity, and slow evolution of the disorder are major obstacles to systematic genotype-phenotype, intervention, and outcome studies, hampering the development of evidence-based diagnostic and therapeutic concepts. To overcome these limitations, the PodoNet Consortium has created an international registry for congenital nephrotic syndrome and childhood-onset steroid-resistant nephrotic syndrome.

Design, setting, participants, & measurements Since August of 2009, clinical, biochemical, genetic, and histopathologic information was collected both retrospectively and prospectively from 1655 patients with childhood-onset steroid-resistant nephrotic syndrome, congenital nephrotic syndrome, or persistent subnephrotic proteinuria of likely genetic origin at 67 centers in 21 countries through an online portal.

Results Steroid-resistant nephrotic syndrome manifested in the first 5 years of life in 64% of the patients. Congenital nephrotic syndrome accounted for 6% of all patients. Extrarenal abnormalities were reported in 17% of patients. The most common histopathologic diagnoses were FSGS (56%), minimal change nephropathy (21%), and mesangioproliferative GN (12%). Mutation screening was performed in 1174 patients, and a genetic disease cause was identified in 23.6% of the screened patients. Among 14 genes with reported mutations, abnormalities in *NPHS2* ($n=138$), *WT1* ($n=48$), and *NPHS1* ($n=41$) were most commonly identified. The proportion of patients with a genetic disease cause decreased with increasing manifestation age: from 66% in congenital nephrotic syndrome to 15%–16% in schoolchildren and adolescents. Among various intensified immunosuppressive therapy protocols, calcineurin inhibitors and rituximab yielded consistently high response rates, with 40%–45% of patients achieving complete remission. Confirmation of a genetic diagnosis but not the histopathologic disease type was strongly predictive of intensified immunosuppressive therapy responsiveness. Post-transplant disease recurrence was noted in 25.8% of patients without compared with 4.5% ($n=4$) of patients with a genetic diagnosis.

Conclusions The PodoNet cohort may serve as a source of reference for future clinical and genetic research in this rare but significant kidney disease.

Clin J Am Soc Nephrol 10: 592–600, 2015. doi: 10.2215/CJN.06260614

Introduction

Approximately 12%–15% of children with idiopathic nephrotic syndrome do not respond to oral steroid therapy (1,2). Steroid-resistant nephrotic syndrome (SRNS) is a challenging clinical condition with highly variable outcomes, and 50% of children progress to ESRD within 15 years (3,4). Although in some patients, temporary or persistent remission is achieved by intensified immunosuppressive therapies (IITs), others exhibit a multidrug-resistant phenotype. Historically, diagnostic evaluation and prognostic classification relied largely on histopathologic assessment. In recent years, abnormalities in a growing number of

genes essential for podocyte development, structure, and function have been identified in patients with congenital nephrotic syndrome (CNS) and SRNS (5). The ongoing discovery of genetic podocytopathies is about to redefine the physiopathologic understanding, diagnostic assessment, prognostic judgment, and therapeutic approaches in childhood-onset SRNS. However, the development of evidence-based management algorithms has been hampered by the low incidence of SRNS, which is estimated at 2–4 per million person-years (1).

To overcome the limitations imposed by the rarity of the disorder, the PodoNet Consortium has created

Due to the number of contributing authors, the affiliations are provided in the Supplemental Material.

Correspondence:
Professor Franz Schaefer, Division of Pediatric Nephrology, Center for Pediatrics and Adolescent Medicine, Im Neuenheimer Feld 430, 69120 Heidelberg, Germany.
Email: franz.schaefer@med.uni-heidelberg.de

an international registry for CNS and childhood-onset SRNS. Over the past 5 years, comprehensive clinical, medication-related, histopathologic, and genetic information from >1600 children with the disorder has been collected in 21 countries. The detailed description of the PodoNet cohort presented here aims to provide a source of reference for future clinical and genetic research in CNS and SRNS.

Materials and Methods

Registry Description

The PodoNet Consortium was formed in 2008 by research groups from Heidelberg, Paris, Rome, Bergamo, Genova, and Ankara. The PodoNet project (www.podonet.org) encompasses clinical, genetic, and experimental research into hereditary podocyte disorders. The clinical activities encompass a web-based international clinical registry and a central biobank for SRNS. To date, investigators from 67 clinical units in 21 countries of Europe, the Middle East, and Latin America have joined the PodoNet Registry Consortium. The registry study protocol was approved by the local institutional review boards/ethics committees, and written informed consent is obtained from the families. Data protection is ensured by pseudonymized data input.

The registry accepts patients under current medical care with childhood-onset (age ≤ 20 years old) SRNS, CNS, or persistent subnephrotic proteinuria with likely genetic disease.

The items collected by the registry are given in Supplemental Appendix. Retrospective and prospective information is collected. The regular data entries are discontinued when RRT is initiated, with a follow-up entry providing information about kidney transplantation and occurrence, treatment, and outcome of post-transplant disease recurrence.

For incident patients, a uniform set of diagnostic and therapeutic algorithms is proposed (www.podonet.org/opencms/opencms/podonet/podonet_en/Algorithms/index.html), but adherence to the protocols is not mandatory for inclusion of patients in the registry.

In the prospective part of the registry, serum and urine samples are collected at 6-month intervals and stored in a central biorepository. Also, DNA samples are collected from all available and consenting patients and any affected relatives. Central genetic screening has been offered for patients with lacking or incomplete screening results.

Definitions

SRNS was defined by persistent proteinuria after 4 weeks of oral steroid therapy (prednisone at 60 mg/m² per day). Subnephrotic proteinuria was defined as 24-hour protein excretion >100 mg but <1 g/m² per day, urine protein-to-creatinine ratio (UPCr) of 0.2 (0.6 if age <2 years old) to 2 mg/mg, or dipstick proteinuria of 1+. Nephrotic-range proteinuria was defined by 24-hour protein excretion ≥ 1 g/m² per day, UPCR>2 mg/mg, or dipstick $\geq 2+$. Complete remission of proteinuria was defined by the attainment of the following criteria during therapy: 24-hour protein excretion <100 mg/m² per d, UPCR<0.2 mg/mg, dipstick of 0 or (+), or in absence of proteinuria information, serum albumin >3.5 g/dl. Partial remission was defined by 24-hour protein excretion of 0.1–1 g/m²/d, UPCR=0.2–2 mg/mg, dipstick of 1+, and if available, serum

albumin >3.0 g/dl or if no proteinuria information was available, serum albumin of 3.0–3.5 g/dl.

Throughout the manuscript, percentages are given relative to all patients with available information regarding a particular topic.

Results

Cohort Structure

From August of 2009 to August of 2014, 1655 patients were registered who were ages 0–20 years old at disease onset. Among these, 261 (15.8%) patients were living in Italy, 227 (13.7%) patients were living in Germany, 226 (13.7%) patients were living in Poland, 189 (11.4%) patients were living in other European countries, 491 (29.7%) patients were living in Turkey or Georgia, 153 (9.2%) patients were living in the Middle East (Syria, Lebanon, Iran, and the United Arab Emirates), and 108 (6.5%) patients were living in Latin America (Chile and Colombia). Parental ethnicity (reported in 1274 families) was Caucasian in 1151 (90.3%) patients, Hispanic in 93 (7.3%) patients, Indian in nine (0.7%) patients, east Asian in five (0.4%) patients, African in three (0.2%) patients, Native American in two (0.2%) patients, and mixed in 11 (0.9%) patients. Among the patients with available respective information, parental consanguinity was reported in 28.6%, and familial disease occurrence was reported in 25.6%. Consanguinity and familial disease occurrence were most common in the Middle Eastern countries (Supplemental Table 1).

In 1544 patients, the age at first disease manifestation was reported (Figure 1). Among these, 207 (13.4%) were incident patients with first disease manifestation within 6 months before enrolment. Of the prevalent patients, 329 (24.6%) patients had been followed for 0.5–2 years, 345 (25.8%) patients had been followed for 2–5 years, 320 (23.9%) patients had been followed for 5–10 years, and 343 (25.6%) patients had been followed for >10 years at the time of enrolment.




Figure 1. | Age at first disease manifestation in children with and without an identified genetic cause of steroid-resistant nephrotic syndrome.

Information Content

Clinical and biochemical information from the time of first disease manifestation was available in 1310 (79.2%) patients. In total, 11,934 clinical follow-up entries were made for 1525 (92.1%) patients. Medication data were entered for 1283 (77.5%) patients and comprised 26,406 individual start, discontinuation, or dose change entries. Renal biopsy information was available from 1368 (82.7%) patients; the results of two sequential biopsies were recorded in 145 patients, and the results of more than two biopsies were recorded in 26 patients. Results of genetic screening were reported for 1174 (70.9%) patients, and information on family medical history was available from 1014 study participants (61.3%). The PodoNet biobank contains DNA samples from 950 patients, one or several serum samples from 472 patients, and urine samples from 450 patients.

First Disease Manifestation

The patient characteristics according to age at first disease manifestation are given in Table 1. Almost three quarters of the cohort presented between 1–11 years of age, whereas 6% were categorized as congenital, 7% were categorized as early infantile (onset age of 3–12 months), and 16% were categorized as adolescent-onset nephrotic syndrome (onset age \geq 12 years old). Among the children who underwent mutation screening, genetic disease detection rate was markedly higher (66%) in patients with CNS than in children first presenting at age 6 years old and older (16%). Regarding clinical and biochemical characteristics, hypoalbuminemia at time of diagnosis was more common in CNS, whereas hypertension was more prevalent in adolescents. Nephrotic-range proteinuria was not yet present in 8.7% of the children at the time of diagnosis, but most of these patients progressed to full nephrotic syndrome during follow-up.

Follow-Up Information

The median (interquartile range [IQR]) duration of follow-up from first manifestation was 3.7 (1.4–7.2) years. At the most recent observation, 61.3% of patients were still without RRT (CKD stage 1: 411 [60.2%], stage 2: 178 [26.1%], stage 3: 55 [8.1%], stage 4: 22 [3.2%], and stage 5: 17 [2.5%]), 11.7% of patients required dialysis, 14.2% of patients had received kidney allografts, 2.3% of patients were deceased, and 10.6% of patients were lost of follow-up. Among the patients without RRT, 33% were in full remission and 22.5% were in partial remission at last observation. Among the patients with full or partial remission at last observation, 66.9% had normal kidney function, 26.4% were in CKD stage 2, 5.1% in CKD stage 3, and 1.6% in CKD stages 4 or 5.

Histopathologic Findings

First kidney biopsies were performed in 1368 patients at a median (IQR) of 2 (1–8) months after first disease manifestation. The most common histopathologic diagnosis was FSGS (56.0%) followed by minimal change nephropathy (MCN; 21.1%), mesangioproliferative GN (MesPGN; 12.4%), and diffuse mesangial sclerosis (DMS; 2.9%). Information on average age, eGFR, and prevalence of hypertension at the time of diagnosis as well as prevalence of genetic disease according to histopathologic diagnosis is given in Table 2. Repeat biopsies were performed in 171 (12.5%)

Table 1. Key characteristics at time of diagnosis and response to intensified immunosuppressive therapy stratified by age at disease onset

Age at Disease Onset	No. of Patients (%)	Genetic Cause Identified N (%)	Nephrotic-Range Proteinuria (%)	Hypertension (%)	Serum Albumin (g/dl)	eGFR ^a (ml/min per 1.73 m ²)	Response to IIT		ESRD at Last Observation (%)	Post-Tx Recurrence N/N _{Tx} (%)
							Partial Remission N/N _{info} (%)	Complete Remission N/N _{info} (%)		
<3 mo	98 (6.3)	55 (66.3)	92.7	10.2	1.7 (0.9)	62 (41–80)	2/11 (18.2)	3/11 (27.3)	60.9	2/34 (5.9)
3–12 mo	106 (6.9)	31 (35.6)	89.7	17.0	2.1 (1.0)	89 (53–132)	3/33 (9.1)	16/33 (48.5)	37.8	0/21 (0.0)
1–5 yr	786 (50.9)	119 (21.8)	93.3	14.3	2.0 (0.9)	105 (78–147)	97/505 (19.2)	231/505 (45.7)	21.5	16/86 (18.6)
6–11 yr	350 (22.7)	40 (15.6)	88.1	16.6	2.6 (1.0)	105 (76–145)	49/208 (23.6)	73/208 (35.1)	27.2	11/36 (30.6)
\geq 12 yr	204 (13.2)	24 (16.1)	89.8	27.9	2.6 (1.0)	93 (64–129)	29/128 (22.7)	42/128 (32.8)	28.6	3/26 (11.5)

Data are given as number (%), mean (SD), or median (interquartile range). IIT, immunosuppressive therapy; N_{info}, number of patients with reported information; Tx, transplant; N_{Tx}, number of patients who received a kidney transplant.

^aMedian (interquartile range)

patients at a median (IQR) of 31 (17–54) months after the initial biopsy. The diagnosis changed in 54% of patients (in most cases from MCN to FSGS [26 of 47] and from MesPGN to FSGS [16 of 33]).

Extrarenal Symptoms

In 17.3% of the patients, one or several extrarenal abnormalities were reported (Table 3). Alterations of the morphology and/or function of the central nervous system (brain anomaly, microcephaly, and/or mental retardation) were the most common extrarenal abnormalities, affecting 5.3% of all patients. Other features included symptoms suggestive of *WT1* disease (sex reversal/urogenital abnormalities and cancer), impaired mitochondrial energy metabolism (myopathy, cardiomyopathy, and impaired hearing), Pierson syndrome (impaired vision), and Schimke syndrome (osteodysplasia).

Mutation Screening Results

In total, 3037 individual gene screens were performed in 1174 of 1655 study participants (70.9%). The extent of genetic screening varied among countries. Among 11 countries contributing at least 30 patients to the registry, screening prevalence was highest in Chile (91% of patients), Syria (84%), and Germany (80%) and lowest in Colombia (26%) and Iran (21%).

The most commonly screened genes were *NPHS2* and *WT1* (performed in 93% and 77% of screened patients, respectively), whereas other podocyte genes were screened more selectively guided by age, histopathology, and/or syndromic features or as part of joint research projects. In addition, 27 patients recently underwent comprehensive screening using a next generation sequencing panel of 31 podocyte genes. Genetic diagnoses were established in 277 (23.6%) patients; among these, mutations in *NPHS2* accounted for 49.8% of patients, mutations in *WT1* accounted for 17.3% of patients, mutations in *NPHS1* accounted for 14.8% of patients, mutations in *SMARCAL1* accounted for 4.3% of patients, and mutations in *PLCE1* accounted for 3.6% of patients. The remaining 10% of patients were attributable to variants in nine different genes (Table 4).

Pharmacologic Therapy

Among 1283 patients with available information on pharmacologic therapy, 42.6% received steroid therapy, 3.5% received cyclophosphamide pulse therapy, 21.2% received oral cyclophosphamide, 66.3% received ciclosporin A, 12.2% received tacrolimus, 24.9% received mycophenolate-mofetil (MMF), and 6.6% received rituximab at any time during the observation period. Renin angiotensin system (RAS) antagonists were applied in 76.1% of patients (angiotensin-converting enzyme inhibitor [72.3%] and/or angiotensin type 1 receptor blocker [33.7%]). In 1234 patients, 5240 drug treatment periods applied for at least 3 weeks (excluding intravenous therapies) were documented.

The distribution of treatment protocols, duration of administration, and treatment responses are summarized in Table 5. Calcineurin inhibitor (CNI)-based therapy showed the highest response rates, with approximately two thirds of patients achieving partial or complete proteinuria remission. Steroid pulse therapy and cyclophosphamid-based protocols

Table 2. Key characteristics according to histopathologic diagnosis sorted by patient age at time of biopsy

Histopathologic Diagnosis	No. of Patients N (%)	Genetic cause N/N _{info} (%)	At First Biopsy			IIT Responsiveness			ESRD at Last Observation (%)
			Age ^a (yr)	Serum Albumin (g/dl)	Hypertension (%)	eGFR ^a (ml/min per 1.73 m ²)	Partial Remission N/N _{info} (%)	Complete Remission N/N _{info} (%)	
CNS	8 (0.6)	3/7 (43.0)	0.2 (0.2–0.7)	1.4 (0.4)	0.0	56 (29–98)	0/1 (0.0)	0/1 (0.0)	57.1
DMS	39 (2.9)	22/35 (62.9)	1.3 (0.6–2.3)	2.0 (0.8)	26.3	59 (37–97)	0/10 (0.0)	1/10 (10.0)	72.2
MCN	283 (21.1)	24/195 (12.3)	4.2 (2.4–7.6)	2.5 (1.0)	15.8	108 (86–155)	41/202 (20.3)	103/202 (51.0)	12.9
MesPGN	166 (12.4)	23/123 (18.7)	4.4 (2.4–8.4)	2.4 (1.0)	15.1	118 (84–162)	26/101 (25.7)	41/101 (40.6)	18.9
FSGS	752 (56.0)	122/555 (22.0)	6.2 (3.0–11.0)	2.7 (1.0)	18.9	97 (68–142)	93/469 (19.8)	183/469 (39.0)	30.9
GGS	17 (1.3)	4/12 (33.3)	6.6 (3.4–9.1)	2.8 (0.8)	17.7	17 (10–85)	0/3 (0.0)	1/3 (33.3)	68.8
MN	11 (0.8)	0/11 (0.0)	7.0 (2.7–13.6)	2.4 (0.9)	27.3	116 (103–138)	3/10 (30.0)	5/10 (50.0)	9.1
MPGN	35 (2.6)	4/25 (16.0)	10.5 (5.8–11.9)	2.9 (0.8)	22.9	112 (61–148)	4/16 (25.0)	6/16 (37.5)	23.5
Other	31 (2.3)	9/27 (33.3)	8.8 (3.2–11.7)	2.7 (1.1)	6.7	45 (29–98)	4/16 (25.0)	6/16 (37.5)	24.1

Information on genetic disease, features at first biopsy, IIT responsiveness, and fraction of ESRD at last observations refers to diagnosis by initial biopsy. Data are given as number (%), mean (SD), or median (interquartile range). CNS, Finnish-type congenital nephrotic syndrome; DMS, diffuse mesangial sclerosis; MCN, minimal change nephropathy; MesPGN, mesangiolipiferative GN; GGS, global glomerulosclerosis; MN, membranous nephropathy; MPGN, membranoproliferative GN; N_{info}, number of patients with reported information.
^aMedian (interquartile range).

Table 3. Extrarenal abnormalities

Type of Extrarenal Abnormality	N	%
Patients without reported extrarenal abnormalities	1368	82.7
Mental retardation	65	3.9
Anomalies of central nervous system	42	2.5
Microcephaly	17	1.0
Visual impairment	32	1.9
Hearing disorder	25	1.5
Anomalies of peripheral nervous system	7	0.4
Myopathy	13	0.8
Cardiomyopathy	6	0.4
Urogenital abnormalities	33	2.0
Impaired sex differentiation	16	1.0
Short stature	84	5.1
Facial dysmorphism	37	2.2
Spondyloepiphyseal dysplasia	8	0.5
Polydactyly	5	0.3
Nail patella syndrome	2	0.1
Cardiac structural disorder	36	2.2
Malignant disorder	21	1.3
Hematologic disorders	7	0.4
Diabetes mellitus	7	0.4
Other endocrine abnormalities	23	1.4
Skin abnormalities	13	0.8
Abnormalities of the gastrointestinal tract	5	0.3
Autoimmune disorder	9	0.5
Connatal cytomegaly virus	9	0.5
Hepatitis B	6	0.4
Hepatitis C	10	0.6

generally showed poor efficacy and did not achieve better results than antiproteinuric treatment with RAS inhibitors, which were frequently coadministered with IIT.

The use of steroid therapy and nonsteroidal IIT increased with age at first manifestation but was not limited to patients beyond a certain threshold age. Steroid therapy (oral and/or intravenous pulses) was reportedly administered in 28.6% of the children with CNS and 59.4% of those with disease manifestation at age 3–12 months old; 24.5% of the children with CNS and 42.5% of those with onset at 3–12 months also received nonsteroidal IIT (CNI, MMF, cyclophosphamide, or rituximab). IIT response rates according to age at first manifestation are given in Table 1.

The initial histopathologic diagnosis did not predict IIT responsiveness in this SRNS cohort. Remission rates differed only marginally between MCN, MesPGN, and FSGS (Table 2).

Among 156 incident patients since the start of the registry in 2009 with disease onset at >3 months, oral steroid resistance, and documented follow-up pharmacotherapy, pulse steroids were initially administered to ascertain steroid resistance in 63 (40.4%) patients, and 97 (62.2%) patients received CNI therapy as first nonsteroidal therapy (39 with and 58 without prior steroid pulses) as recommended in the PodoNet consensus treatment algorithm; seven (4.5%) patients received other nonsteroidal IITs, 21 (13.5%) patients received RAS only, and seven (4.5%) patients were continued on oral steroids only.

In 115 (42%) of the patients in whom a genetic diagnosis was established at some time during follow-up, IIT protocols were applied; in 12 patients, complete proteinuria remission was observed during IIT.

Late Outcomes

Children with CNS and early infantile nephrotic syndrome were more likely to have ESRD at the last documented observation than older children (Table 1). With respect to histopathologic diagnosis, patients with DMS and global glomerulosclerosis had the highest likelihood of ESRD (Table 2). Patients with FSGS were more likely to progress than those with MCN.

In total, 217 patients underwent kidney transplantation. Proteinuria recurrence was reported in 33 (15.2%) of these patients. The risk of disease recurrence increased with age at first disease manifestation (Table 1).

The histopathologic diagnoses at last (native) kidney biopsy in patients with post-transplant disease recurrence were FSGS in 22 (68.8%) patients, MesPGN in four (12.5%) patients, global glomerulosclerosis in two (6.3%) patients, MCN in two (6.3%) patients, membranoproliferative GN in one (3.1%) patient, DMS in one (3.1%) patient, and unknown in one patient.

Although post-transplant disease recurrence occurred in 28.5% of patients without a genetic diagnosis, recurrence was noted in four (4.5%) patients of those diagnosed with genetic disease. In all of these patients, *NPHS2* mutations had been identified.

Discussion

In this work, we present essential reference information for pediatric SRNS. With >1600 patients enrolled in 67 centers on three continents, the PodoNet Registry cohort represents, by far, the largest collection of clinicopathologic and genetic information assembled to date.

The registry provides important information regarding the distribution of clinical and histopathologic phenotypes in childhood SRNS. We also compiled valuable information about extrarenal disease manifestations. Neurologic abnormalities were most commonly reported followed by a wide range of anomalies concerning various organs and tissues. Altogether, syndromic phenotypes accounted only for a small fraction of patients, and SRNS occurred as an isolated kidney disease in 83% of the patients in the cohort.

Historically, renal histopathology has been used as a key criterion for diagnostic and prognostic categorization in children with SRNS. Here, we provide demographic, clinical, and genetic information on the individual histopathologic entities in the pediatric SRNS population. The three major histologic diagnoses (FSGS, MCN, and MesPGN) presented with similar degrees of hypoalbuminemia and comparable prevalence of hypertension. Patients with FSGS were slightly older and more often had reduced eGFR at the time of diagnosis.

FSGS was observed almost two times as often as MCN and MesPGN together. The predominance of FSGS contrasts slightly with previous single-center surveys in south Asian and African populations, where MCN was equally or even more common than FSGS (6–8). This variation might be related to differences in ethnic composition, biopsy indication

Table 4. Results of genetic screening studies

Gene	Patients Screened	Patients with Causative Mutation	Screened Positive (%)
<i>NPHS2</i>	1088	138	12.7
<i>WT1</i>	902	48	5.3
<i>NPHS1</i>	208	41	19.7
<i>SMARCAL1</i>	68	12	17.6
<i>PLCE1</i>	75	10	13.3
<i>PTPRO</i>	45	6	13.3
<i>LAMB2</i>	84	5	6.0
<i>INF2</i>	112	4	3.6
<i>COQ6</i>	30	3	10.0
<i>MYO1E</i>	48	2	4.2
<i>TRPC6</i>	96	1	1.0
<i>COQ2</i>	56	1	1.8
<i>LMX1B</i>	27	1	3.7
<i>ADCK4</i>	27	1	3.7
<i>PDSS2</i>	56	0	0.0
<i>ACTN4</i>	59	0	0.0
<i>CD2AP</i>	56	0	0.0
All	1174	277	23.6

policies, or periods of observation analyzed. A trend toward an increasing incidence of FSGS has been noted in several studies (9–11).

The other, much rarer histopathologic entities DMS, membranoproliferative GN, and membranous nephropathy typically presented with milder hypoalbuminemia and a slightly higher prevalence of hypertension than MCN, MesPGN, and FSGS. Patients with DMS usually manifested at earlier age with established renal failure.

Abnormalities in podocyte genes are considered likely in children with multidrug-resistant nephrotic syndrome (12,13). In 21 countries participating in PodoNet, genetic screening was performed in more than two thirds of the affected children. Of these, almost one quarter of the patients tested were screened positive for one of the monogenic disorders known to cause SRNS. In agreement with published numbers (14), the detection rate was highest in children with CNS and significantly lower with increasing age at first disease manifestation during the first 6 years of life. However, even among patients first presenting at adolescent age, a genetic disease cause could be established in more than one of seven patients. A detailed analysis of the genetic findings in the adolescent segment of the cohort has recently been published (15). The observed gene detection rate is most likely an underestimate, because until recently, the high cost and processing times of conventional Sanger sequencing allowed for screening of only a few genes selected according to age at disease onset, histopathology findings, and in familial cases, mode of inheritance. Also, within the lifetime of the registry, several new genes have been identified (5)—some as part of the activities of the PodoNet consortium (16,17). Our findings indicate that SRNS may be explained by rare abnormalities in a large number of genes. The advent of next generation sequencing will soon allow us to routinely screen all genes associated with SRNS. The detection rates per gene and per patient published in this report, which were obtained mainly

by classic screening strategies and conventional technologies, may serve as a benchmark for future efforts to establish genetic diagnoses in patients with SRNS.

Another noteworthy aspect of this study is the association of genetic disease causes with histopathologic diagnoses. Our findings underline the limited value of kidney biopsies in distinguishing genetic from nongenetic disease etiologies. Genetic abnormalities were found in 22% of patients with FSGS, 19% of patients with MesPGN, and 12% of patients with MCN. Close associations with specific genetic disorders were limited to DMS (*WT1* and *PLCE1* nephropathies) and CNS (*NPHS1* disease).

Detailed information on pharmacotherapy is available from >1200 children in the registry. The wide range of reported monotherapies and combined therapies reflects the complexity of SRNS management with frequent medication changes because of lack of efficiency or—observed or anticipated—adverse effects. Moreover, significant variation was noted regarding first- and second-line drug choices across the participating centers and countries, and treatment preferences seemed to change with time. Among the patients diagnosed with SRNS since the start of the PodoNet Registry, 40% adhered to the internal consensus recommendation to ascertain steroid resistance by intravenous steroid pulse therapy, and 62% used CNI as first-choice second-line IIT. Remarkably, steroid therapy and even IIT protocols were applied in some 25% of children presenting with CNS. These insights illustrate the large current variability of global clinical practice patterns.

Our preliminary analysis of proteinuria response patterns with different IIT protocols allows several tentative conclusions. First, steroid pulses and cyclophosphamide are of very limited, if any, efficacy in children with SRNS. Second, 40%–50% of patients with SRNS seem to respond to CNI by achieving complete remission of proteinuria. Third, rituximab may be an equally effective therapeutic option as CNI. Fourth, MMF may be less efficacious than CNI and rituximab in SRNS. Fifth, nonspecific antiproteinuric therapy by RAS inhibition is associated with complete remission of proteinuria in some 25% and partial remission in another 20% of patients with SRNS, an effect to be aware of in the evaluation of treatment effects and account for in the design of future pharmacotherapeutic trials in this condition.

These main findings are well in line with the existing body of evidence derived from observational studies and clinical trials (18–26) and provide a valuable reference framework for future clinical research. More detailed analyses of intervention-response patterns, relapse rates during and after discontinuation of treatment, adverse effect patterns, etc. are beyond the scope of this report but will be performed.

Whereas proteinuria was multidrug resistant in the vast majority of genetic cases, a few patients with genetic diagnoses apparently exhibited responsiveness to IIT. Although beyond the scope of this study, these peculiar cases deserve a detailed analysis of the nature of the genetic abnormalities diagnosed and the pharmacotherapies applied.

A diagnostically and prognostically relevant feature of SRNS is post-transplant disease recurrence, which was reported in 15% of >200 allograft recipients in the registry. Proteinuria recurrence occurred almost exclusively in patients without a genetic diagnosis; four patients with apparent recurrence despite genetic disease likewise will require additional evaluation.

Table 5. Medication protocols applied in 1234 patients

Medication Protocol	No. of Patients	No. of Periods	Cotreated with RAS Antagonists (%)	Cotreated with Oral Steroids (%)	Treatment Duration (mo)	Response to Therapy	
						Partial Remission N/N _{info} (%)	Complete Remission N/N _{info} (%)
RAS antagonists only	623	950	—	63.3	5.2 (1.9–17.1)	109/531 (20.5)	141/531 (26.6)
Oral steroids only	722	1119	—	—	2.0 (1.4–4.7)	53/567 (9.3)	62/567 (10.9)
Steroid pulses	541	1114	51.0	72.4	0.1 (0.1–0.9)	42/279 (15.1)	25/279 (9.0)
CPH (oral or pulse)	273	336	39.0	92.0	2.5 (1.6–3.0)	11/131 (8.4)	22/131 (16.8)
CNI	806	1210	70.7	89.5	8.3 (3.0–23.0)	153/707 (21.6)	311/707 (44.0)
MMF	177	219	80.4	71.7	6.7 (3.0–14.9)	19/148 (12.8)	57/148 (38.5)
CNI + MMF	137	167	81.4	76.0	7.7 (3.0–23.9)	26/119 (21.8)	57/119 (47.9)
Rituximab	79	91	87.9	76.9	0.0 (0.0–4.6)	10/66 (15.2)	29/66 (43.9)

Treatment duration is given as median and interquartile range. Renin angiotensin system (RAS) antagonists are angiotensin-converting enzyme inhibitor and/or angiotensin type 1 receptor blocker. CPH, cyclophosphamide; CNI, calcineurin inhibitor (cyclosporin A or tacrolimus); MMF, mycophenolate-mofetil.

Finally, our study underlines the clinical usefulness of genetic testing in children with SRNS. Establishing a genetic diagnosis is far superior to histopathologic disease classification in predicting IIT responsiveness and post-transplant disease recurrence in patients with SRNS. However, because hereditary forms of SRNS encompass a large number of genes, it will take the routine use of the emerging next-generation sequencing technologies for simultaneous assessment of all known podocytopathy genes to materialize the full potential of genetic screening.

Acknowledgments

We acknowledge the continued support of patient families and medical staff at all participating centers.

The PodoNet project has received financial support from E-Rare (German Ministry of Education and Research), European Union 7th Framework Programme (EUReOmics) Grant 2012-305608, Chilean Research Agency (Fondecyt) Grant 11090045, Polish Ministry of Science and Education Grant N402631840, and German Research Foundation Grant Scha 477/11-1. F.O. was supported by Scientific and Technological Research Council of Turkey Grant 108S417 and Hacettepe University Infrastructure Project Grants 06A101008 and 011A101003.

PodoNet Collaborators: Chile: Marta Azocar, Hospital Luis Calvo Mackenna-Facultad de Chile, Santiago; Lily Quiroz, Roberto del Rio Childrens Hospital, Santiago; Colombia: Lina Maria Serna Higuaita, Hospital Pablo Toban Uriba, Medellín; Czech Republic: Jiří Dušek, University Hospital Motol, Prague; France: Bruno Ranchin, Centre de Reference des Maladies Renales Rare/University Hospital, Lyon; Michel Fischbach, University Children's Hospital, Strasbourg; Georgia: Tinatin Davitaia, M. Iashvili Children's Central Hospital, Tbilisi; Germany: Jutta Gellermann, Charité University Hospital, Berlin; Jun Oh, University Children's Hospital, Hamburg; Anette Melk, Medical University Children's Hospital, Hannover; Franz Schaefer, Heidelberg University Center for Pediatrics and Adolescent Medicine, Heidelberg; Marianne Wigger, University Children's Hospital, Rostock; Greece: Fotios Papachristou, Hippokraton General Hospital, Aristotle University, Thessaloniki; Hungary: Peter Sallay, Semmelweis University Hospital, Budapest; Iran: Alaleh Gheissari, St. Al Zahra Hospital, Isfahan University of Medical Science, Isfahan; Italy: Giuseppe Remuzzi, Clinical Research Centre of Rare Diseases "Aldo & Cele Daccò," Istituto di Ricerche Farmacologiche "Mario Negri," Bergamo; Andrea Pasini, S.Orsola-Malpighi Hospital, Bologna; Gian Marco Ghiggeri, Istituto G. Gaslini, Genova; Gianluigi Ardissino, IRCCS Fondazione Ospedale Maggiore Policlinico, Milano; Elisa Benetti, Ascenda Ospedaliera University, Padua; Francesco Emma, Ospedale Pediatrico Bambino, Rome; Lebanon: Bilal Aoun, Rafic Hariri University Hospital, Beirut; Pauline Abou-Jaoudé, Notre Dame de Secours University Hospital, Byblos; Lithuania: Augustina Jankauskiene, Vilnius University Children's Hospital, Vilnius; Poland: Anna Wasilewska, University Hospital, Bialystok; Lidia Hyla-Klekot, Pediatrics and Oncology Center, Chorzow; Aleksandra Zurowska, Medical University, Gdansk; Dorota Drozd, Jagiellonian University Medical College, Krakow; Marcin Tkaczyk, Polish Mothers Memorial Hospital Research Institute, Lodz; Halina Borzecka, Medical University, Lublin; Magdalena Silska, Poznan University of Medical Sciences, Poznan; Tomasz Jarmolinski, Pommeranian Academy of Medicine, Szczecin; Agnieszka Firszt-Adamczyk, Wojewodzki Hospital, Torun; Joanna Ksiazek, Centrum Zdrowia Dziecka, Warsaw; Elzbieta Kuzma-Mroczkowska, University Hospital, Warsaw; Anna Medynska, Medical University, Wroclaw; Maria Szczepanska, Medical University of Silesia in Katowice, Zabrze; Portugal: Alberto Caldas Afonso, Hospital

S.Joao Pediatrics, Porto; Helena Jardim, Centre Hospitalar, Porto; Serbia: Amira Peco-Antic, University Children's Hospital, Belgrade; Radovan Bogdanovic, Institute of Mother and Healthcare of Serbia, Belgrade; Sweden: Rafael T. Krmar, Karolinska University Children's Hospital, Stockholm; Switzerland: Giacomo D. Simonetti, University Children's Hospital, Bern; Syria: Bassam Saeed, Kidney Hospital, Damascus; Turkey: Ali Anarat, Cukurova University Children's Hospital, Adana; Aysin Bakkaloglu, Hacettepe University Medical Faculty, Ankara; Ayse Balat, Gaziantep University Medical Faculty, Gaziantep; Z. Esra Baskin, Baskent University Hospital, Ankara; Nilgun Cakar, Diskapi Children's Hospital, Ankara; Ozlem Erdogan, Sami Ulus Children's Hospital, Ankara; Birsin Özcakar, Ankara University Medical School, Ankara; Fatih Ozaltin, Hacettepe University, Ankara; Onur Sakallioğlu, Gulhane Military Academy of Medicine, Ankara; Oguz Soylemezoglu, Gazi University Hospital, Ankara; Sema Akman, Akdeniz University, Antalya; Faysal Gok, Gulhane Military Academy of Medicine, Gulhane; Salim Caliskan, Cerrahpasa Medical Faculty, Istanbul University, Istanbul; Cengiz Candan, Goztepe Hospital, Istanbul; Sevinc Emre, Istanbul Medical Faculty, Istanbul; Sevgi Mir, Ege University Medical Faculty, Izmir; Ipek Akil, Celal Bayar University, Manisa; Pelin Ertan, Celal Bayar University, Manisa; Ozan Özkaya, Ondokuz Mayıs University, Samsun; Mukaddes Kalyoncu, Karadeniz University, Trabzon; United Arab Emirates: Eva Simkova, Dubai Hospital, Dubai; Entesar Alhammadi, Al Qassimi Hospital, Sharjah; Ukraine: Roman Sobko, Western Ukrainian Specialised Medical Centre, Lviv.

Disclosures

None.

References

- McKinney PA, Feltbower RG, Brocklebank JT, Fitzpatrick MM: Time trends and ethnic patterns of childhood nephrotic syndrome in Yorkshire, UK. *Pediatr Nephrol* 16: 1040–1044, 2001
- Kim JS, Bellew CA, Silverstein DM, Aviles DH, Boineau FG, Vehaskari VM: High incidence of initial and late steroid resistance in childhood nephrotic syndrome. *Kidney Int* 68: 1275–1281, 2005
- Mekahli D, Liutkus A, Ranchin B, Yu A, Bessenay L, Girardin E, Van Damme-Lombaerts R, Palcoux JB, Cachat F, Lavocat MP, Bourdat-Michel G, Nobili F, Cochat P: Long-term outcome of idiopathic steroid-resistant nephrotic syndrome: A multicenter study. *Pediatr Nephrol* 24: 1525–1532, 2009
- Zagury A, Oliveira AL, Montalvão JA, Novaes RH, Sá VM, Moraes CA, Tavares MS: Steroid-resistant idiopathic nephrotic syndrome in children: Long-term follow-up and risk factors for end-stage renal disease. *J Bras Neurol* 35: 191–199, 2013
- Saleem MA: New developments in steroid-resistant nephrotic syndrome. *Pediatr Nephrol* 28: 699–709, 2013
- Mubarak M, Lanewala A, Kazi JI, Akhter F, Sher A, Fayyaz A, Bhatti S: Histopathological spectrum of childhood nephrotic syndrome in Pakistan. *Clin Exp Nephrol* 13: 589–593, 2009
- Nammalwar BR, Vijayakumar M, Prahlad N: Experience of renal biopsy in children with nephrotic syndrome. *Pediatr Nephrol* 21: 286–288, 2006
- Bhimma R, Adhikari M, Asharam K: Steroid-resistant nephrotic syndrome: The influence of race on cyclophosphamide sensitivity. *Pediatr Nephrol* 21: 1847–1853, 2006
- Srivastava T, Simon SD, Alon US: High incidence of focal segmental glomerulosclerosis in nephrotic syndrome of childhood. *Pediatr Nephrol* 13: 13–18, 1999
- Bonilla-Felix M, Parra C, Dajani T, Ferris M, Swinford RD, Portman RJ, Verani R: Changing patterns in the histopathology of idiopathic nephrotic syndrome in children. *Kidney Int* 55: 1885–1890, 1999
- Banaszak B, Banaszak P: The increasing incidence of initial steroid resistance in childhood nephrotic syndrome. *Pediatr Nephrol* 27: 927–932, 2012
- Ruf RG, Lichtenberger A, Karle SM, Haas JP, Anacleto FE, Schultheiss M, Zalewski I, Imm A, Ruf EM, Mucha B, Bagga A, Neuhaus T, Fuchshuber A, Bakkaloglu A, Hildebrandt F; Arbeitsgemeinschaft Für Pädiatrische Nephrologie Study Group: Patients with mutations in NPHS2 (podocin) do not respond to standard steroid treatment of nephrotic syndrome. *J Am Soc Nephrol* 15: 722–732, 2004
- Büscher AK, Kranz B, Büscher R, Hildebrandt F, Dworniczak B, Pennekamp P, Kuwertz-Bröking E, Wingen AM, John U, Kemper M, Monnens L, Hoyer PF, Weber S, Konrad M: Immunosuppression and renal outcome in congenital and pediatric steroid-resistant nephrotic syndrome. *Clin J Am Soc Nephrol* 5: 2075–2084, 2010
- Hinkes BG, Mucha B, Vlangos CN, Gbadegesin R, Liu J, Hasselbacher K, Hangan D, Ozaltin F, Zenker M, Hildebrandt F; Arbeitsgemeinschaft für Paediatrische Nephrologie Study Group: Nephrotic syndrome in the first year of life: Two thirds of cases are caused by mutations in 4 genes (NPHS1, NPHS2, WT1, and LAMB2). *Pediatrics* 119: e907–e919, 2007
- Lipska BS, Iatropoulos P, Maranta R, Caridi G, Ozaltin F, Anarat A, Balat A, Gellermann J, Trautmann A, Erdogan O, Saeed B, Emre S, Bogdanovic R, Azocar M, Balasz-Chmielewska I, Benetti E, Caliskan S, Mir S, Melk A, Ertan P, Baskin E, Jardim H, Davitaia T, Wasilewska A, Drodz D, Szczepanska M, Jankauskiene A, Higueta LM, Ardissino G, Ozkaya O, Kuzma-Mroczkowska E, Soylemezoglu O, Ranchin B, Medynska A, Tkaczyk M, Peco-Antic A, Akil I, Jarmolinski T, Firszt-Adamczyk A, Dusek J, Simonetti GD, Gok F, Gheissari A, Emma F, Krmar RT, Fischbach M, Printza N, Simkova E, Mele C, Ghiggeri GM, Schaefer F; PodoNet Consortium: Genetic screening in adolescents with steroid-resistant nephrotic syndrome. *Kidney Int* 84: 206–213, 2013
- Mele C, Iatropoulos P, Donadelli R, Calabria A, Maranta R, Cassis P, Buelli S, Tomasoni S, Piras R, Krendel M, Bettoni S, Morigi M, Delledonne M, Pecoraro C, Abbate I, Capobianchi MR, Hildebrandt F, Otto E, Schaefer F, Macchiardi F, Ozaltin F, Emre S, Ibsirlioglu T, Benigni A, Remuzzi G, Noris M; PodoNet Consortium: MYO1E mutations and childhood familial focal segmental glomerulosclerosis. *N Engl J Med* 365: 295–306, 2011
- Ozaltin F, Ibsirlioglu T, Taskiran EZ, Baydar DE, Kaymaz F, Buyukcelik M, Kilic BD, Balat A, Iatropoulos P, Asan E, Akarsu NA, Schaefer F, Yilmaz E, Bakkaloglu A; PodoNet Consortium: Disruption of PTPRO causes childhood-onset nephrotic syndrome. *Am J Hum Genet* 89: 139–147, 2011
- Hodson EM, Willis NS, Craig JC: Interventions for idiopathic steroid-resistant nephrotic syndrome in children. *Cochrane Database Syst Rev* 11: CD003594, 2010
- Lombel RM, Gipson DS, Hodson EM; Kidney Disease: Improving Global Outcomes: Treatment of steroid-sensitive nephrotic syndrome: New guidelines from KDIGO. *Pediatr Nephrol* 28: 415–426, 2013
- Ehrich JH, Geerlings C, Zivicnjak M, Franke D, Geerlings H, Gellermann J: Steroid-resistant idiopathic childhood nephrosis: Overdiagnosed and undertreated. *Nephrol Dial Transplant* 22: 2183–2193, 2007
- Gellermann J, Ehrich JH, Querfeld U: Sequential maintenance therapy with cyclosporin A and mycophenolate mofetil for sustained remission of childhood steroid-resistant nephrotic syndrome. *Nephrol Dial Transplant* 27: 1970–1978, 2012
- Gulati A, Sinha A, Gupta A, Kanitkar M, Sreenivas V, Sharma J, Mantan M, Agarwal I, Dinda AK, Hari P, Bagga A: Treatment with tacrolimus and prednisolone is preferable to intravenous cyclophosphamide as the initial therapy for children with steroid-resistant nephrotic syndrome. *Kidney Int* 82: 1130–1135, 2012
- Gulati A, Sinha A, Jordan SC, Hari P, Dinda AK, Sharma S, Srivastava RN, Moudgil A, Bagga A: Efficacy and safety of treatment with rituximab for difficult steroid-resistant and -dependent nephrotic syndrome: Multicentric report. *Clin J Am Soc Nephrol* 5: 2207–2212, 2010
- Hamasaki Y, Yoshikawa N, Nakazato H, Sasaki S, Iijima K, Nakanishi K, Matsuyama T, Ishikura K, Ito S, Kaneko T, Honda M; for Japanese Study Group of Renal Disease in Children: Prospective 5-year follow-up of cyclosporine treatment in

- children with steroid-resistant nephrosis. *Pediatr Nephrol* 28: 765–771, 2013
25. Roberti I, Vyas S: Long-term outcome of children with steroid-resistant nephrotic syndrome treated with tacrolimus. *Pediatr Nephrol* 25: 1117–1124, 2010
 26. Aizawa-Yashiro T, Tsuruga K, Watanabe S, Oki E, Ito E, Tanaka H: Novel multidrug therapy for children with cyclosporine-resistant or -intolerant nephrotic syndrome. *Pediatr Nephrol* 26: 1255–1261, 2011

Received: June 23, 2014 **Accepted:** December 22, 2014

Published online ahead of print. Publication date available at www.cjasn.org.

This article contains supplemental material online at <http://cjasn.asnjournals.org/lookup/suppl/doi:10.2215/CJN.06260614/-/DCSupplemental>.

Online Appendix to:

Trautmann et al.

**Spectrum of Steroid-Resistant and Congenital Nephrotic Syndrome in Children:
The PodoNet Registry Cohort**

Information collected in PodoNet Registry

Information on ethnic background and parental consanguinity, any relevant family history, age, clinical symptoms and biochemical features at first disease manifestation, initial and any subsequent renal histopathological findings, extrarenal abnormalities, and the results of genetic screening are recorded. In addition, clinical, biochemistry and medication updates are obtained every 6 months, with additional entries requested at time of major treatment changes. The following information is entered: date, height, weight, a semiquantitative assessment of edema status, serum albumin and creatinine, CsA/tacrolimus blood levels, proteinuria (order of preference: 24-hour protein excretion, spot urine protein/creatinine ratio, semiquantitative urine dipstick), type, dose and date of start, discontinuation and dose change of immunosuppressive medications and RAS inhibitors, and treatment complications including weight gain, hypertension, bone disease, infections, mood disorders, cataract, leukopenia, anemia, hair loss, hypertrichosis, gingival hyperplasia, and diabetes mellitus.

Table S1. Distribution of countries of origin with rates of parental consanguinity, familial disease occurrence and genetic diagnosis. Countries contributing less than 10 patients are summarized as 'Other European' or 'Other Middle East'.

N_{info} = Number of patients with available information on given item.

Country	N	Parental consanguinity N/ N_{info} (%)	Familial disease N/ N_{info} (%)	Identified genetic disease N/ N_{info} (%)
Georgia	14	0/13 (0.0)	0/10 (0.0)	1/11 (9.1)
Greece	12	0/5 (0.0)	0/11 (0.0)	0/7 (0.0)
Sweden	12	0/12 (0.0)	1/5 (20.0)	0/8 (0.0)
Other European	10	0/9 (0.0)	2/10 (20.0)	2/9 (22.2)
Lithuania	31	0/20 (0.0)	5/23 (21.7)	6/11 (54.5)
Serbia	59	1/47 (2.1)	6/47 (12.7)	6/37 (16.2)
Colombia	43	1/37 (2.7)	4/27 (14.8)	1/11 (9.1)
Italy	261	4/141 (2.8)	30/148 (20.3)	38/189 (20.1)
Poland	226	3/75 (4.0)	46/138 (33.3)	34/175 (19.4)
Chile	65	2/49 (4.1)	6/39 (15.4)	13/59 (22.0)
Portugal	21	2/19 (10.5)	3/18 (16.7)	4/15 (26.7)
France	32	3/28 (10.7)	0/12 (0.0)	15/23 (65.2)
Czech Republic	12	1/8 (12.5)	0/12 (0.0)	2/11 (18.2)
Germany	227	11/47 (23.4)	20/88 (22.7)	58/181 (32.0)
Other Middle East	12	3/10 (30.0)	2/8 (25.0)	3/8 (37.5)
Turkey	477	193/415 (46.5)	109/351 (31.1)	81/351 (23.1)
Syria	61	37/61 (60.7)	23/60 (38.3)	11/51 (21.6)
Iran	80	45/73 (61.6)	3/6 (50.0)	2/17 (11.8)
All	1,655	306/1,070 (28.6)	260/1,014 (25.6)	277/1,174 (23.6)

Spectrum of Steroid Resistant and Congenital Nephrotic Syndrome in Children: The PodoNet Registry Cohort

Agnes Trautmann¹, Monica Bodria², Fatih Ozaltin³, Alaleh Gheisari⁴, Anette Melk⁵, Marta Azocar⁶, Ali Anarat⁷, Salim Caliskan⁸, Francesco Emma⁹, Jutta Gellermann¹⁰, Jun Oh¹¹, Esra Baskin¹², Joanna Ksiazek¹³, Giuseppe Remuzzi¹⁴, Ozlem Erdogan¹⁵, Sema Akman¹⁶, Jiri Dusek¹⁷, Tinatin Davitaia¹⁸, Ozan Özkaya¹⁹, Fotios Papachristou²⁰, Agnieszka Firszt-Adamczyk²¹, Tomasz Urasinski²², Sara Testa²³, Rafael T. Krmar²⁴, Lidia Hyla-Klekot²⁵, Andrea Pasini²⁶, Z. Birsin Özcahar²⁷, Peter Sallay²⁸, Nilgun Cakar²⁹, Monica Galanti³⁰, Joelle Terzic³¹, Bilal Aoun³², Alberto Caldas Afonso³³, Hanna Szymanik-Grzelak³⁴, Beata S. Lipska³⁵, Sven Schnaidt³⁶, Franz Schaefer¹, for the PodoNet Consortium*

¹ Division of Pediatric Nephrology, University Center for Pediatrics and Adolescent Medicine, Heidelberg, Germany

² Division of Nephrology, Dialysis and Transplantation, Istituto Giannina Gaslini, Genoa, and Department of Clinical Medicine, Nephrology and Health Sciences, University of Parma, Italy

³ Department of Pediatric Nephrology, Nephrogenetics Laboratory and Center for Biobanking and Genomics, Hacettepe University, Ankara, Turkey

⁴ Pediatric Nephrology Department, Isfahan University of Medical Science, St. Al Zahra Hospital, Isfahan, Iran

⁵ Department of Pediatric Nephrology, Medical University Children's Hospital, Hannover, Germany

⁶ Department of Pediatric Nephrology, Hospital Luis Calvo Mackenna-Facultad de Chile, Santiago de Chile

⁷ Department of Pediatric Nephrology, Cukurova University Medical Faculty, Adana, Turkey

⁸ Department of Pediatric Nephrology, Cerrahpasa Medical Faculty, Istanbul University, Istanbul, Turkey

⁹ Nephrology and Dialysis Unit, Ospedale Pediatrico Bambino, Rome, Italy

¹⁰ Department of Pediatric Nephrology, Charite University Hospital, Berlin, Germany

¹¹ Department of Pediatric Nephrology, University Children's Hospital, Hamburg, Germany

¹² Department of Pediatric Nephrology, Baskent University Hospital, Ankara, Turkey

¹³ Department of Pediatric Nephrology, Centrum Zdrowia Dziecka, Warsaw, Poland

¹⁴ Clinical Research Centre for Rare Diseases "Aldo & Cele Daccò", IRCCS - Istituto di Ricerche Farmacologiche "Mario Negri" and Unit of Nephrology, Azienda Ospedaliera Papa Giovanni XXXI, Bergamo, Italy

¹⁵ Department of Pediatric Nephrology, Dr. Sami Ulus Children's Hospital, Ankara, Turkey

¹⁶ Department of Pediatric Nephrology, Akdeniz University, Antalya, Turkey

¹⁷ 1st Department of Pediatrics, University Hospital Motol, Prague, Czech Republic

¹⁸ Department of Pediatrics, M.Iashvili Children's Central Hospital, Tbilisi, Georgia

¹⁹ Department of Pediatric Nephrology, Ondokuz Mayıs University, Samsun, Turkey

²⁰ 1st Pediatric Department, Hippokraton General Hospital, Aristotle University, Thessaloniki, Greece

²¹ Department of Pediatric Nephrology, Wojewodzki Hospital, Torun, Poland

²² Clinic of Pediatrics, Pomeranian Academy of Medicine, Szczecin, Poland

²³ Pediatric Nephrology and Dialysis, Fondazione IRCCS Ca'Granda, Ospedale Maggiore Policlinico, Milano, Italy

²⁴ Division of Pediatrics, Department of Clinical Science, Intervention and Technology, Karolinska University Hospital, Stockholm, Sweden

²⁵ Department of Pediatric Nephrology, Pediatrics and Oncology Center, Chorzow, Poland

²⁶ Pediatric Nephrology and Dialysis Unit, S.Orsola-Malpighi Hospital, Bologna, Italy

²⁷ Department of Pediatric Nephrology, Ankara University Medical School, Ankara, Turkey

²⁸ 1st Department of Pediatrics, Semmelweis University, Budapest, Hungary

²⁹ Department of Pediatric Nephrology, Diskapi Children's Hospital, Ankara, Turkey

³⁰ Pediatric Nephrology Unit, Roberto del Rio Children's Hospital, Santiago de Chile

³¹ Pediatric Nephrology and Dialysis Unit, University Children's Hospital, Strasbourg

³² Pediatric Nephrology Unit, Rafic Hariri University Hospital, Beirut, Lebanon

³³ Department of Pediatric Nephrology, Hospital S. Joao Pediatrics, Porto, Portugal

³⁴ Department of Pediatric Nephrology, University of Warsaw, Warsaw, Poland

³⁵ Department of Biology and Genetics, Medical University of Gdansk, Gdansk, Poland

³⁶ Institute of Medical Biometry and Informatics, University of Heidelberg, Germany