

The Provider's Role in Conservative Care and Advance Care Planning for Patients with ESRD

Gregorio T. Obrador

Clin J Am Soc Nephrol 11: 750–752, 2016. doi: 10.2215/CJN.03150316

Conservative care (CC), also known as conservative management, is being gradually recognized as a viable therapeutic alternative for patients with advanced CKD in the United States (1,2). The Renal Physician Association's clinical practice guideline on shared decision making in the appropriate initiation of and withdrawal from dialysis recommends to inform patients with stage 4 or 5 CKD and patients with ESRD about their prognosis and all treatment options, including CC (3). Although evidence is limited, current data suggest that patients with ESRD ≥ 75 years of age with high levels of comorbid conditions and/or poor functional status may not benefit from dialysis in survival and/or health-related quality of life. Indeed, hospitalization rates decrease and home deaths increase when these patients receive comprehensive CC compared with dialysis (4–7).

Despite its potential benefits, several caveats limit implementation of CC, including (1) lack of a precise definition, (2) poor characterization of the CKD population that might benefit from it, (3) limited observational evidence of its potential benefits, and (4) scarce data on patient-centered outcomes (8). A recent survey regarding practice patterns of CC in the United Kingdom highlighted some of these limitations (9). Despite that all but one of 67 (of a total of 71) adult renal units reported provision of CC, the absolute number of patients could not be calculated because of lack of agreement on when a patient is receiving CC. Indeed, terminology varied substantially among renal units, with conservative management being the most frequently used term (46%); 80% of units reported a need for better evidence comparing outcomes of CC versus dialysis, and 65% considered it appropriate to enter patients into a randomized trial. Interestingly, the majority of renal units (88%) provided primary care physicians (PCPs) with information/advice regarding CC.

Advance care planning (ACP) is essential to decision making regarding therapeutic options for patients with advanced CKD, including CC. ACP involves communicating information to the patient and family (or other caregiver) about the current clinical condition, prognosis, and treatment options within the context of the patient's values, goals, and preferences, which will ultimately guide shared decision making (3). The purpose of the latter is to align the treatment with patient and

family goals, values, and preferences. The team caring for patients with advanced CKD should become involved in ACP early in the illness and in a flexible manner, because the health status and patient and family goals may change over time (8). Unfortunately, current evidence suggests that discussions about prognosis and end of life care are uncommon and that decision making is integrated poorly into CKD care, which leads many patients and their families to be ill prepared for living with CKD or dealing with end of life decisions (10,11). Although still limited, current evidence suggests that ACP is associated with improved end of life care, reduced hospitalizations and inappropriate use of life-sustaining treatments, increased use of hospice and supportive care, and greater compliance with patients' end of life wishes (12,13).

In this issue of the *Clinical Journal of the American Society of Nephrology*, the results of two studies examining the provider's role in CC and ACP for patients with advanced CKD are reported (14,14).

In the first study, Parvez *et al.* (14) conducted a nationwide survey to examine nephrologists' and PCPs' practices, attitudes, and knowledge regarding CC in the United States.

Parvez *et al.* (14) reported confusion about terminology (*i.e.*, >40% of both nephrologists and PCPs believed that CC and palliative care were the same). Also, respondents had an underappreciation that pain is a common symptom (*i.e.*, <40% of both nephrologists and PCPs endorsed pain management), and there was a lack of knowledge about CC (*i.e.*, 20%–30% of nephrologists and PCPs responded that CC could serve as a bridge to kidney transplantation).

Although both nephrologists and PCPs reported similar practices about discussing CC with their patients with advanced CKD (61% versus 54.3%, respectively; $P=0.17$), there were significant differences regarding barriers to discussing CC and knowledge of CC. Fewer nephrologists than PCPs reported as barriers difficulty in determining eligibility (14.3% versus 42.5%, respectively) and limited information about its effectiveness (24.5% and 49.6%, respectively). In contrast, nephrologists were more likely to endorse difficulty in determining whether a patient with CKD would benefit from CC than PCPs (52.8% versus 36.2%, respectively). As expected, nephrologists were more knowledgeable than PCPs about outcomes of CC

Department of Medicine, Universidad Panamericana School of Medicine, Mexico City, Mexico

Correspondence: Dr. Gregorio T. Obrador, Universidad Panamericana School of Medicine, Donatello 59, Col. Insurgentes Mixcoac, 03920 Mexico City, Mexico. Email: gbrador@up.edu.mx

versus dialysis and the Renal Physician Association's guidelines regarding withdrawal or withholding of dialysis.

The results of the work by Parvez *et al.* (14) should be interpreted in light of two significant limitations, namely a low response rate and likely selection bias. After inviting a simple random sample of 16,193 nephrologists and PCPs through three waves of emails, only 431 physicians responded to the survey (crude response rate of 2.7%). The majority of respondents were white men in their 30s and 40s, and nearly one half were in private practice. Also, close to two thirds were nephrologists, and of these, more were in academic practice compared with PCPs. Both the low response rate and the likely selection bias of the respondents limit the generalizability of the results. Indeed, most providers surprisingly reported that a discussion about CC was not difficult to initiate or time consuming and that they were not concerned how their patients would react after such a discussion.

In the second study, O'Hare *et al.* (15) conducted semi-structured interviews at the Veterans Affairs (VA) Puget Sound Healthcare System with 26 providers (74% of a total of 35 invited providers) from various disciplines and specializations who take care of patients with advanced CKD to evoke their perspectives on ACP. Among the 26 providers, there were 13 physicians (from cardiology, intensive care, nephrology, palliative care, psychiatry, and vascular surgery), six nurses (including nurses and nurse practitioners in palliative care, nephrology, and dialysis), three dialysis technicians, two dietitians, and two social workers.

O'Hare *et al.* (15) identified four overlapping themes: (1) a complex and fragmented medical care for this population across settings and providers and over time; (2) lack of a shared understanding and vision of ACP; (3) unclear definition of authority and responsibility for ACP; and (4) lack of collaboration and communication regarding ACP among all providers. Although the results reflected the practice of a small number of providers from a single center, they were consistent with published literature in other settings and populations. However, the single-center design limits their generalizability to settings other than the VA.

These two studies highlight several key points that strongly influence the provider's role in CC and ACP for patients with advanced CKD (14,15). First and foremost, they show confusion regarding terminology for CC. To address this issue, a recent Kidney Disease Improving Global Outcomes (KDIGO) Controversies Conference on Supportive Care in CKD defined CC as "planned holistic patient-centered care for patients with glomerular filtration rate category (G) 5 CKD, that includes interventions to delay progression of kidney disease and minimize risk of adverse events or complications, shared decision-making, active symptom management, detailed communication (including ACP), psychological support, social and family support, and cultural and spiritual domains of care" (8). The KDIGO definition explicitly states that comprehensive CC does not include dialysis (8).

Second, the studies point out the need for more evidence regarding optimal patient selection for CC and comparison of outcomes of CC versus dialysis (16). Although the latter is subject to lead time bias related to imprecisions with GFR estimation and difficulties with randomizing patients,

additional evidence will help address these barriers consistently identified by providers, individualize discussions with patients and families, and facilitate decision making. Third, they show the need to promote a shared vision of ACP and interdisciplinary collaboration among different providers involved in caring for patients with advanced CKD (2).

For the time, it seems reasonable to disseminate formal recommendations and guidelines regarding CC and ACP, particularly among PCPs (3). Also, information should be provided on a variety of tools and online resources, including skills-based training, to facilitate ACP for the health care team, patients, and family with CKD and ultimately, shared decision making (17–20). Additional research is needed to improve patient-centered care as a means of achieving better health outcomes and greater satisfaction among patients with advanced CKD (8).

Disclosures

None.

References

- Treit K, Lam D, O'Hare AM: Timing of dialysis initiation in the geriatric population: Toward a patient-centered approach. *Semin Dial* 26: 682–689, 2013
- Johnson DS, Kapoian T, Taylor R, Meyer KB: Going upstream: Coordination to improve CKD care. *Semin Dial* 29: 125–134, 2016
- Renal Physicians Association: *Shared Decision-Making in the Appropriate Initiation of and Withdrawal from Dialysis*, 2nd Ed., Rockville, MD, Clinical Practice Guideline, 2010
- Murtagh FE, Marsh JE, Donohoe P, Ekbal NJ, Sheerin NS, Harris FE: Dialysis or not? A comparative survival study of patients over 75 years with chronic kidney disease stage 5. *Nephrol Dial Transplant* 22: 1955–1962, 2007
- Carson RC, Juszczak M, Davenport A, Burns A: Is maximum conservative management an equivalent treatment option to dialysis for elderly patients with significant comorbid disease? *Clin J Am Soc Nephrol* 4: 1611–1619, 2009
- Chandna SM, Da Silva-Gane M, Marshall C, Warwicker P, Greenwood RN, Farrington K: Survival of elderly patients with stage 5 CKD: Comparison of conservative management and renal replacement therapy. *Nephrol Dial Transplant* 26: 1608–1614, 2011
- Hussain JA, Mooney A, Russon L: Comparison of survival analysis and palliative care involvement in patients aged over 70 years choosing conservative management or renal replacement therapy in advanced chronic kidney disease. *Palliat Med* 27: 829–839, 2013
- Davison SN, Levin A, Moss AH, Jha V, Brown EA, Brennan F, Murtagh FE, Naicker S, Germain MJ, O'Donoghue DJ, Morton RL, Obrador GT: Executive summary of the KDIGO Controversies Conference on Supportive Care in Chronic Kidney Disease: Developing a roadmap to improving quality care. *Kidney Int* 88: 447–459, 2015
- Okamoto I, Tonkin-Crine S, Rayner H, Murtagh FE, Farrington K, Caskey F, Tomson C, Loud F, Greenwood R, O'Donoghue DJ, Roderick P: Conservative care for ESRD in the United Kingdom: A national survey. *Clin J Am Soc Nephrol* 10: 120–126, 2015
- Morton RL, Tong A, Howard K, Snelling P, Webster AC: The views of patients and carers in treatment decision making for chronic kidney disease: Systematic review and thematic synthesis of qualitative studies. *BMJ* 340: c112, 2010
- O'Hare AM, Armistead N, Schrag WL, Diamond L, Moss AH: Patient-centered care: An opportunity to accomplish the "Three Aims" of the National Quality Strategy in the Medicare ESRD program. *Clin J Am Soc Nephrol* 9: 2189–2194, 2014
- Brinkman-Stoppelenburg A, Rietjens JA, van der Heide A: The effects of advance care planning on end-of-life care: A systematic review. *Palliat Med* 28: 1000–1025, 2014

13. Detering KM, Hancock AD, Reade MC, Silvester W: The impact of advance care planning on end of life care in elderly patients: Randomised controlled trial. *BMJ* 340: c1345, 2010
14. Parvez S, Abdel-Kader K, Pankratz VS, Song M-K, Unruh M: Provider knowledge, attitudes, and practices surrounding conservative management for patients with advanced CKD. *Clin J Am Soc Nephrol* 11: 812–820, 2016
15. O'Hare AM, Szarka J, McFarland LV, Taylor JS, Sudore RL, Trivedi R, Reinke LF, Vig EK: Provider perspectives on advanced care planning for patients with kidney disease: Whose job is it anyway? *Clin J Am Soc Nephrol* 11: 855–866, 2016
16. Tam-Tham H, Thomas CM: Does the evidence support conservative management as an alternative to dialysis for older patients with advanced kidney disease? [published online ahead of print March 17, 2016]. *Clin J Am Soc Nephrol* doi:CJN.01910216
17. Bekker HL, Winterbottom A, Gavaruzzi T, Mooney A, Wilkie M, Davies S, Crane D, Tupling K, Mathers N; Yorkshire Dialysis Decision Aid (YoDDA) Team: *The Dialysis Decision Aid Booklet: Making The Right Choices for You*, Peterborough, United Kingdom, Kidney Research UK, 2015
18. Schell JO, Arnold RM: NephroTalk: Communication tools to enhance patient-centered care. *Semin Dial* 25: 611–616, 2012
19. Alberta Health Services: Alberta Health Services Advance Care Planning. Available at: <http://www.albertahealthservices.ca/9098.asp>. Accessed March 17, 2016
20. Respecting Choices: Advance Care Planning. Available at: <http://respectingchoices.org/>. Accessed March 17, 2016

Published online ahead of print. Publication date available at www.cjasn.org.

See related articles, "Provider Knowledge, Attitudes, and Practices Surrounding Conservative Management for Patients with Advanced CKD," and "Provider Perspectives on Advance Care Planning for Patients with Kidney Disease: Whose Job Is It Anyway?," on pages 812–820 and 855–866, respectively.