

Preservation of Residual Kidney Function and Urine Volume in Patients on Dialysis

Raymond T. Krediet

Clin J Am Soc Nephrol 12: 377–379, 2017. doi: 10.2215/CJN.00330117

Chronic dialysis treatment usually starts before the development of anuria to prevent the major complications of the uremic syndrome. Consequently, most patients will still have urine production at this time. Hemodialysis (HD) has been the main dialysis modality for chronic kidney failure since its start in the early sixties of the last century. Clinical observations at that time showed that most patients became anuric while on dialysis treatment. As a consequence, the preponderance of outcome studies on the effect of HD dose on survival either excluded patients with urine production or assumed that none were present. This policy was also adopted in the guidelines on HD dose that were part of the Dialysis Outcomes Quality Initiative in 1997 (1). The stormy development of continuous ambulatory peritoneal dialysis (CAPD) in the late seventies enabled nephrologists to observe a better preservation of residual kidney function with CAPD compared with HD. This observation was first published from France in 1983 (2) and confirmed in many other studies since, including an analysis of The Netherlands Cooperative Study on the Adequacy of Dialysis (NECOSAD) (3). This analysis also showed that BP and comorbidity were associated with residual renal function in the whole dialysis population investigated and that the rapid decline in the initial 3 months was associated with episodes of hypovolemia. Studies in patients treated with automated peritoneal dialysis (PD) have given variable results, but taking all of the results together, it can be concluded that its decline is similar to that in CAPD (4).

The excretory kidney function consists of glomerular filtration and tubular secretion and reabsorption. In the presence of normal kidney function, about 99% of the glomerular filtrate is reabsorbed. Consequently, urine production is not influenced by the GFR but is influenced by tubular transport mechanisms. A different situation is present in patients on PD. In these patients, urine volume accounted for one half of the variance in GFR (5). A relationship between urine production and GFR is also present in patients on HD (6). Here, the relationship between urine production and GFR in the interdialytic interval had an explained variance of 67%. Apparently, tubuloglomerular feedback mechanisms are disturbed or function differently in patients on dialysis. Inulin clearance is the gold standard for determination of GFR in humans, but its determination

is cumbersome. Creatinine clearance overestimates GFR due to some tubular secretion, and clearance of urea gives an underestimation because of reabsorption, especially in the presence of hypovolemia. The mean of urea and creatinine clearance is closely related to inulin clearance in both patients on HD (6) and patients on PD (7). Therefore, the mean of the two clearances can be considered the reference method for GFR determination in patients on dialysis. This requires a timed and accurate urine collection. To overcome cumbersome urine collections, formulas have been developed for GFR estimation on the basis of plasma concentrations of creatinine and urea. The Modification of Diet in Renal Disease (MDRD) equation (8) is widely used and extensively validated but not in predialysis patients or patients on dialysis. It consists of sex and race in the numerator and creatinine and age in the denominator. Recently, a new equation aimed for use with patients on dialysis, $eGFR_{urea,creat}$ was developed by some of the same group (9). It consists of urea in the numerator and creatinine in the denominator. Also, serum concentrations of low molecular proteins that are removed from the body by glomerular filtration and subsequent reabsorption and breakdown in proximal tubular cells can be used for GFR estimation. β_2 -Microglobulin and cystatin C were related to residual urine production in both HD (10) and PD (11). Equations have been developed for GFR estimations from serum cystatin C in patients on HD and patients on PD (12). By using serum concentrations of low molecular weight proteins, it is assumed that GFR is the only determinant of these. The practical value of measurement or estimation of residual GFR or urine production is especially relevant when they are associated with the risk of death. A reanalysis of the Canada-USA Study performed on patients on incident PD showed that urine volume but not measured GFR was associated with a 36% reduction of the relative risk of death (13). The NECOSAD reported some similar results. Measured GFR was related to survival, whereas urine production only became significant after removal of GFR from the model (14). In HD, renal Kt/V_{urea} was evidently associated with survival (15). The importance of measured GFR (16) or urine production (17) for survival of patients on HD has been confirmed in other prospective studies. The performance of $eGFR$ in the MDRD equation in patients just before the start of dialysis is different. A

Division of Nephrology, Department of Medicine, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Correspondence: Prof. Raymond T. Krediet, Room F4-215, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands. Email: r.t.krediet@amc.uva.nl

number of studies, including an analysis by the European Renal Association-European Dialysis and Transplant Association Registry (18), reported a higher mortality on dialysis of patients who started dialysis with the highest eGFR values, despite extensive corrections for potentially confounding factors. The explanation for this unexpected finding is the dependency of plasma creatinine on muscle mass, which is, in a situation of dialysis initiation, more important for death than the effect of glomerular filtration (19).

A nationwide prospective cohort study (20) in 1946 patients on incident HD and patients on PD from Korea, published in this issue of the *Clinical Journal of the American Society of Nephrology*, shows that urine volume and, to a lesser extent, measured GFR were significantly associated with survival after adjustment for potential confounders but that eGFR_{urea,creat} was not (21). In contrast, the association between eGFR on the basis of serum β 2-microglobulin and survival was significant. This is the largest study on residual kidney function so far. It confirms that urine production is the most important kidney function parameter related to survival of patients on dialysis and also, that eGFR on the basis of plasma concentrations of small solutes only provides insufficient or even misleading information on this issue. Serum concentrations of small proteins may be an alternative but need more investigation.

The importance of urine volume requires a focus on the prevention of hypovolemia. However, hypervolemia should also be avoided, meaning that maintaining euvolemia can be regarded as sailing between Scylla and Charybdis. This difficult task for patients and doctors is easier in the presence of a substantial urine production. Despite the increased relative importance of glomerular filtration in the determination of urine volume in patients on dialysis, pharmacologic modifications of parts of the tubular system are still effective. This has been extensively shown for furosemide, and also recently, for tolvaptan, PD with icodextrin, and angiotensin-converting enzyme inhibitors or angiotensin receptor blockers, although not very convincingly.

Furosemide should be given in high dosages, because the concentration in the urine of the proximal tubule determines its effect. Active secretion of furosemide into the lumen of the proximal tubule occurs by a transport mechanism that requires high plasma concentrations because of the impaired function of the proximal tubule. Farther distally, urine furosemide inhibits sodium reabsorption in the thick ascending limb of the loop of Henle. It is immediately effective in both patients on HD (21,22) and patients on PD (23), in whom it also increased the fractional excretion of sodium and potassium. Furosemide administration has no effect on GFR or its decline with time on dialysis (23,24). Tolvaptan has been studied extensively in patients on dialysis, but only one study has been published on its effect on urine production in a small number of incident diabetic patients on PD (25). After 1 year, urine volume and renal creatinine clearance had remained stable, whereas they decreased in the control group. It is well known that dialysis solutions containing icodextrin as an osmotic agent increase peritoneal ultrafiltration, especially during long dialysis dwells, but only one recent study reported an effect on the time course of urine volume without an effect on residual GFR (26). It is likely that circulating oligosaccharides generated by breakdown of absorbed icodextrin increase the osmotic load for

the kidneys, leading to a higher urine production. A number of studies have been published on possible effects of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers on preservation of residual renal function but without consistent results (27).

Maintenance of euvolemia is the most important issue in prolongation of survival of patients on dialysis. Preservation of residual kidney function and especially, urine production has an important contribution to the prevention of overhydration. Using eGFR on the basis of plasma concentrations of creatinine and/or urea gives misleading information in patients on dialysis, which makes the use of timed urine collections unavoidable. The role of serum concentrations of low molecular weight proteins has not been established yet. Furosemide in high dosages is the only therapeutic intervention currently available to increase urine production, but it has no effect on the other aspects of residual kidney function.

Disclosures

None.

References

- Owen W Jr., Roberts L; National Kidney Foundation: NKF-DOQI clinical practice guidelines for hemodialysis adequacy. *Am J Kidney Dis* 30[Suppl 2]: S15–S66, 1997
- Rottembourg J, Issad B, Gallego JL, Degoulet P, Aime F, Gueffaf B, Legrain M: Evolution of residual renal function in patients undergoing maintenance haemodialysis or continuous ambulatory peritoneal dialysis. *Proc Eur Dial Transplant Assoc* 19: 397–403, 1983
- Jansen MAM, Hart AAM, Korevaar JC, Dekker FW, Boeschoten EW, Krediet RT; NECOSAD Study Group: Predictors of the rate of decline of residual renal function in incident dialysis patients. *Kidney Int* 62: 1046–1053, 2002
- Bieber SD, Burkart J, Golper TA, Teitelbaum I, Mehrotra R: Comparative outcomes between continuous ambulatory and automated peritoneal dialysis: A narrative review. *Am J Kidney Dis* 63: 1027–1037, 2014
- Johnson DW, Mudge DW, Sturtevant JM, Hawley CM, Campbell SB, Isbel NM, Hollett P: Predictors of decline of residual renal function in new peritoneal dialysis patients. *Perit Dial Int* 23: 276–283, 2003
- van Olden RW, van Acker BAC, Koomen GCM, Krediet RT, Arisz L: Time course of inulin and creatinine clearance in the interval between two haemodialysis treatments. *Nephrol Dial Transplant* 10: 2274–2280, 1995
- Van Olden RW, Krediet RT, Struijk DG, Arisz L: Measurement of renal function in patients treated with continuous ambulatory peritoneal dialysis. *J Am Soc Nephrol* 7: 745–750, 1996
- Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D; Modification of Diet in Renal Disease Study Group: A more accurate method to estimate glomerular filtration rate from serum creatinine: A new prediction equation. *Ann Intern Med* 130: 461–470, 1999
- Shafi T, Michels WM, Levey AS, Inker LA, Dekker FW, Krediet RT, Hoekstra T, Schwartz GJ, Eckfeldt JH, Coresh J: Estimating residual kidney function in dialysis patients without urine collection. *Kidney Int* 89: 1099–1110, 2016
- Kabanda A, Jadoul M, Pochet JM, Lauwerys R, van Ypersele de Strihou C, Bernard A: Determinants of the serum concentrations of low molecular weight proteins in patients on maintenance hemodialysis. *Kidney Int* 45: 1689–1696, 1994
- Kabanda A, Goffin E, Bernard A, Lauwerys R, van Ypersele de Strihou C: Factors influencing serum levels and peritoneal clearances of low molecular weight proteins in continuous ambulatory peritoneal dialysis. *Kidney Int* 48: 1946–1952, 1995
- Hoek FJ, Korevaar JC, Dekker FW, Boeschoten EW, Krediet RT: Estimation of residual glomerular filtration rate in dialysis patients from the plasma cystatin C level. *Nephrol Dial Transplant* 22: 1633–1638, 2007
- Bargman JM, Thorpe KE, Churchill DN; CANUSA Peritoneal Dialysis Study Group: Relative contribution of residual renal function and peritoneal clearance to adequacy of dialysis: A

- reanalysis of the CANUSA study. *J Am Soc Nephrol* 12: 2158–2162, 2001
14. Termorshuizen F, Korevaar JC, Dekker FW, van Manen JG, Boeschoten EW, Krediet RT; NECOSAD Study Group: The relative importance of residual renal function compared with peritoneal clearance for patient survival and quality of life: An analysis of the Netherlands Cooperative Study on the Adequacy of Dialysis (NECOSAD)-2. *Am J Kidney Dis* 41: 1293–1302, 2003
 15. Termorshuizen F, Dekker FW, van Manen JG, Korevaar JC, Boeschoten EW, Krediet RT; NECOSAD Study Group: Relative contribution of residual renal function and different measures of adequacy to survival in hemodialysis patients: An analysis of the Netherlands Cooperative Study on the Adequacy of Dialysis (NECOSAD)-2. *J Am Soc Nephrol* 15: 1061–1070, 2004
 16. Shemin D, Bostom AG, Laliberty P, Dworkin LD: Residual renal function and mortality risk in hemodialysis patients. *Am J Kidney Dis* 38: 85–90, 2001
 17. Shafi T, Jaar BG, Plantinga LC, Fink NE, Sadler JH, Parekh RS, Powe NR, Coresh J: Association of residual urine output with mortality, quality of life, and inflammation in incident hemodialysis patients: The choices for healthy outcomes in caring for end-stage renal disease (CHOICE) study. *Am J Kidney Dis* 56: 348–358, 2010
 18. Stel VS, Dekker FW, Ansell D, Augustijn H, Casino FG, Collart F, Finne P, Ioannidis GA, Salomone M, Traynor JP, Zurriaga O, Verrina E, Jager KJ: Residual renal function at the start of dialysis and clinical outcomes. *Nephrol Dial Transplant* 24: 3175–3182, 2009
 19. Grootendorst DC, Michels WM, Richardson JD, Jager KJ, Boeschoten EW, Dekker FW, Krediet RT; NECOSAD Study Group: The MDRD formula does not reflect GFR in ESRD patients. *Nephrol Dial Transplant* 26: 1932–1937, 2011
 20. Lee MJ, Park JT, Park KS, Kwon YE, Oh HJ, Yoo TH, Kim YL, Kim YS, Yang CW, Kim NH, Kang SW, Han SH: Prognostic Value of Residual Urine Volume, GFR by 24-hour Urine Collection, and eGFR in Patients Receiving Dialysis. *Clin J Am Soc Nephrol* 12: 426–434, 2017
 21. van Olden RW, van Meyel JJM, Gerlag PGG: Acute and long-term effects of therapy with high-dose furosemide in chronic hemodialysis patients. *Am J Nephrol* 12: 351–356, 1992
 22. Sjolund J, Garcia Anton D, Bayes LY, Hoekstra T, Dekker FW, Munoz Mendoza J: Diuretics, limited ultrafiltration, and residual renal function in incident hemodialysis patients: A case series. *Semin Dial* 29: 410–415, 2016
 23. van Olden RW, Guchelaar HJ, Struijk DG, Krediet RT, Arisz L: Acute effects of high-dose furosemide on residual renal function in CAPD patients. *Perit Dial Int* 23: 339–347, 2003
 24. Medcalf JF, Harris KPG, Walls J: Role of diuretics in the preservation of residual renal function in patients on continuous ambulatory peritoneal dialysis. *Kidney Int* 59: 1128–1133, 2001
 25. Hiramoto T, Hobo A, Hayasaki T, Kabu K, Furuta S: A pilot study examining the effects of tolvaptan on residual renal function in peritoneal dialysis for diabetics. *Perit Dial Int* 35: 552–558, 2015
 26. Chang TI, Ryu DR, Yoo TH, Kim HJ, Kang EW, Kim H, Chang JH, Kim DK, Moon SJ, Yoon SY, Han SH; Yonsei Associate Network CHronic Kidney Disease Trial (YACHT) investigators: Effect of icodextrin solution on the preservation of residual renal function in peritoneal dialysis patients: A randomized controlled study. *Medicine (Baltimore)* 95: e2991, 2016
 27. Zhang L, Zeng X, Fu P, Wu HM: Angiotensin-converting enzyme inhibitors and angiotensin receptor blockers for preserving residual kidney function in peritoneal dialysis patients. *Cochrane Database Syst Rev* 6: CD009120, 2014

Published online ahead of print. Publication date available at www.cjasn.org.

See related article, “Prognostic Value of Residual Urine Volume, GFR by 24-hour Urine Collection, and eGFR in Patients Receiving Dialysis,” on pages 426–434.