





# Comparative Effects of Etelcalcetide and Maxacalcitol on Serum Calcification Propensity in Secondary Hyperparathyroidism

## A Randomized Clinical Trial

Tetsuo Shoji <sup>1,2</sup>, Shinya Nakatani <sup>3</sup>, Daijiro Kabata,<sup>4</sup> Katsuhito Mori,<sup>5</sup> Ayumi Shintani <sup>4</sup>, Hisako Yoshida,<sup>4</sup> Kanae Takahashi,<sup>4</sup> Keiko Ota,<sup>6</sup> Hisako Fujii,<sup>7</sup> Shinichiro Ueda,<sup>8</sup> Shinichi Nishi,<sup>9</sup> Tatsuya Nakatani,<sup>10</sup> Minoru Yoshiyama,<sup>11</sup> Kiyoshi Goto,<sup>12</sup> Takayoshi Hamada,<sup>13</sup> Masahito Imanishi,<sup>14</sup> Eiji Ishimura,<sup>15</sup> Sosuke Kagitani,<sup>16</sup> Yoshikazu Kato,<sup>17</sup> Yasuro Kumeda,<sup>18</sup> Kiyoshi Maekawa,<sup>19</sup> Takayasu Matsumura,<sup>20</sup> Harumi Nagayama,<sup>21</sup> Yasue Obi,<sup>22</sup> Yoshiteru Ohno,<sup>23</sup> Yoshinori Sai,<sup>24</sup> Mayumi Sakurai,<sup>25</sup> Satoshi Sasaki,<sup>26</sup> Kaori Shidara,<sup>27</sup> Shigeichi Shoji <sup>28</sup>, Yoshihiro Tsujimoto,<sup>29</sup> Kenjiro Yamakawa,<sup>30</sup> Hideaki Yasuda,<sup>31</sup> Shozo Yodoi,<sup>32</sup> Masaaki Inaba,<sup>2,3,5</sup> and Masanori Emoto<sup>2,3,5</sup>

Due to the number of contributing authors, the affiliations are listed at the end of this article.

### Correspondence:

Prof. Tetsuo Shoji, Department of Vascular Medicine, Osaka City University Graduate School of Medicine; and Vascular Science Center for Translational Research, Osaka City University Graduate School of Medicine, 1-4-3, Asahi-machi, Abeno-ku, Osaka 545-8585, Japan. Email: t-shoji@med.osaka-cu.ac.jp

### Abstract

**Background and objectives** Vitamin D receptor activators and calcimimetics (calcium-sensing receptor agonists) are two major options for medical treatment of secondary hyperparathyroidism. A higher serum calcification propensity (a shorter T<sub>50</sub> value) is a novel surrogate marker of calcification stress and mortality in patients with CKD. We tested a hypothesis that a calcimimetic agent etelcalcetide is more effective in increasing T<sub>50</sub> value than a vitamin D receptor activator maxacalcitol.

**Design, setting, participants, & measurements** A randomized, multicenter, open-label, blinded end point trial with active control was conducted in patients with secondary hyperparathyroidism undergoing hemodialysis in Japan. Patients were randomly assigned to receive intravenous etelcalcetide 5 mg thrice weekly (etelcalcetide group) or intravenous maxacalcitol 5 or 10 μg thrice weekly (maxacalcitol group). The primary, secondary, and tertiary outcomes were changes in T<sub>50</sub> value, handgrip strength, and score of the Dementia Assessment Sheet for Community-Based Integrated Care System from baseline to 12 months, respectively.

**Results** In total, 425 patients from 23 dialysis centers were screened for eligibility, 326 patients were randomized (etelcalcetide, *n* = 167; control, *n* = 159), and 321 were included in the intention-to-treat analysis (median age, 66 years; 113 women [35%]). The median (interquartile range) of T<sub>50</sub> value was changed from 116 minutes (interquartile range, 90–151) to 131 minutes (interquartile range, 102–176) in the maxacalcitol group, whereas it was changed from 123 minutes (interquartile range, 98–174) to 166 minutes (interquartile range, 127–218) in the etelcalcetide group. The increase in T<sub>50</sub> value was significantly greater in the etelcalcetide group (difference in change, 20 minutes; 95% confidence interval, 7 to 34 minutes; *P* = 0.004). No significant between-group difference was found in the change in handgrip strength or in the Dementia Assessment Sheet for Community-Based Integrated Care System score.

**Conclusions** Etelcalcetide was more effective in increasing T<sub>50</sub> value than maxacalcitol among patients on hemodialysis with secondary hyperparathyroidism. There was no difference in handgrip strength or cognition between the two drugs.

**Clinical Trial registry name and registration number:** VICTORY; UMIN000030636 and jRCTs051180156

CJASN 16: 599–612, 2021. doi: <https://doi.org/10.2215/CJN.16601020>

### Introduction

Mineral bone disorder in CKD is considered as one of the important factors explaining the high mortality rate in patients with CKD, including those receiving hemodialysis (1–4). Secondary hyperparathyroidism results in high turnover bone, bone mineral loss, calcification of soft tissues including arteries, and a higher risk of cardiovascular disease and mortality.

There are two major options for medical treatment for secondary hyperparathyroidism, namely, vitamin D receptor activators (VDRAs) and calcimimetics (calcium-sensing receptor agonists). The Kidney Disease Improving Global Outcomes (KDIGO) 2017 clinical practice guideline (5) suggests the use of calcimimetics, calcitriol, vitamin D analogs, or a combination of calcimimetics with calcitriol or vitamin D analogs in

patients with CKD undergoing dialysis requiring parathyroid hormone (PTH)-lowering therapy. The guideline does not recommend calcimimetics or VDRA as a preferred choice of treatment. Choice may depend on other outcomes than PTH levels that are important for individual patients, such as mortality, calcification stress, sarcopenia, and cognition. So far, however, no such comparison was made between the two classes of drugs (6).

Phosphate plays important roles in vascular calcification (7) and mortality (3,4). The toxicity of phosphate is thought to be exerted by calciprotein particles, nanosized colloidal particles consisted of calcium phosphate crystals stabilized by a glycoprotein fetuin-A (8,9). The primary calciprotein particles <100 nm are considered as physiologic carriers of calcium and phosphate. However, after they undergo spontaneous transformation to the secondary calciprotein particles >100 nm, these larger calciprotein particles have highly cytotoxic and proinflammatory properties (10). The propensity of transformation to the secondary calciprotein particles can be assessed by measuring time required to transform into the secondary calciprotein particles *in vitro*, called serum calcification propensity ( $T_{50}$ ) (11). A shorter  $T_{50}$ , which means a higher calcification propensity, was associated with arterial calcification (12), arterial stiffness (13), and a higher risk of all-cause death in cohorts of dialysis-independent (14) and -dependent patients with CKD (15), and also in kidney transplant recipients (16). Thus,  $T_{50}$  could be a surrogate marker of poor clinical outcomes attributable to calcification stress.

Sarcopenia and cognitive impairment have emerged as important health concerns in aging societies (17), and vitamin D may play some roles (18). Vitamin D receptor knockout mice show abnormal development of skeletal muscles (19). Observational human studies reported associations of vitamin D status with physical performance (20) and cognitive decline (21) in the elderly and in patients with CKD (22,23) as well.

We hypothesized that treatment with a calcimimetic agent is more effective in suppressing  $T_{50}$  compared with treatment with a VDRA, whereas a VDRA may be better in preventing decline in muscle strength and/or cognition in patients with secondary hyperparathyroidism undergoing hemodialysis.

## Materials and Methods

### Study Oversight

This trial named "VDRA vs. Intravenous Calcimimetics in the Treatment Of Renal patients with secondary hYperparathyroidism" (VICTORY) was designed and overseen by the representative principal investigator and the steering committee at Osaka City University, and it was conducted in accordance with the principles of the Declaration of Helsinki; the Ethical Guidelines for Medical and Health Research Involving Human Subjects by the Ministry of Health, Labor and Welfare, Japan (December 22, 2014; revised on February 28, 2017); and the Clinical Trial Act (act no. 16 of April 14, 2017). The protocol was reviewed and approved by the Ethical Committee of Osaka City University Graduate School of Medicine (approval no. 3879) and the Osaka City University Hospital Certified Review Board (approval no. OCU0011). All participants gave written

informed consent before the study. To address the Clinical Trial Act, which came into force on April 1, 2019, this trial was registered in the Japan Registry of Clinical Trials after enrollment of patients, in addition to the registration in the University Hospital Medical Information Network Clinical Trials Registry, which was made before the patient enrollment.

### Study Design

This clinical trial with a randomized, open-label, blinded end point design compared the effects of a calcimimetic agent etelcalcetide (intervention group) versus a VDRA maxacalcitol (control group) on the change in  $T_{50}$  from baseline to 12 months in patients with secondary hyperparathyroidism receiving maintenance hemodialysis. The protocol and statistical analysis plan are available in Supplemental Material. Supplemental Figure 1 shows the synopsis of this trial.

### Study Population

The participants were patients with secondary hyperparathyroidism aged 20–80 years who were receiving hemodialysis thrice weekly, whose serum intact PTH was  $\geq 241$  pg/ml and serum corrected calcium was  $\geq 8.4$  mg/dl, and who were not taking any calcimimetics or VDRA at randomization. These laboratory values were derived from the 2012 version of the clinical practice guideline by the Japanese Society for Dialysis Therapy (24). Detailed inclusion and exclusion criteria are listed in Supplemental Table 1. Eligible patients were screened at 23 dialysis facilities in Osaka, Japan.

### Randomization

Randomization (1:1) using computer-generated random numbers with a permuted block scheme was performed. Randomization was stratified by sex (men or women), age (<65 or  $\geq 65$  years old), and serum phosphate level (<6.0 or  $\geq 6.0$  mg/dl). Allocation was distributed through the web-based system called Research Electronic Data Capture (<https://projectredcap.org/about/>) (25,26) at Osaka City University (<http://www.hosp.med.osaka-cu.ac.jp/self/hyokac/redcap/index.shtml>).

### Intervention and Follow-Up in the Etelcalcetide Group

The patients allocated to the etelcalcetide (intervention) group were assigned to receive intravenous etelcalcetide thrice weekly administered *via* blood access at the end of each dialysis session. The initial dose was set at 5 mg following directions on the package insert of etelcalcetide hydrochloride (PARSABIV; Ono Pharmaceuticals, Co. Ltd., Osaka, Japan). Dose adjustment was permitted to achieve the target range of intact PTH (between 60 and 240 pg/ml) and to avoid hypocalcemia in the dose range between 2.5 and 15 mg per shot. Oral VDRA were allowed only to avoid hypocalcemia even after the dose of etelcalcetide was reduced to 2.5 mg per shot. Intravenous VDRA preparations were prohibited, but they could be used if hypocalcemia could not be relieved by dose reduction of etelcalcetide and use of an oral VDRA. Temporal cessation of etelcalcetide was made if corrected calcium was <7.5 mg/dl, and etelcalcetide was restarted when corrected

calcium returned to  $\geq 8.4$  mg/dL. In this group, “discontinuation of study drug” was defined when etelcalcetide was not administered continuously for 42 days or longer regardless of the reason.

### Intervention and Follow-Up in the Maxacalcitol Group

The patients allocated to the maxacalcitol (control) group were assigned to receive intravenous maxacalcitol thrice weekly administered *via* blood access at the end of each dialysis session. The initial dose was set at 5  $\mu$ g thrice weekly in patients with serum intact PTH  $< 500$  pg/ml and 10  $\mu$ g thrice weekly in patients with intact PTH  $\geq 500$  pg/ml following directions on the package insert of maxacalcitol (OXAROL; Chugai Pharmaceuticals, Co. Ltd., Tokyo, Japan). Dose adjustment was permitted to achieve the target range of intact PTH (between 60 and 240 pg/ml) and to avoid hypercalcemia in the dose range between 2.5 and 10  $\mu$ g per shot. Oral calcimimetics (cinacalcet and evocalcet) were allowed only to avoid hypercalcemia even after the dose of maxacalcitol was reduced to 2.5  $\mu$ g per shot. Intravenous calcimimetic agent (etelcalcetide) was prohibited, but it could be used if hypercalcemia could not be relieved by dose reduction of maxacalcitol and use of an oral calcimimetic agent. Temporal cessation of maxacalcitol was made if corrected calcium was  $> 11.5$  mg/dl, and maxacalcitol was restarted when corrected calcium returned to  $< 11.0$  mg/dl. In this group, “discontinuation of study drug” was defined when maxacalcitol was not administered continuously for 42 days or longer regardless of the reason.

### Study Outcomes and Adverse Events

The primary, secondary, and tertiary efficacy outcomes were the changes from study visit at 0–12 months in  $T_{50}$  (11), handgrip strength (27), and cognition (28), respectively.  $T_{50}$  was measured using the method by Pasch *et al.* (11) in our laboratory (29). Cognition was assessed by the Dementia Assessment Sheet for Community-Based Integrated Care System (DASC-21) (28). Details of these methods are available in Supplemental Tables 2 and 3.

Supplemental Table 4 shows other subsidiary outcomes, including (1) changes in  $T_{50}$  at visits from 0 to 3 months and from 0 to 6 months; (2) presence or absence of achieving the target ranges of serum corrected calcium, phosphate, and intact PTH (24) at 3, 6, and 12 months; and (3)  $> 30\%$  reduction in intact PTH from baseline at 3, 6, and 12 months.

The safety assessment included all serious adverse events (SAEs), hypocalcemia and hypercalcemia requiring cessation of study drug, symptoms (nausea, vomiting, dysgeusia, diarrhea, itching, irritability, and others), and falls.

### Other Measurements

Serum levels of albumin, calcium, phosphate, intact PTH, intact fibroblast growth factor-23 (FGF23), fetuin-A, and magnesium were centrally measured at SRL, Inc. (Tokyo, Japan). Other clinical data, including laboratory measurements at each site, were collected from the medical records.

### Interim Analysis and Stopping Guideline

The independent monitoring committee conducted interim data viewing to review the status of recruitment and interim safety profile and gave a recommendation to continue the trial as planned. However, no interim analysis of the primary outcome was done for possible early stopping of recruitment. We recruited participants as much as possible within the scheduled recruitment period.

### Sample Size Determination

Sample size was computed comparing mean  $T_{50}$  at 12 months between the two groups. Because there are few published data available on  $T_{50}$ , we estimated  $T_{50}$  value on the basis of the reported association between  $T_{50}$  and phosphate level by Pasch *et al.* (15). The mean (SD)  $T_{50}$  value at 12 months among the control group was assumed to be 215 (84) minutes on the basis of the mean  $T_{50}$  level among patients on hemodialysis with elevated PTH levels in the work by Pasch *et al.* (15). The mean  $T_{50}$  at 12 months in the intervention group was assumed to be 240 minutes according to the average change of phosphate level among the patients using etelcalcetide in the work by Block *et al.* (30) and that for the patients using maxacalcitol in the work by Hayashi *et al.* (31). At two-tailed significance level of 5%, 178 patients are required to achieve 80% statistical power. On the basis of the above calculations, assuming that the frequency of dropout is approximately 10%, it would be necessary to enroll 200 participants per group for a total of 400 participants.

### Statistical Analyses

We defined three populations for analysis: a full analysis set, a per-protocol set, and a safety analysis set. The detailed definitions are available in the protocol and statistical analysis plan (Supplemental Material). The primary efficacy analysis was done with full analysis set according to the principle of intention to treat. Missing data were imputed by the last observation carried forward method. In the per-protocol set, last observation carried forward was done up to the time of discontinuation of treatment or withdrawal from the study. To describe results, continuous data were summarized as medians and interquartile ranges, and categorical variables were summarized as numbers and percentages. For comparisons of the outcomes between the allocation groups, we performed multivariable-adjusted linear regression analyses with adjustment for the stratification variables used for randomization, including age, sex, and phosphate level at the visit at 0 months. Least square means and 95% confidence intervals (95% CIs) were calculated for each group, and the estimated absolute differences between the groups were reported with 95% CIs. Although we planned to transform the residuals when normality assumption was not met, no transformation was done (excluding intact FGF23) because the residuals were deemed to follow normal distribution. All of these statistical inferences were made at two-sided significance level of 5%. All calculations were conducted using R version 4.0.2 (<https://cran.r-project.org/>).

## Results


### Flow and Baseline Characteristics of Participants

Figure 1 shows the flow of participants. The participant recruitment was open on February 1, 2018; the first patient was enrolled on February 28, 2018; and the last patient was enrolled on January 9, 2019. The last day of follow-up was December 27, 2019. Eligibility was assessed in 425 pre-enrolled patients. Following a washout period for up to 8 weeks, 99 patients were found not to meet the eligibility criteria (serum corrected calcium and intact PTH levels in most cases). Finally, because of the scheduled period for enrollment, we ended the enrollment and randomization with 326 patients. Of the 326 patients who were randomized, two patients were excluded from the safety analysis set because they did not receive the study drug. Additionally, three patients were excluded from the full analysis set because of protocol violation. Further, two patients were

excluded from the per-protocol set because they were found ineligible during intervention. Thus, the safety analysis set, the full analysis set, and the per-protocol set included 324, 321, and 319 participants, respectively. Table 1 gives the baseline characteristics of the two groups in the full analysis set.

### Changes in Laboratory Data and Medications

Figure 2 shows the changes in select laboratory data. Intact PTH levels were decreased following the interventions in both groups. Phosphate level showed a modest decrease in the etelcalcetide group, whereas it stayed almost unchanged in the maxacalcitol group. Corrected calcium and calcium  $\times$  phosphate product decreased in the etelcalcetide group, whereas they increased in the maxacalcitol group. FGF23 showed a decrease in the etelcalcetide



**Figure 1.** | This chart indicates the flow of participants. SAE, serious adverse event.

**Table 1. Baseline characteristics of participants in a randomized clinical trial comparing the effects of etelcalcetide and maxacalcitol on serum calcification propensity**

Characteristics	Etelcalcetide Group, n=165	Maxacalcitol Group, n=156
Age, yr	66 (55–71)	66 (57–71)
Men, N (%)	110 (67)	98 (63)
Women, N (%)	55 (33)	58 (37)
Duration of hemodialysis, yr	6.2 (3.1–14.8)	7.1 (3.7–14.3)
Diabetic kidney disease, N (%)	69 (42)	56 (36)
<b>Medical history, N (%)</b>		
Cerebral infarction	12 (7)	10 (6)
Cerebral hemorrhage	5 (3)	1 (1)
Subarachnoidal hemorrhage	0 (0)	1 (1)
Myocardial infarction	8 (5)	11 (7)
Coronary revascularization	15 (9)	21 (13)
Amputation of ischemic limb	5 (3)	4 (3)
Leg artery intervention/bypass	3 (2)	12 (8)
Atrial fibrillation	13 (8)	8 (5)
Bone fracture	13 (8)	23 (15)
Falls in the preceding year	19 (12)	19 (12)
Systolic BP, mm Hg	145 (132–163)	151 (136–163)
Diastolic BP, mm Hg	78 (69–88)	80 (71–87)
Height, cm	162 (157–169)	162 (154–168)
Weight, kg	60 (52–69)	59 (50–68)
Body mass index, kg/m <sup>2</sup>	22.5 (20.4–25.2)	22.2 (20.0–25.5)
<b>Serum biochemistry</b>		
Albumin, g/dl	3.7 (3.5–3.9)	3.7 (3.5–3.9)
Intact PTH, pg/ml	418 (313–563)	436 (338–600)
Corrected calcium, mg/dl	9.2 (8.8–9.6)	9.2 (8.9–9.5)
Phosphate, mg/dl	5.5 (4.6–6.2)	5.6 (4.9–6.6)
Magnesium, mg/dl	2.6 (2.4–2.8)	2.6 (2.4–2.8)
Intact FGF23, pg/ml	3500 (1290–8300)	4100 (1140–10,625)
Fetuin-A, μg/ml	253 (216–310)	254 (222–299)
<b>Phosphate binder use, N (%)</b>		
Calcium carbonate	85 (52)	81 (52)
Lanthanum carbonate	83 (50)	82 (53)
Ferric citrate hydrate	32 (19)	30 (19)
Sucroferric oxyhydroxide	17 (10)	10 (6)
Sevelamer hydrochloride	35 (21)	28 (18)
Bixalomer	12 (7)	13 (8)
Magnesium oxide use, N (%)	2 (1)	5 (3)
<b>Hemodialysis conditions</b>		
Time of dialysis per week, h	12.0 (12.0–12.3)	12.0 (12.0–12.0)
Surface area of dialyzer membrane, m <sup>2</sup>	2.1 (1.8–2.1)	2.1 (1.5–2.1)
Kt/V	1.50 (1.33–1.71)	1.50 (1.33–1.70)
Blood flow rate, ml/h	230 (200–250)	220 (200–250)
Dialysate flow rate, ml/h	500 (500–500)	500 (500–500)
<b>Dialysate calcium concentration, mEq/L, N (%)</b>		
2.5	25 (15)	25 (16)
2.75	87 (53)	85 (54)
3.0	53 (32)	46 (29)

Continuous variables were summarized as medians (interquartile ranges), and categorical variables were summarized as numbers (percentages). PTH, parathyroid hormone; FGF23, fibroblast growth factor-23.


group, whereas an increase was observed in the maxacalcitol group. Fetuin-A and magnesium stayed unchanged in both groups.

Figure 3 summarizes changes in medications related to CKD mineral bone disorder. In the etelcalcetide group, discontinuation of the study drug occurred in ten patients. Use of calcium carbonate was increased, whereas use of lanthanum carbonate was decreased. Oral alfacalcidol was used in 44 of 144 patients (31%) at the visit at 12 months to treat hypocalcemia, but intravenous VDRA were not used in this group. In the maxacalcitol group, four patients discontinued the study drug, and 17 patients received oral calcimimetics (cinacalcet or evocalcet). Use of calcium carbonate was decreased, whereas use of lanthanum carbonate and use of

sucroferric oxyhydroxide were increased. Cinacalcet and evocalcet were used in some patients, but etelcalcetide was not used in this group. Dialysate calcium concentration was not changed during the trial in either group.

### Primary, Secondary, and Tertiary Outcomes

Table 2 gives the results on T<sub>50</sub>, handgrip strength, and DASC-21 at the visits at 0 and 12 months and the changes in these measurements during 12 months in each group. Figure 4 graphically depicts these data. The change in T<sub>50</sub> was significantly greater by 20 minutes in the etelcalcetide group versus the maxacalcitol group. There was no significant between-group difference in the change in



**Figure 2.** | The graphs depict the changes in laboratory data during the trial. Circles and vertical bars represent medians and interquartile ranges. Differences in calcium (Ca)  $\times$  phosphate (P) product and intact fibroblast growth factor-23 (FGF23) levels are significant at visits at 3, 6, and 12 months (M;  $P < 0.001$  for all). *Post hoc* between-group comparison showed significant differences in serum P and corrected Ca levels at visits at 3, 6, and 12 M ( $P < 0.001$  for all). These comparisons were done after adjustment for age, sex, and serum P at baseline. PTH, parathyroid hormone.

handgrip strength or in the change in DASC-21. A *post hoc* sensitivity analysis using a mixed effect model also showed a significant difference in the primary outcome between the groups at the visits at 3, 6, and 12 months ( $P < 0.001$  for all).

#### Other Subsidiary Efficacy Outcomes

Other subsidiary efficacy outcomes are summarized in Table 3.


#### Additional Efficacy Analyses with the Per-Protocol Set

Supplemental Table 5 shows analyses of the primary, secondary, and tertiary efficacy outcomes using the per-protocol set. These results were essentially the same with the results using the full analysis set.

#### Adverse Events

Table 4 summarizes adverse events. The total numbers of reports of SAEs were 41 (32 cases; 19.4%) in the





**Figure 3.** | These graphs show the changes in medications during the trial. The study drug (etelcalcetide or maxacalcitol) was continued in most of the participants in each group. In the etelcalcetide group, the use of alfacalcidol, the increased use of calcium carbonate, and the decreased use of lanthanum carbonate were noted. Mg, magnesium; VDRA, vitamin D receptor activator.

etelcalcetide group and 70 (44 cases; 27.7%) in the maxacalcitol group. The most frequent symptom was itching, and it was comparable between the two groups. The numbers of falls were comparable between the two groups. Temporal cessation of study drug occurred in 33 occasions in 30 participants (18.2%) in the etelcalcetide group and in ten occasions in ten participants (6.3%) in the maxacalcitol group. These dose interruptions were all due to


hypocalcemia and hypercalcemia in the etelcalcetide group and the maxacalcitol group, respectively.

**Discussion**

This study compared the effects of etelcalcetide and maxacalcitol on T<sub>50</sub>, handgrip strength, and DASC-21 as the primary, secondary, and tertiary outcomes, respectively, in

Outcomes	Etelcalcetide Group	Maxacalcitol Group	Between-Group Difference
<b>T<sub>50</sub>, min</b>			
0 mo	123 (98–174) n=165	116 (90–151) n=156	
3 mo	141 (117–175) n=149	108 (84–145) n=148	
6 mo	153 (117–190) n=148	105 (79–134) n=141	
12 mo	166 (127–218) n=143	131 (102–176) n=133	
Change from 0 to 12 mo	29 (16 to 43) n=165	9 (–4 to 22) n=156	20 (7 to 34) P=0.004
<b>Handgrip strength, kg</b>			
0 mo	23.5 (17.4–28.4) n=164	22.7 (16.4–29.0) n=155	
12 mo	22.9 (17.8–28.4) n=142	21.6 (16.1–27.8) n=135	
Change from 0 to 12 mo	0.0 (–1.0 to 1.1) n=164	–0.1 (–1.1 to 0.9) n=155	0.2 (–0.9 to 1.2) P=0.76
<b>DASC-21, points</b>			
0 mo	22.0 (21.0–23.0) n=162	21.0 (21.0–23.0) n=150	
12 mo	21.5 (21.0–23.0) n=140	21.0 (21.0–23.0) n=133	
Change from 0 to 12 mo	–0.0 (–1.0 to 1.0) n=162	0.6 (–0.5 to 1.6) n=150	–0.6 (–1.6 to 0.5) P=0.29

Measurements at each visit were summarized as medians (interquartile ranges) with number of patients. Changes from 0 to 12 months and between-group differences were least square means (95% confidence intervals), which were estimated with imputation of missing data by the last observation carried forward method. T<sub>50</sub>, serum calcification propensity; 0 mo, visit at baseline; 3 mo, visit at 3 months; 6 mo, visit at 6 months; 12 mo, visit at 12 months; DASC-21, Dementia Assessment Sheet for Community-Based Integrated Care System.



**Figure 4.** | The increase in T<sub>50</sub> (primary outcome) was greater in the etelcalcetide group than the maxacalcitol group. No difference was significant in the change in handgrip strength (secondary outcome) or DASC-21 score (tertiary outcome) between the groups. The left panels show medians (interquartile ranges), and the right panels show least square means (95% confidence intervals) of the changes from 0 to 12 M in serum calcification propensity (T<sub>50</sub>), handgrip strength, and Dementia Assessment Sheet for Community-Based Integrated Care System (DASC-21). *P* values are for between-group comparison. The individual values are given in Table 2.

patients with secondary hyperparathyroidism undergoing hemodialysis. The change in T<sub>50</sub> following the intervention for 12 months was significantly greater in the etelcalcetide group than in the maxacalcitol group, whereas there was no difference in the change in handgrip strength or DASC-21 between the two groups.

The observed between-group difference in the change in T<sub>50</sub> can be explained mainly by the different effects of the two drugs on phosphate and calcium levels, because T<sub>50</sub> was reported to correlate inversely with serum phosphate and calcium levels (15). In this trial, *post hoc* analyses showed significant between-group differences in serum phosphate and corrected calcium levels at the visits at

3 months ( $P=0.005$ ), 6 months ( $P<0.001$ ), and 12 months ( $P<0.001$ ). Serum magnesium, another factor that increases T<sub>50</sub> (32,33), remained unchanged in both groups. There is a report that serum fetuin-A concentration was increased in patients with secondary hyperparathyroidism after treatment with a VDRA paricalcitol (34). Because fetuin-A is a potent inhibitor of calcification (8) and the key glycoprotein component of calciprotein particles (35), an increase in serum fetuin-A could result in an increase in T<sub>50</sub> (14). However, there was no change in serum fetuin-A level in either the maxacalcitol group or the etelcalcetide group.

Potential roles of vitamin D in muscular health have been suggested by experimental studies (19) and observational



**Table 3. Other subsidiary efficacy outcomes**

Outcomes	Etelcalcetide Group	Maxacalcitol Group	P Value
Change in T <sub>50</sub> from 0 to 3 mo, min	16 (6 to 26)	−11 (−21 to −0)	<0.001
Change in T <sub>50</sub> from 0 to 6 mo, min	18 (6 to 31)	−16 (−28 to −3)	<0.001
<b>Corrected Ca in target range, mo, n/N (%)</b>			
3	75/149 (50)	92/148 (62)	0.06
6	91/148 (62)	90/141 (64)	0.94
12	95/142 (67)	83/132 (63)	0.49
<b>P in target range, mo, n/N (%)</b>			
3	110/149 (74)	104/148 (70)	0.66
6	117/148 (79)	90/141 (64)	0.002
12	101/142 (71)	81/132 (61)	0.08
<b>Intact PTH in target range, mo, n/N (%)</b>			
3	53/149 (36)	80/148 (54)	<0.001
6	71/148 (48)	76/141 (54)	0.17
12	74/142 (52)	74/132 (56)	0.39
<b>Ca, P, and intact PTH all in target range, mo, n/N (%)</b>			
3	18/149 (12)	41/148 (28)	<0.001
6	38/148 (26)	30/141 (21)	0.36
12	37/142 (26)	33/132 (25)	0.87
<b>&gt;30% reduction in intact PTH, mo, n/N (%)</b>			
3	73/149 (49)	121/148 (82)	<0.001
6	101/148 (68)	109/141 (77)	0.04
12	102/142 (72)	107/132 (81)	0.04
<b>Ca × P product, mo, mg<sup>2</sup>/dl<sup>2</sup></b>			
3	43 (40 to 45)	54 (51 to 56)	<0.001
6	41 (39 to 44)	54 (51 to 56)	<0.001
12	45 (42 to 47)	54 (51 to 56)	<0.001
<b>Fetuin-A, mo, μg/ml</b>			
3	243 (230 to 256)	260 (247 to 273)	0.01
6	246 (233 to 258)	241 (229 to 254)	0.50
12	259 (247 to 270)	261 (250 to 273)	0.63
<b>Intact FGF23, mo, pg/ml</b>			
3	850 (661–11,093)	3312 (2578–4256)	<0.001
6	721 (556–934)	4729 (3652–6125)	<0.001
12	836 (629–1109)	4853 (3660–6434)	<0.001

Changes in T<sub>50</sub> and other laboratory data excluding intact FGF23 are shown as least square means (95% confidence intervals). For intact FGF23, these values were calculated after logarithmic transformation, and the table gives the values by exponential retransformation. Laboratory data in target ranges are given in number of patients in target range (*n*), number of patients at each visit (*N*), and percentage in parentheses. Statistical analysis was done by adjusting for age, sex, and P level at baseline. The target ranges recommended by the clinical practice guideline by the Japanese Society for Dialysis Therapy (2012) were corrected C between 8.4 and 10.0 mg/dl, P between 3.5 and 6.0 mg/dl, and intact PTH between 60 and 240 pg/ml (24). The changes in T<sub>50</sub> from 0 to 3 months and from 0 to 6 months were calculated with missing values being imputed by the last observation carried forward method. The numbers of patients in the target ranges are shown by the numbers of patients at each visit without imputation. All tests for between-group comparison were adjusted for age and serum P at baseline. T<sub>50</sub>, serum calcification propensity; Ca, calcium; 3, visit at 3 months; 6, visit at 6 months; 12, visit at 12 months; P, phosphate; PTH, parathyroid hormone; FGF23, fibroblast growth factor-23.

human studies (20,36). However, this trial showed no effect of maxacalcitol or etelcalcetide on handgrip strength. This is consistent with the study by Westerberg *et al.* (37) that showed no effect of high-dose cholecalciferol on handgrip strength in patients with secondary hyperparathyroidism and CKD stages 3 and 4. A longer study period might be needed. Previous observational studies might have been confounded by some factors, such as higher sunshine exposure in individuals with higher physical activity. Vitamin D action may be more important in muscle development in early stages than the maintenance of muscular function in later stages of life.

There was no change in DASC-21 score in either group, and there was no intergroup difference in the changes in DASC-21. However, the apparently neutral effects on DASC-21 should be interpreted carefully. The very low baseline scores of DASC-21 indicate almost full performance of cognition and activity of daily living levels, which

made it difficult to detect improvement in these outcomes, if any, by the so-called ceiling effect. Thus, the results of this study regarding cognition and activity of daily living are inconclusive.

How can we translate our finding on T<sub>50</sub> into hard clinical end points? According to the *post hoc* analysis of the EVOLVE trial (15) in patients with secondary hyperparathyroidism undergoing hemodialysis, a shorter T<sub>50</sub> value was associated with higher risks of all-cause mortality and myocardial infarction, with hazard ratios of 1.10 (95% CI, 1.02 to 1.17) and 1.38 (95% CI, 1.19 to 1.60) per one SD of T<sub>50</sub>, respectively. According to another study in patients with CKD predialysis (14), a longitudinal increase of aortic pulse wave velocity was inversely associated with serum T<sub>50</sub> at baseline, with an odds ratio of 0.52 (95% CI, 0.31 to 0.85) per one SD of T<sub>50</sub>. In this trial, the observed between-group difference of the change in T<sub>50</sub> during 12 months of treatment (20 minutes) corresponds to 0.36 SD of T<sub>50</sub> at

**Table 4. Adverse events in safety analysis set**

Adverse Events	Etelcalcetide Group, <i>n</i> =165	Maxacalcitol Group, <i>n</i> =159
<b>Serious adverse events</b>		
Any serious adverse events	41 (32; 19%)	70 (44; 28%)
All-cause death	5 (5; 3%)	6 (6; 4%)
Any hospitalization	38 (30; 18%)	65 (40; 25%)
Cardiac events	8 (5; 3%)	4 (4; 3%)
Cerebrovascular events	1 (1; 1%)	6 (6; 4%)
Peripheral vascular event	9 (9; 5%)	14 (10; 6%)
Infectious disease	3 (3; 2%)	20 (14; 9%)
Malignancy	3 (3; 2%)	10 (6; 4%)
Other serious adverse events	17 (15; 9%)	16 (13; 8%)
<b>Symptoms</b>		
Nausea	11 (9; 5%)	24 (17; 11%)
Vomiting	10 (8; 5%)	16 (12; 8%)
Dysgeusia/cacogeusia	7 (4; 2%)	5 (3; 2%)
Diarrhea	22 (16; 10%)	10 (7; 4%)
Itching	58 (30; 18%)	72 (38; 24%)
Irritability/jitteriness	20 (15; 9%)	24 (16; 10%)
Falls	47 (28; 17%)	38 (27; 17%)
Hypercalcemia requiring temporal cessation of study drug	0 (0; 0%)	10 (10; 6%)
Hypocalcemia requiring temporal cessation of study drug	33 (30; 18%)	0 (0; 0%)
The table gives numbers of event (number of patients; percentage of participants) by group. Some patients had repeated adverse events in the same category.		

baseline. Thus, it may be translated into modest effects on hard clinical end points and aortic stiffness. Although we had collected data on SAEs, including mortality and hospitalization, for safety assessment, the follow-up period for 12 months may not be long enough. Clearly, we need further studies to directly address this issue.

The KDIGO clinical practice guideline does not prioritize calcimimetics or VDRA in the treatment for secondary hyperparathyroidism (5). In choosing a treatment, considerations may be important regarding not only the effect on PTH but also influence on outcomes such as cardiovascular disease, calcification stress, sarcopenia, and cognition. Previous randomized trials in CKD failed to show the beneficial effects of VDRA on left ventricular mass (38,39) or cardiovascular events (40) as compared with placebo or standard care. This study provides novel evidence that treatment with etelcalcetide had a greater effect of calcification propensity reduction than treatment with maxacalcitol, whereas no difference was found on strength or cognition. Numerically fewer patients in the etelcalcetide group experienced SAEs, which may affect the choice to some extent.

There are several limitations in this study. First, this study did not compare the effects of the two study drugs on clinical hard end points as the primary outcome as discussed above. Second, this randomized trial was not a double-blinded one. However, this hardly affected the result on the primary efficacy outcome of  $T_{50}$  because  $T_{50}$  was measured by persons who were masked to the information on assigned treatment. Third, the results of this study in Japan may not be applicable to patients in other countries where background characteristics of patients and the target ranges of intact PTH are different. Fourth, there was temporal cessation of etelcalcetide in 30 patients and use of oral alfacalcidol in 44 patients in the etelcalcetide group due to hypocalcemia, whereas ten patients skipped

maxacalcitol and 17 patients received oral cinacalcet or evocalcet in the maxacalcitol group due to hypercalcemia. This “contamination” of treatments may underestimate the true effects of these drugs. Fifth, this study was underpowered because only 326 patients were enrolled as compared with the target number of 400. Sixth, the  $T_{50}$  values in this trial were much lower than those reported for dialysis-dependent and -independent patients with CKD from Western countries. Although the reason was unknown, it hardly affects the conclusion. On the other hand, the strengths of this study include the randomized design, the 12 months of follow-up, and the use of intravenously administered drugs as study drugs that prohibited the issue of nonadherence.

In conclusion, this study showed that treatment with etelcalcetide was more effective in increasing  $T_{50}$ , namely in suppressing calcification propensity, than treatment with maxacalcitol in patients with secondary hyperparathyroidism undergoing hemodialysis. No difference was found on the effects on handgrip strength or DASC-21 between the two groups. These results provide novel evidence in the treatment of secondary hyperparathyroidism. Further studies are needed to confirm these results in other settings and to examine the potential effects on hard clinical outcomes.

#### Disclosures

M. Emoto reports receiving grants and personal fees from Medtronic, Nippon Boehringer Ingelheim, Novo Nordisk Pharma, Ono Pharmaceutical, Sumitomo Dainippon Pharma, Taisho Pharma, and Takeda Pharmaceutical; receiving grants from Roche DC Japan and Teijin Pharma limited; receiving research funding from Chugai Pharmaceutical, Eli Lilly, Fuji Pharmaceutical, Kissei Pharmaceutical, Kowa, Kyowa Kirin, Mitsubishi Tanabe Pharma, Mochida Pharmaceutical, Sanofi, and Torii Pharmaceutical; receiving personal fees from Bristol-Myers Squibb, Chugai Pharmaceutical,

Daiichi Sankyo Pharma Development, Eli Lilly, Johnson & Johnson, Kissei Pharmaceutical, Kowa, Kyowa Kirin, Mitsubishi Tanabe, Nikkiso, Nipro, Otsuka Pharmaceutical, Sanofi, Sanwa Kagaku Kenkyusho, and Terumo; and receiving personal fees from Novartis outside the submitted work. T. Hamada reports employment with Malie Medical Clinic, personal fees from Kyowa Kirin, and personal fees from Torii Pharmaceutical outside the submitted work. M. Imanishi reports employment with Ishikiriseiki Hospital; receiving personal fees from Astellas Pharma Inc., Baxter Limited, Bayer Yakuhin, Chugai Pharmaceutical, Fuso Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Mitsubishi Tanabe Pharma, MSD, Nordisk Pharma, Novartis, Novo, Ono Pharmaceutical, Otsuka Pharmaceutical, Sanwa Kagaku Kenkyusho, Takeda Pharmaceutical, and Torii Pharmaceutical; and receiving personal fees from Eli Lilly outside the submitted work. M. Inaba reports receiving grants and personal fees from Bayer Yakuhin, Chugai Pharmaceutical, Kyowa Kirin, and Ono Pharmaceutical; receiving grants from Asahi Kasei Pharmaceutical, Novartis, Roche DC Japan, Sumitomo Dainippon Pharma, Takeda Pharmaceutical, and Teijin Pharma Limited; receiving personal fees from Daiichi Sankyo Co., Ltd., Kissei Pharmaceutical, Kyowa Kirin Co., Ltd., Pfizer, and Torii Pharmaceutical; and receiving personal fees from Daiichi Sankyo Pharma Development outside the submitted work. E. Ishimura reports receiving grants and personal fees from Daiichi Sankyo Pharma Development; receiving personal fees from Astellas Pharma Inc., Bayer Yakuhin, Chugai Pharmaceutical, Fuso Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Nipro, Novo Nordisk Pharma, Ono Pharmaceutical, Takeda Pharmaceutical, and Torii Pharmaceutical; and receiving personal fees from Sanofi outside the submitted work. D. Kabata reports receiving grants and personal fees from Kyowa Kirin; grants from Daiichi Sankyo Pharma Development; personal fees from Chugai Pharmaceutical; and personal fees from Bayer Yakuhin outside the submitted work. D. Kabata also reports that the Department of Medical Statistics of Osaka City University received the analysis and consultation fee from the Department of Vascular Medicine of Osaka City University. Y. Kato reports receiving personal fees from Kissei Pharmaceutical and receiving personal fees from Torii Pharmaceutical outside the submitted work. Y. Kumeda reports receiving research funding from Kyowa Kirin and Ono Pharmaceutical; personal fees from Ono Pharmaceutical; and personal fees from Kyowa Kirin outside the submitted work. T. Matsumura reports receiving personal fees from Mitsubishi Tanabe Pharma and Ono Pharmaceutical and receiving personal fees from Nippon Boehringer Ingelheim outside the submitted work. K. Mori reports receiving grants and personal fees from Torii Pharmaceutical; receiving personal fees from Bayer Yakuhin, Chugai Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Ono Pharmaceutical, and Takeda Pharmaceutical outside the submitted work; receiving personal fees from Astellas Pharma Inc. outside the submitted work; receiving research funding from Mitsubishi Tanabe Pharma Corporation; and speakers bureau for Astellas Pharma Inc., Bayer Yakuhin, Chugai Pharmaceutical, Daiichi Sankyo Company, Eli Lilly Japan K.K., Kissei Pharmaceutical, Kyowa Kirin, Mitsubishi Tanabe Pharma Corporation, Nippon Boehringer Ingelheim, Ono Pharmaceutical, and Takeda Pharmaceutical. S. Nakatani reports receiving personal fees from Astellas Pharma Inc., Bayer Yakuhin, Chugai Pharmaceutical, Kenkyusho, Kyowa Kirin, MSD, Ono Pharmaceutical, Sanwa Kagaku, and Torii Pharmaceutical and personal fees from Otsuka Pharmaceutical outside the submitted work. T. Nakatani reports receiving grants and personal fees from Astellas Pharma Inc., Chugai Pharmaceutical, Kissei Pharmaceutical, Nippon Shinyaku, Ono Pharmaceutical, Pfizer, Sanofi, Takeda

Pharmaceutical, Teijin Pharma limited, and Torii Pharmaceutical; receiving grants from Taiho Pharmaceutical; receiving personal fees from Asahi Kasei, AstraZeneca, Bayer Yakuhin, Bristol-Myers Squibb, Daiichi Sankyo Pharma Development, Janssen Pharmaceutical, Kyowa Kirin, MSD, Novartis, and Taisho Pharma; and receiving personal fees from Tsumura outside the submitted work. S. Nishi reports receiving grants and personal fees from Astellas Pharma Inc., Bayer Yakuhin, Chugai Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Ono Pharmaceutical, Takeda Pharmaceutical, and Torii Pharmaceutical; receiving personal fees from Fuso Pharmaceutical and Mylan; and receiving personal fees from Sanwa Kagaku Kenkyusho outside the submitted work. Y. Ohno reports receiving personal fees from Chugai Pharmaceutical, Daiichi Sankyo Pharma Development, Kissei Pharmaceutical, Kyowa Kirin, Mitsubishi Tanabe Pharma, and Takeda Pharmaceutical and receiving personal fees from Kowa outside the submitted work. Y. Sai reports employment with Suminodo Clinic; consultancy agreements with Kissei Pharmaceutical; and receiving personal fees from Kissei Pharmaceutical outside the submitted work. S. Sasaki reports receiving personal fees from AbbVie, Astellas Pharma Inc., Kyowa Kirin, Mitsubishi Tanabe Pharma, Ono Pharmaceutical, Sumitomo Dainippon Pharma, Taisho Pharma, and Takeda Pharmaceutical and receiving personal fees from Nippon Boehringer Ingelheim outside the submitted work. A. Shintani reports receiving grants and personal fees from Kyowa Kirin and Takeda Pharmaceutical; receiving personal fees from Bayer Yakuhin, Chugai Pharmaceutical, Nipro, and Ono Pharmaceutical; and receiving personal fees from Torii Pharmaceutical outside the submitted work. S. Shoji reports receiving personal fees from Astellas Pharma Inc., AstraZeneca, Bayer Yakuhin, Chugai Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Mitsubishi Tanabe Pharma, Mochida Pharmaceutical, Nippon Boehringer Ingelheim, Novo Nordisk Pharma, Ono Pharmaceutical, Otsuka Pharmaceutical, Sanwa Kagaku Kenkyusho, Taisho Pharma, Teijin Pharma Limited, and Torii Pharmaceutical and receiving personal fees from Toa Eiyo outside the submitted work. T. Shoji reports consultancy agreements with Sanwa Kagaku Kenkyusho; receiving grants and personal fees from Bayer Yakuhin; receiving personal fees from Astellas Pharma Inc., Chugai Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Ono Pharmaceutical, Sanwa Kagaku Kenkyusho, and Torii Pharmaceutical outside the submitted work; receiving personal fees from Teijin Pharma limited outside the submitted work; serving on the editorial boards of *CJASN* and *Renal Replacement Therapy*; and serving as associate editor of the *Journal of Atherosclerosis and Thrombosis*. Y. Tsujimoto reports employment with Inoue Hospital; receiving personal fees from Bayer Yakuhin, Chugai Pharmaceutical, Kissei Pharmaceutical, Kyowa Kirin, Ono Pharmaceutical, and Torii Pharmaceutical; and receiving personal fees from Astellas Pharma Inc. outside the submitted work. S. Ueda reports receiving grants and personal fees from Astellas Pharma Inc., Bayer Yakuhin, Chugai Pharmaceutical, Daiichi Sankyo Pharma Development, MSD, and Takeda Pharmaceutical; receiving grants from Bristol-Myers Squibb and Kowa; receiving personal fees from Bristol-Myers Squibb, Kowa, Kyowa Kirin, and Ono Pharmaceutical; and receiving personal fees from Pfizer outside the submitted work. K. Yamakawa reports employment with Shirasagi Hospital. H. Yasuda reports receiving personal fees from Kissei Pharmaceutical and Kyowa Kirin and receiving personal fees from Torii Pharmaceutical outside the submitted work. H. Yoshida reports receiving personal fees from Sumitomo Dainippon Pharma and receiving personal fees from Baxter Limited outside the submitted work. M. Yoshiyama reports receiving grants and personal fees from Astellas Pharma Inc., Bayer Yakuhin, Daiichi

Sankyo Pharma Development, MSD, Nippon Boehringer Ingelheim, Ono Pharmaceutical, Otsuka Pharmaceutical, Pfizer Sumitomo Dainippon Pharma, Takeda Pharmaceutical, Teijin Pharma Limited, and Toray Medical; receiving grants from Biotronik Japan, Boston Scientific, Edwards Lifesciences, Goodman, Japan Lifeline, Kowa, Sanofi, and Terumo; receiving personal fees from Chugai Pharmaceutical, Kyowa Kirin, Mylan, Sanwa, and Torii Pharmaceutical; and receiving personal fees from Kissei Pharmaceutical outside the submitted work. All remaining authors have nothing to disclose.

### Funding

This study was funded by a grant from Ono Pharmaceutical (to T. Shoji).

### Acknowledgments

The authors acknowledge valuable contributions of the following collaborators: Ms. Masayo Sasagawa at the Department of Metabolism, Endocrinology and Molecular Medicine; Dr. Takahisa Terada at Kitatatsumi Shirasagi Clinic; Dr. Hideki Masaki at Hirano Shirasagi Clinic; Ms. Yumi Ikehara at the University of the Ryukyus Hospital; Dr. Yoshie Fukunaga at Osaka University Hospital; Dr. Eriko Komiya at Osaka University Hospital; and Ms. Ryoko Yagi, Ms. Junko Matsumura, Ms. Mariko Oka, Ms. Yuka Matsumoto, Ms. Yuri Matsumoto, Ms. Masayo Kitano, and Ms. Toshiko Seki at the Center for Clinical Research and Innovation, Osaka University Hospital. The authors thank the patients who participated in this study.

The results of this study were presented at the High-Impact Clinical Trials Symposium during the 65th Annual Meeting of Japanese Society for Dialysis Therapy in Osaka, Japan (November 2, 2020; web version), and the abstract was published in Japanese.

The funder played no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

### Data Sharing Statement

Individual participant data that underline the results reported in this article (text, tables, figures, and appendices) after de-identification, study protocol, and statistical analysis plan will be shared with researchers who provide a methodologically sound proposal to achieve aims in the approved proposal. Data sharing will be available in a period beginning 3 months and ending 5 years following article publication. Proposals should be directed to t-shoji@med.osaka-cu.ac.jp.

### Supplemental Material

This article contains the following supplemental material online at <http://cjasn.asnjournals.org/lookup/suppl/doi:10.2215/CJN.16601020/-/DCSupplemental>.

Supplemental Material. Study protocol of the VICTORY trial and statistical analysis plan.

Supplemental Table 1. Inclusion and exclusion criteria.

Supplemental Table 2. Assay method of serum calcification propensity ( $T_{50}$ ).

Supplemental Table 3. Methods for handgrip strength and DASC-21 measurements.

Supplemental Table 4. List of primary, secondary, tertiary, and other subsidiary outcomes.

Supplemental Table 5. Additional efficacy analysis with per-protocol set.

Supplemental Figure 1. Synopsis of this trial plan.

### References

- Moe S, Drüeke T, Cunningham J, Goodman W, Martin K, Olgaard K, Ott S, Sprague S, Lameire N, Eknoyan G; Kidney Disease: Improving Global Outcomes (KDIGO): Definition, evaluation, and classification of renal osteodystrophy: A position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 69: 1945–1953, 2006
- Wanner C, Amann K, Shoji T: The heart and vascular system in dialysis. *Lancet* 388: 276–284, 2016
- Block GA, Klassen PS, Lazarus JM, Ofsthun N, Lowrie EG, Chertow GM: Mineral metabolism, mortality, and morbidity in maintenance hemodialysis. *J Am Soc Nephrol* 15: 2208–2218, 2004
- Taniguchi M, Fukagawa M, Fujii N, Hamano T, Shoji T, Yokoyama K, Nakai S, Shigematsu T, Iseki K, Tsubakihara Y; Committee of Renal Data Registry of the Japanese Society for Dialysis Therapy: Serum phosphate and calcium should be primarily and consistently controlled in prevalent hemodialysis patients. *Ther Apher Dial* 17: 221–228, 2013
- Ketteler M, Block GA, Evenepoel P, Fukagawa M, Herzog CA, McCann L, Moe SM, Shroff R, Tonelli MA, Toussaint ND, Vervloet MG, Leonard MB: Executive summary of the 2017 KDIGO Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD) guideline update: What's changed and why it matters. *Kidney Int* 92: 26–36, 2017
- Wetmore JB, Gurevich K, Sprague S, Da Roza G, Buerkert J, Reiner M, Goodman W, Cooper K: A randomized trial of cinacalcet versus vitamin D analogs as monotherapy in secondary hyperparathyroidism (PARADIGM). *Clin J Am Soc Nephrol* 10: 1031–1040, 2015
- Jono S, McKee MD, Murry CE, Shioi A, Nishizawa Y, Mori K, Mori H, Giachelli CM: Phosphate regulation of vascular smooth muscle cell calcification. *Circ Res* 87: E10–E17, 2000
- Heiss A, DuChesne A, Denecke B, Grötzinger J, Yamamoto K, Renné T, Jahn-Dechent W: Structural basis of calcification inhibition by alpha 2-HS glycoprotein/fetuin-A. Formation of colloidal calciprotein particles. *J Biol Chem* 278: 13333–13341, 2003
- Heiss A, Jahn-Dechent W, Endo H, Schwahn D: Structural dynamics of a colloidal protein-mineral complex bestowing on calcium phosphate a high solubility in biological fluids. *Bio-interphases* 2: 16–20, 2007
- Aghagolzadeh P, Radpour R, Bachtler M, van Goor H, Smith ER, Lister A, Odermatt A, Feelisch M, Pasch A: Hydrogen sulfide attenuates calcification of vascular smooth muscle cells via KEAP1/NRF2/NQO1 activation. *Atherosclerosis* 265: 78–86, 2017
- Pasch A, Farese S, Gräber S, Wald J, Richterling W, Floege J, Jahn-Dechent W: Nanoparticle-based test measures overall propensity for calcification in serum. *J Am Soc Nephrol* 23: 1744–1752, 2012
- Bundy JD, Cai X, Scialla JJ, Dobre MA, Chen J, Hsu CY, Leonard MB, Go AS, Rao PS, Lash JP, Townsend RR, Feldman HI, de Boer IH, Block GA, Wolf M, Smith ER, Pasch A, Isakova T; CRIC Study Investigators: Serum calcification propensity and coronary artery calcification among patients with CKD: The CRIC (Chronic Renal Insufficiency Cohort) study. *Am J Kidney Dis* 73: 806–814, 2019
- Smith ER, Ford ML, Tomlinson LA, Rajkumar C, McMahon LP, Holt SG: Phosphorylated fetuin-A-containing calciprotein particles are associated with aortic stiffness and a procalcific milieu in patients with pre-dialysis CKD. *Nephrol Dial Transplant* 27: 1957–1966, 2012
- Smith ER, Ford ML, Tomlinson LA, Bodenham E, McMahon LP, Farese S, Rajkumar C, Holt SG, Pasch A: Serum calcification propensity predicts all-cause mortality in predialysis CKD. *J Am Soc Nephrol* 25: 339–348, 2014
- Pasch A, Block GA, Bachtler M, Smith ER, Jahn-Dechent W, Arampatzis S, Chertow GM, Parfrey P, Ma X, Floege J: Blood calcification propensity, cardiovascular events, and survival in patients receiving hemodialysis in the EVOLVE trial. *Clin J Am Soc Nephrol* 12: 315–322, 2017
- Keyzer CA, de Borst MH, van den Berg E, Jahn-Dechent W, Arampatzis S, Farese S, Bergmann IP, Floege J, Navis G, Bakker SJ, van Goor H, Eisenberger U, Pasch A: Calcification propensity and survival among renal transplant recipients. *J Am Soc Nephrol* 27: 239–248, 2016

17. Maggio M, Lauretani F: Prevalence, incidence, and clinical impact of cognitive-motoric risk syndrome in Europe, USA, and Japan: Facts and numbers update 2019. *J Cachexia Sarcopenia Muscle* 10: 953–955, 2019
18. Molina P, Carrero JJ, Bover J, Chauveau P, Mazzaferro S, Torres PU; European Renal Nutrition (ERN) and Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD) Working Groups of the European Renal Association-European Dialysis Transplant Association (ERA-EDTA): Vitamin D, a modulator of musculoskeletal health in chronic kidney disease. *J Cachexia Sarcopenia Muscle* 8: 686–701, 2017
19. Endo I, Inoue D, Mitsui T, Umaki Y, Akaike M, Yoshizawa T, Kato S, Matsumoto T: Deletion of vitamin D receptor gene in mice results in abnormal skeletal muscle development with deregulated expression of myoregulatory transcription factors. *Endocrinology* 144: 5138–5144, 2003
20. Wicherts IS, van Schoor NM, Boeke AJ, Visser M, Deeg DJ, Smit J, Knol DL, Lips P: Vitamin D status predicts physical performance and its decline in older persons. *J Clin Endocrinol Metab* 92: 2058–2065, 2007
21. Miller JW, Harvey DJ, Beckett LA, Green R, Farias ST, Reed BR, Olichney JM, Mungas DM, DeCarli C: Vitamin D status and rates of cognitive decline in a multiethnic cohort of older adults. *JAMA Neurol* 72: 1295–1303, 2015
22. Gordon PL, Sakkas GK, Doyle JW, Shubert T, Johansen KL: Relationship between vitamin D and muscle size and strength in patients on hemodialysis. *J Ren Nutr* 17: 397–407, 2007
23. Zahed N, Chehrizi S, Falaknasi K: The evaluation of relationship between vitamin D and muscle power by micro manual muscle tester in end-stage renal disease patients. *Saudi J Kidney Dis Transpl* 25: 998–1003, 2014
24. Fukagawa M, Yokoyama K, Koiwa F, Taniguchi M, Shoji T, Kazama JJ, Komaba H, Ando R, Kakuta T, Fujii H, Nakayama M, Shibagaki Y, Fukumoto S, Fujii N, Hattori M, Ashida A, Iseki K, Shigematsu T, Tsukamoto Y, Tsubakihara Y, Tomo T, Hirakata H, Akizawa T; CKD-MBD Guideline Working Group; Japanese Society for Dialysis Therapy: Clinical practice guideline for the management of chronic kidney disease-mineral and bone disorder. *Ther Apher Dial* 17: 247–288, 2013
25. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG: Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 42: 377–381, 2009
26. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, McLeod L, Delacqua G, Delacqua F, Kirby J, Duda SN; REDCap Consortium: The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 95: 103208, 2019
27. Dam M, Neelemaat F, Struijk-Wieling T, Weijs PJ, van Jaarsveld BC: Physical performance and protein-energy wasting in patients treated with nocturnal haemodialysis compared to conventional haemodialysis: Protocol of the DiapriFIT study. *BMC Nephrol* 18: 144, 2017
28. Awata S, Sugiyama M, Ito K, Ura C, Miyamae F, Sakuma N, Niikawa H, Okamura T, Inagaki H, Ijuin M: Development of the dementia assessment sheet for community-based integrated care system. *Geriatr Gerontol Int* 16[Suppl 1]: 123–131, 2016
29. Nakatani S, Mori K, Sonoda M, Nishide K, Uedono H, Tsuda A, Emoto M, Shoji T: Association between serum zinc and calcification propensity (T<sub>50</sub>) in patients with type 2 diabetes mellitus and *in vitro* effect of exogenous zinc on T<sub>50</sub>. *Biomedicines* 8: 337, 2020
30. Block GA, Bushinsky DA, Cunningham J, Druke TB, Ketteler M, Kewalramani R, Martin KJ, Mix TC, Moe SM, Patel UD, Silver J, Spiegel DM, Sterling L, Walsh L, Chertow GM: Effect of etelcalcetide vs placebo on serum parathyroid hormone in patients receiving hemodialysis with secondary hyperparathyroidism: Two randomized clinical trials. *JAMA* 317: 146–155, 2017
31. Hayashi M, Tsuchiya Y, Itaya Y, Takenaka T, Kobayashi K, Yoshizawa M, Nakamura R, Monkawa T, Ichihara A: Comparison of the effects of calcitriol and maxacalcitol on secondary hyperparathyroidism in patients on chronic haemodialysis: A randomized prospective multicentre trial. *Nephrol Dial Transplant* 19: 2067–2073, 2004
32. Bressendorff I, Hansen D, Schou M, Pasch A, Brandt L: The effect of increasing dialysate magnesium on serum calcification propensity in subjects with end stage kidney disease: A randomized, controlled clinical trial. *Clin J Am Soc Nephrol* 13: 1373–1380, 2018
33. Bressendorff I, Hansen D, Schou M, Silver B, Pasch A, Bouchelouche P, Pedersen L, Rasmussen LM, Brandt L: Oral magnesium supplementation in chronic kidney disease stages 3 and 4: Efficacy, safety, and effect on serum calcification propensity—A prospective randomized double-blinded placebo-controlled clinical trial. *Kidney Int Rep* 2: 380–389, 2016
34. Manenti L, Vaglio A, Pasquali S: Increased fetuin-A levels following treatment with a vitamin D analog. *Kidney Int* 78: 1187, 2010
35. Jähnen-Dechent W, Heiss A, Schäfer C, Ketteler M: Fetuin-A regulation of calcified matrix metabolism. *Circ Res* 108: 1494–1509, 2011
36. Visser M, Deeg DJ, Lips P; Longitudinal Aging Study Amsterdam: Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): The longitudinal aging study Amsterdam. *J Clin Endocrinol Metab* 88: 5766–5772, 2003
37. Westerberg PA, Sterner G, Ljunggren Ö, Isaksson E, Elvarson F, Dezfolian H, Linde T: High doses of cholecalciferol alleviate the progression of hyperparathyroidism in patients with CKD stages 3–4: Results of a 12-week double-blind, randomized, controlled study. *Nephrol Dial Transplant* 33: 466–471, 2018
38. Thadhani R, Appelbaum E, Chang Y, Pritchett Y, Bhan I, Agarwal R, Zoccali C, Wanner C, Lloyd-Jones D, Cannata J, Thompson T, Audhya P, Andress D, Zhang W, Ye J, Packham D, Singh B, Zehnder D, Manning WJ, Pachika A, Solomon SD: Vitamin D receptor activation and left ventricular hypertrophy in advanced kidney disease. *Am J Nephrol* 33: 139–149, 2011
39. Wang AY, Fang F, Chan J, Wen YY, Qing S, Chan IH, Lo G, Lai KN, Lo WK, Lam CW, Yu CM: Effect of paricalcitol on left ventricular mass and function in CKD—The OPERA trial. *J Am Soc Nephrol* 25: 175–186, 2014
40. Shoji T, Inaba M, Fukagawa M, Ando R, Emoto M, Fujii H, Fujimori A, Fukui M, Hase H, Hashimoto T, Hirakata H, Honda H, Hosoya T, Ikari Y, Inaguma D, Inoue T, Isaka Y, Iseki K, Ishimura E, Itami N, Ito C, Kakuta T, Kawai T, Kawanishi H, Kobayashi S, Kumagai J, Maekawa K, Masakane I, Minakuchi J, Mitsuiki K, Mizuguchi T, Morimoto S, Murohara T, Nakatani T, Negi S, Nishi S, Nishikawa M, Ogawa T, Ohta K, Ohtake T, Okamura M, Okuno S, Shigematsu T, Sugimoto T, Suzuki M, Tahara H, Takemoto Y, Tanaka K, Tominaga Y, Tsubakihara Y, Tsujimoto Y, Tsuruya K, Ueda S, Watanabe Y, Yamagata K, Yamakawa T, Yano S, Yokoyama K, Yorioka N, Yoshiyama M, Nishizawa Y; J-DAVID Investigators: Effect of oral alfacalcidol on clinical outcomes in patients without secondary hyperparathyroidism receiving maintenance hemodialysis: The J-DAVID randomized clinical trial. *JAMA* 320: 2325–2334, 2018

Received: October 22, 2020 Accepted: January 22, 2021

Published online ahead of print. Publication date available at [www.cjasn.org](http://www.cjasn.org).

## AFFILIATIONS

- <sup>1</sup>Department of Vascular Medicine, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>2</sup>Vascular Science Center for Translational Research, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>3</sup>Department of Metabolism, Endocrinology and Molecular Medicine, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>4</sup>Department of Medical Statistics, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>5</sup>Department of Nephrology, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>6</sup>Center for Clinical Research and Innovation, Osaka City University Hospital, Osaka, Japan
- <sup>7</sup>Department of Drug and Food Evaluation, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>8</sup>Department of Clinical Pharmacology and Therapeutics, University of the Ryukyus, Okinawa, Japan
- <sup>9</sup>Division of Nephrology and Kidney Center, Kobe University Graduate School of Medicine, Kobe, Japan
- <sup>10</sup>Department of Urology, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>11</sup>Department of Cardiovascular Medicine, Osaka City University Graduate School of Medicine, Osaka, Japan
- <sup>12</sup>Ono Naika Clinic, Osaka, Japan
- <sup>13</sup>Marie Clinic, Higashi Osaka, Japan
- <sup>14</sup>Ishikiri Seiki Hospital, Higashi Osaka, Japan
- <sup>15</sup>Meijibashi Hospital, Matsubara, Japan
- <sup>16</sup>Kawamura Clinic, Osaka, Japan
- <sup>17</sup>Shirasagi Minami Clinic, Osaka, Japan
- <sup>18</sup>Minami Osaka Clinic, Osaka, Japan
- <sup>19</sup>Fujiidera Shirasagi Clinic, Fujiidera, Japan
- <sup>20</sup>Ikeda Hospital, Higashi Osaka, Japan
- <sup>21</sup>Nagayama Clinic, Sennan-gun, Japan
- <sup>22</sup>Obi Clinic, Osaka, Japan
- <sup>23</sup>Ohno Memorial Hospital, Osaka, Japan
- <sup>24</sup>Suminodo Clinic, Daito, Japan
- <sup>25</sup>Jurakukai Clinic, Osaka, Japan
- <sup>26</sup>Sasaki Medical Clinic, Osaka, Japan
- <sup>27</sup>Ikeda Clinic, Higashi Osaka, Japan
- <sup>28</sup>Shirasagi Hospital, Osaka, Japan
- <sup>29</sup>Aijinkai Healthcare Corporation Inoue Hospital, Suita, Japan
- <sup>30</sup>Shirasagi Clinic, Osaka, Japan
- <sup>31</sup>Okada Clinic, Osaka, Japan
- <sup>32</sup>Yodoi Hospital, Osaka, Japan