

Relative Blood Volume Changes Underestimate Total Blood Volume Changes during Hemodialysis

Judith J. Dasselaar,^{*†} Marjolijn N. Lub-de Hooge,^{‡§} Jan Pruijm,[§] Hugo Nijhuis,[§] Anneke Wiersum,^{*} Paul E. de Jong,[†] Roel M. Huisman,^{*†} and Casper F.M. Franssen^{*†}

^{*}Dialysis Center Groningen, [†]Department of Internal Medicine, Division of Nephrology, [‡]Hospital and Clinical Pharmacy, [§]Department of Nuclear Medicine and Molecular Imaging, University Medical Center Groningen and University of Groningen, Groningen, Netherlands

Background: Measurements of relative blood volume changes (Δ RBV) during hemodialysis (HD) are based on hemoconcentration and assume uniform mixing of erythrocytes and plasma throughout the circulation. However, whole-body hematocrit (Ht) is lower than systemic Ht. During HD, a change in the ratio between whole-body to systemic Ht (F cell ratio) is likely to occur as a result of a net shift of low Ht blood from the microcirculation to the macrocirculation. Hence, Δ RBV may differ significantly from total blood volume changes (Δ TBV). Therefore, this study compared Δ RBV and Δ TBV during HD.

Design, setting, participants, and measurements: Plasma and erythrocyte volumes were measured using ¹²⁵I- and ¹²³I-radioiodinated albumin and ⁵¹Cr-labeled erythrocytes, respectively. After validation of the standardized method in two patients on a nondialysis day, seven patients completed the protocol during HD. ¹²⁵I-albumin and ⁵¹Cr-labeled erythrocytes were administered 20 min before the start of HD. ¹²³I-albumin was administered at 160 min into the HD session to quantify and correct for ¹²⁵I-albumin leakage. Δ RBV was measured continuously throughout HD. The F cell ratio was derived from whole-body and systemic Ht.

Results: Total ultrafiltration volume was 2450 ± 770 ml. TBV declined from 5905 ± 824 to 4877 ± 722 ml during HD. Thus, TBV declined $17.3 \pm 4.4\%$, whereas the RBV decline was only $8.2 \pm 3.7\%$ ($P = 0.001$). The F cell ratio increased from 0.896 ± 0.036 to 0.993 ± 0.049 during HD ($P = 0.002$).

Conclusions: Δ RBV significantly underestimates Δ TBV during HD. The rise in F cell ratio strongly suggests that during HD, blood translocates from the microcirculation to the macrocirculation, probably as a cardiovascular compensatory mechanism in response to hypovolemia.

Clin J Am Soc Nephrol 2: 669-674, 2007. doi: 10.2215/CJN.00880207

Technological advances have allowed the development of continuous, noninvasive measurement of relative blood volume changes (Δ RBV) during hemodialysis (HD). Continuous registration of Δ RBV during HD with ultrafiltration (UF) is advocated as a tool to maintain an adequate volume of the intravascular compartment to avoid HD hypotension (1–3). However, the use of Δ RBV measurements in clinical practice depends on its accuracy in reflecting the change in whole-body blood volume or total blood volume (Δ TBV).

Noninvasive measurements of Δ RBV are based on hemoconcentration of blood constituents that remain in the circulation during HD with UF. Mostly erythrocytes are used for these measurements. Δ RBV devices that are based on hemoconcentration of afferent blood can adequately represent Δ TBV only in case of uniform mixing of plasma and erythrocytes throughout the different vascular beds of the circulation (4–6). However,

this assumption is not valid: The whole-body hematocrit (Ht) is lower than the Ht of arterial or venous blood (7,8). The difference is due to a dynamic reduction in microvascular Ht in capillaries and venules ($<200 \mu\text{m}$), known as the Fahraeus effect (9). The difference between arterial or venous Ht and whole-body Ht is expressed as the F cell ratio (the ratio of whole-body Ht to arterial or venous Ht) and approximates 0.91 in nondialysis individuals (10,11). The lack of uniform mixing of erythrocytes throughout the circulation would not induce a divergence between Δ RBV and Δ TBV calculation if the difference in Ht between the different vascular beds would remain constant during HD—in other words, if the F cell ratio would not change (12). However, a change in F cell ratio may occur during HD with UF (13,14). As a consequence, the observed Δ RBV may differ significantly from the Δ TBV (13).

To our knowledge, no studies to date that have compared absolute Δ TBV, which is the sum of erythrocyte and plasma volume, using methods that are generally accepted as the gold standard: Radioiodinated albumin (¹²⁵I) and ⁵¹Cr-labeled erythrocytes (15,16), during HD with UF with its relative counterpart: Δ RBV measurements as measured by hemoconcentration of afferent blood. In this study we compared hemoconcentration-based Δ RBV and Δ TBV measured by repetitive measurement of

Received February 21, 2007. Accepted March 26, 2007.

Published online ahead of print. Publication date available at www.cjasn.org.

Address correspondence to: Dr. Judith J. Dasselaar, Department of Internal Medicine, Division of Nephrology, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, Netherlands. Phone: +31-50-3615497; Fax: +31-50-3615403; E-mail: jj.dasselaar@dcg.umcg.nl

absolute plasma and erythrocyte volumes during HD with UF in eight patients. In addition, we calculated the F cell ratio because a change in this factor may explain a possible discrepancy between Δ RBV and Δ TBV measurements.

Materials and Methods

Patients

The standardized method for determining plasma and erythrocyte volumes (radioiodinated ^{125}I -albumin and ^{51}Cr , respectively) was evaluated for its stability for repeated measurements in two HD patients on a nondialysis day in a prestudy test. Subsequently, eight patients underwent measurements of absolute plasma and erythrocyte volumes during a single HD session according to the same protocol.

To be eligible, patients had to fulfill the following criteria: (1) Non-hypotension-prone male HD patients who were treated with HD for at least 6 mo and had stable values for hemoglobin (>7 mmol/L); (2) unaltered erythropoietin dosage for at least 2 mo; (3) an arteriovenous fistula without recirculation as established by Transonic flow measurements (Transonic Systems, Ithaca, NY); and (4) no residual renal function (diuresis <100 ml/24 h). Exclusion criteria were (1) the absence of informed consent, (2) Recent hemorrhage, and (3) diabetes. The last exclusion criterion was applied because participating patients were not allowed to eat or drink during the study.

Study Protocol

Determination of Erythrocyte and Plasma Volume. Plasma and erythrocyte volume measurements are routinely used to assess erythrocyte and plasma volume in the diagnosis of polycythemias. The standard technique for measuring plasma volume involves the use of ^{125}I -labeled albumin (15,16). For the measurement of the erythrocyte volume, ^{51}Cr -labeled erythrocytes are commonly used (15,16). The principle is based on the so-called "isotope dilution technique." Volumes are calculated from the known dosage administered and the radioactivity concentration measured in whole blood or plasma (15). A single dose of ^{51}Cr -labeled erythrocytes will suffice for the duration of this study because the radioactivity half-life is 28 d.

Because transcappillary albumin leakage occurs in both healthy control subjects and HD patients (17), it is not possible to measure reliably the plasma volume over a longer period of time when using only one radioiodinated marker. We therefore determined plasma volume using two radioiodinated albumin injections. We chose ^{125}I -albumin and ^{123}I -albumin because these isotopes differ in γ -ray energy (27 to 32 and 159 keV, respectively) and, thus, allow separate detection in the γ counter. The administration of the second radioiodinated isotope, 180 min after administration of the first isotope, allows quantification of and correction for the transcappillary escape rate of albumin (TER_{alb}). The quantification of TER_{alb} was based on the first ^{125}I plasma volume measurement and the ^{125}I plasma volume determination after 180 min of HD because at this time point, the plasma volume as assessed simultaneously by ^{125}I and ^{123}I should yield identical results. In this study, we assumed that TER_{alb} was constant during HD in the individual participant. All values for plasma volume displayed in this study are corrected for TER_{alb} .

Radiolabeling. ^{51}Cr labeling of erythrocytes and ^{123}I -albumin labeling were performed under aseptic conditions in the radiopharmacy unit of the Nuclear Medicine Department under supervision of a hospital pharmacist. ^{51}Cr -sodium chromate and ^{123}I -sodium iodide (^{123}I -NaI) were obtained from GE Health (Eindhoven, Netherlands). ^{125}I -human serum albumin (^{125}I -HSA) was obtained from Merck Frosst Canada (Kirkland, QU, Canada).

An extensive description of the used methods is described in detail

elsewhere (15,16). In short, the ^{51}Cr -sodium chromate solution was added to the patients' erythrocytes. The mixture was incubated for 45 to 60 min at a temperature of 37°C. Thereafter, 25 mg of ascorbic acid was added to the labeled erythrocyte suspension. A known amount of the well-mixed labeled erythrocyte suspension was injected together with a known amount of ^{125}I -HSA to the patient.

HSA (Cealb 20%; Sanquin, Amsterdam, Netherlands) was labeled in the radiopharmacy department with ^{123}I -NaI using a standard iodogen (Pierce Biotechnology, Rockford, IL) method. The mixture was purified over a PD-10 column (Sephadex G-25M; Amersham Biosciences AB, Uppsala, Sweden) and overnight dialysis (Slide-A-Lyzer; Pierce Biotechnology). Radiochemical purity was $>98\%$.

Protocol. The study protocol was identical during the prestudy test and the HD study except for the start of HD ($T = 0$) during the latter study. HD commenced 20 min after the first injection with the radiopharmaceuticals. This time interval was chosen to allow adequate distribution of the radioiodinated isotopes (15,16). For avoidance of the widely known influence of postural changes on blood volume (18) and F cell ratio (19,20), patients were placed in a supine position 20 min before blood sampling for erythrocyte labeling with ^{51}Cr and remained in this position until the study protocol was completed (from 100 min before the start of HD until the stop of HD). After completion of the erythrocyte labeling with ^{51}Cr , the patients' own blood supplemented with ^{125}I -albumin was administered. The interval between blood sampling for the labeling of the erythrocytes and the re-administration of the isotopes was 60 min. During the prestudy test and the HD study, ^{123}I was given 180 min after the administration of ^{125}I (160 min into the HD session). Blood was sampled four times at 10-min intervals immediately after both injections that contained the isotope(s). After four blood samples, the interval between two samplings was increased to 20 min. In total, blood was sampled 17 times. At each sampling, a total of 5.5 ml of blood was withdrawn. Each sampling included erythrocyte and plasma volume calculation as well as determination of whole-body and systemic Ht. Because the average Ht in our patient group was 0.40 ± 0.05 at the start of HD, the sampling of a total of 93.5 ml of blood equals an erythrocyte and plasma volume of approximately 37.4 and 56.1 ml, respectively. In the calculations of erythrocyte and plasma volume, we made no corrections for this sampling of blood because this plasma volume is very small in comparison with the expected change in plasma volume and because this erythrocyte volume is negligible compared with the patients' erythrocyte volume. A standard dialysis needle was used for blood sampling. A vein on the nondialysis access side was cannulated for the purpose of administering the radiopharmaceuticals.

Patients were asked to refrain from coffee, tea, alcohol, and tobacco use from the night before the study until completion of the study. With the exception of a glass of water (125 ml) at room temperature after 20 and 200 min into the HD session, patients were not allowed to eat or drink throughout the study to avoid any influence of food intake on blood volume (21). BP and heart rate were measured simultaneously during each time blood was sampled.

The study was approved by the Medical Ethics Committee of the University Medical Center Groningen. Written informed consent was obtained from all participating patients. The study was performed in accordance with the principles of the Declaration of Helsinki and guidelines for Good Clinical Practice.

Dialysis Settings. All eight patients underwent dialysis for 4 h with bicarbonate dialysis in a single session. A low-flux polysulfone hollow-fiber dialyzer (F8; Fresenius Medical Care, Bad Homburg, Germany) was used. Blood flow rates and dialysate flow rates were set at 250 and 500 ml/min, respectively. Dialysate composition was as follows: Sodium 139 mmol/L, potassium 1.0 mmol/L, calcium 1.5

mmol/L, magnesium 0.5 mmol/L, chloride 108 mmol/L, bicarbonate 34 mmol/L, acetate 3.0 mmol/L, and glucose 1.0 g/L. Dialysate temperature was set at 36.0°C. HD was performed with the Integra HD apparatus (Gambro-Hospal, Lyon, France). Δ RBV were measured with Hemoscan (Gambro-Hospal). In addition, we assessed Δ RBV from changes in Ht (Δ RBV-Ht) in afferent blood because this method is considered to be the reference method for determining Δ RBV during HD (5,22).

Calculations

The following calculations were used in this study:

Total blood volume = erythrocyte volume + plasma volume

Whole-body Ht = erythrocyte volume/(erythrocyte volume + plasma volume)

F cell ratio = whole-body Ht/venous Ht

Δ TBV (in %) = $[(TBV_0 - TBV_t)/TBV_0] \times 100$

Δ RBV-Ht (in %) = $[(Ht_0/Ht_t) - 1] \times 100$

in which TBV_0 and Ht_0 , and TBV_t and Ht_t represent TBV and Ht at the start of HD and at a certain moment during HD, respectively.

Statistical Analyses

All data were analyzed using GraphPad Prism version 4.00 for Windows (GraphPad Software, San Diego, CA). Data are presented as mean \pm SD unless stated otherwise. Comparisons were made with a paired *t* test or an unpaired *t* test when appropriate. In addition, Bland-Altman analysis was used for the evaluation of differences between Δ RBV and Δ TBV. From these results, the Pearson correlation coefficient *r* was derived. *P* < 0.05 was considered significant.

Results

Patients

Patient characteristics of the two patients in the prestudy test were as follows: Age 67 and 62 yr; dialysis vintage 2 yr and 9 mo and 13 yr, respectively. Cause of renal failure was hypertension and IgA nephropathy, respectively. Plasma volume was 4204 and 3955 ml at the start of the study and 4266 and 4070 ml at the stop of the study for patients 1 and 2, respectively. Erythrocyte volume was 2086 and 2331 ml at the start of the study and 2092 and 2357 ml at the stop of the study for patients 1 and 2, respectively. The studies in both patients indicated that plasma and erythrocyte volumes were stable over the time frame of the study. The F cell ratio at the start of

the study was 0.87 and 0.88 and at the end of the study was 0.86 and 0.88 for patients 1 and 2, respectively. The TER_{alb} was 3.3%/h in both patients.

Next, eight patients were included in the HD study, and all finished the protocol that consisted of plasma and erythrocyte measurements during HD with UF. The results from one patient were excluded from analysis because the erythrocyte volume was unstable.

Patient characteristics of the seven patients from the HD study were as follows: Mean age 56 ± 20 yr (range 27 to 74 yr). The mean time on dialysis was 3.1 ± 1.9 yr (range 1 to 6 yr). The cause of renal failure was hypertension (*n* = 4), chronic pyelonephritis (*n* = 1), membranoproliferative glomerulonephritis (*n* = 1), and Wegener's granulomatosis (*n* = 1). Pre- and post-HD weight was 74.8 ± 8.6 and 72.4 ± 8 kg, respectively. Pre- and post-HD systolic BP was 134 ± 13 and 130 ± 20 mmHg, respectively. Pre- and post-HD diastolic BP was 77 ± 17 and 73 ± 14 mmHg, respectively. Heart rate was 65 ± 9 before HD and 70 ± 15 bpm after HD. No hypotensive episodes occurred during any of the HD sessions.

Erythrocyte and Plasma Volume

In Table 1, the erythrocyte and plasma volumes are shown for the individual patients throughout the HD session. In addition, the individual data for TBV, UF volume, and TER_{alb} are displayed. No significant changes in erythrocyte volume were found. Plasma volume declined from 3784 ± 648 at the start of HD to 2741 ± 563 ml at the end of HD (*P* = 0.0001). TBV declined from 5905 ± 824 at the start of HD to 4877 ± 722 ml at the end of HD (*P* = 0.0001). Mean total UF volume was 2450 ± 770 ml. Mean TER_{alb} was $9.1 \pm 1.3\%/h$.

Table 2 shows the individual change in both TBV and RBV throughout the HD session. In all patients, Δ RBV underestimated the decline in TBV during the second half of the HD session. Figure 1 shows the mean course of Δ TBV and Δ RBV throughout the study. TBV declined $17.3 \pm 4.4\%$ at the end of HD, whereas the observed RBV decline was only $8.2 \pm 3.7\%$ (*P* = 0.001). The difference between Δ TBV and Δ RBV was significant (*P* < 0.05) from 120 min onward. Δ RBV-Ht yielded identical results to Δ RBV (data not shown). There was consid-

Table 1. EV, PV, TBV, UF, and transcapillary escape rate of albumin in the individual patients^a

Patient	EV (ml)					PV (ml)					TBV (ml)		UF (ml)	TER_{alb} (%/h)
	0 min	60 min	120 min	180 min	240 min	0 min	60 min	120 min	180 min	240 min	0 min	240 min		
1	2312	2396	2189	2480	2413	4836	4710	4367	4193	3873	7148	6286	1800	7.6
2	1539	1618	1540	1582	1612	3376	3115	2897	2785	2601	4915	4213	1200	9.8
3	2102	2252	2127	2372	2032	4338	4172	3560	3380	2941	6440	4973	3030	9.0
4	2064	2099	2250	2197	2174	3888	3367	2995	2781	2451	5951	4625	3520	9.4
5	2330	2423	2255	2407	2352	3698	3385	3065	2731	2435	6028	4788	2660	11.5
6	2573	2257	2434	2327	2366	3479	3212	3122	3137	2763	6052	5129	2320	9.1
7	1924	1901	1937	1963	2008	2875	2714	2471	2291	2121	4799	4128	2620	9.1
Mean	2121	2135	2105	2190	2137	3784	3525	3211	3043	2741	5905	4877	2450	9.1

^a0 min and 240 min denote the start and end of the hemodialysis session, respectively. EV, erythrocyte volume; PV, plasma volume; TBV, total blood volume; TER_{alb} , transcapillary escape rate of albumin; UF, ultrafiltration.

Table 2. Δ TBV and Δ RBV at hourly intervals in the individual patients^a

Patient	Δ TBV (%)				Δ RBV (%)			
	60 min	120 min	180 min	240 min	60 min	120 min	180 min	240 min
1	-0.6	-8.3	-8.9	-12.1	-0.1	-1.9	-2.9	-3.4
2	-3.7	-9.7	-10.0	-14.3	-4.4	-5.6	-5.9	-5.7
3	-0.3	-11.7	-14.5	-22.8	-3.4	-5.5	-7.3	-8.1
4	-8.2	-11.9	-19.1	-22.3	-5.5	-7.6	-12.1	-13.9
5	-3.7	-11.7	-13.7	-20.6	-2.1	-3.9	-5.3	-7.2
6	-9.6	-8.2	-14.1	-15.3	-3.2	-4.4	-5.5	-6.8
7	-3.8	-8.1	-13.1	-14.0	-6.4	-8.9	-10.5	-12.2
Mean	-4.3	-10.0	-13.3	-17.3	-3.6	-5.4	-7.0	-8.2

^a Δ RBV, change in relative blood volume; Δ TBV, change in total blood volume.




Figure 1. The mean course of total blood volume changes (Δ TBV; \circ) and relative blood volume changes (Δ RBV; \bullet) during hemodialysis (HD) of the seven patients. Data are means \pm SD. * $P < 0.05$ versus the start of the HD session.

erable interindividual variation in the magnitude of the difference between Δ RBV and Δ TBV (Figure 2). Overall, bias was -5.2% with a 95% limit of agreement from -13.7 to 3.4% . A significant correlation was found when the difference and average of Δ RBV and Δ TBV were compared: $r = 0.42$ ($P = 0.0002$). The extent of the difference between Δ RBV and Δ TBV did not differ between the patients with a UF volume <2450 ml ($n = 3$; mean -5.3 versus -13.9%) in comparison with the four patients with a UF volume >2450 ml ($n = 4$; mean -10.4 versus -19.9%).

F Cell Ratio

In all patients the F cell ratio rose during HD (Table 3). The difference between the F cell ratio at the start of HD in comparison with the F cell ratio during HD was significant ($P < 0.05$) from 60 min into the HD session onward (Figure 3). The F cell ratio increased from 0.896 ± 0.036 at the start of HD to 0.993 ± 0.049 at the stop of the HD session ($P = 0.002$).

Discussion

This study shows directly that Δ RBV as assessed by hemoconcentration of afferent blood significantly underesti-




Figure 2. Bland-Altman plots comparing Δ RBV and Δ TBV at 60, 120, 180, and 240 min into the HD session. Reference lines indicate mean difference \pm 95% limit of agreement. Correlation between difference and average: $r = 0.42$ ($P = 0.0002$). A difference above zero indicates that Δ RBV overestimates Δ TBV; a difference below zero indicates that Δ RBV underestimates Δ TBV. \bullet , patient 1; \circ , patient 2; \blacktriangledown , patient 3; \blacklozenge , patient 4; $*$, patient 5; \square , patient 6; \triangle , patient 7.

mate the TBV decrease during HD with UF. The discrepancy is probably caused by translocation of "low-Ht" blood from the microcirculation to the macrocirculation as reflected by a significant rise in F cell ratio.

The Δ TBV differed significantly from the change in RBV from 120 min onward. Δ RBV devices that are based on hemoconcentration of afferent blood can represent Δ TBV only when the following two prerequisites are met: First, there should be a constant volume of circulating erythrocytes in the circulation (4–6). This assumption is true during HD in the absence of blood leak and hemolysis (13). Second, there should be uniform mixing of erythrocytes and plasma throughout the circulation. This assumption is not true because Ht in the microcirculation is lower than in the macrocirculation (7,8). The lack of uniform mixing of erythrocytes throughout the circulation would probably not induce a great error in Δ RBV measurements as long as the difference in Ht between the different vascular beds would remain constant during HD—in other words, if the F cell ratio

Table 3. Individual hourly F cell ratios

Patient	F Cell Ratio				
	0 min	60 min	120 min	180 min	240 min
1	0.91	0.92	0.93	0.97	1.05
2	0.84	0.93	0.95	0.96	0.99
3	0.92	0.95	0.96	0.98	0.97
4	0.95	0.97	1.00	0.99	1.00
5	0.90	0.99	0.98	1.06	1.06
6	0.87	0.90	0.93	0.92	0.95
7	0.88	0.90	0.88	0.90	0.93
Mean	0.896	0.938	0.947	0.972	0.993




Figure 3. The mean course of the F cell ratio (○) during HD of all seven patients. *P < 0.05 versus the start of the HD session.

would not change (19,23). This study, however, clearly demonstrates that the F cell ratio increases markedly during HD. We hypothesize that this reflects translocation of blood with a lower Ht from the microcirculation to the macrocirculation, thereby inducing a dilution of the central blood compartment. Therefore, the hemoconcentration of blood that reaches the ΔRBV monitor is not representative of the hemoconcentration of the TBV. As a result, ΔRBV monitors underestimate the real ΔTBV during HD.

Mitra *et al.* (13) indirectly established that the F cell ratio might rise during HD. These authors used Indocyanine green to determine plasma volume. Erythrocyte volume was subsequently calculated from plasma volume and venous Ht. However, the use of calculated erythrocyte volumes necessitates the input of an estimated average F cell ratio because an individual F cell ratio cannot be derived from plasma volume alone. This may induce an error because population values for the F cell ratio have been reported as 0.864 (24) and 0.91 (25). On an individual level, F cell ratios have been reported to display wide scattering from 0.76 to 1.15 (25). Therefore, the input of a mean F cell ratio for different patients may introduce a substantial error in calculating erythrocyte volumes in patients who have an F cell ratio that differs from the mean value (15,25). In our study, the mean pre-HD F cell ratio differed from

the mean population values mentioned. In addition, our patients displayed a wide interindividual variability in F cell ratio (from 0.84 to 0.95 before HD). Nevertheless, our study does support the hypothesis from Mitra *et al.* In their study, they could not explain the apparent rise in erythrocyte volume other than by an increase in F cell ratio during HD with UF (13). Our direct F cell measurements underline this hypothesis. It is interesting that the difference between ΔRBV and ΔTBV that we found was of the same order of magnitude as in the study of Mitra *et al.* (8.2 ± 3.7 and $7.7 \pm 10.6\%$, respectively).

In this study, we used ¹²⁵I-albumin as a second plasma volume marker to quantify and correct for TER_{alb} from ¹²⁵I-albumin during HD. In healthy control subjects, TER_{alb} has been demonstrated to range from 4.3 to 7.4%/h (17,26–28). Hildebrandt *et al.* (17) showed a mean TER_{alb} of 9.6% in HD patients before the start of HD, and Geers *et al.* (26) found a TER_{alb} of $9.8 \pm 2.6\%$ in patients with the nephrotic syndrome. As a mean, we found an intradialytic TER_{alb} of $9.1 \pm 1.3\%$, which is in line with the aforementioned studies.

This study has some important pathophysiologic and clinical implications. It shows directly that the F cell ratio rises during HD. This observation strongly suggests that during HD, translocation of blood from the microcirculation to the macrocirculation takes place, which probably is a cardiovascular compensatory mechanism in response to hypovolemia. The underestimation of the real TBV decline may explain why some patients develop hypotension at a relatively small RBV decline. In addition, the poor predictive value of ΔRBV for the occurrence of dialysis hypotension (reviewed by Dasselaar *et al.* [12]) may well partly be explained by variability in the divergence between ΔRBV and ΔTBV (and thus a variable rise in F cell ratio) during HD. In this study, the extent of the divergence between ΔRBV and ΔTBV indeed differed considerably between individuals. Some patients displayed a relatively small difference between ΔRBV and ΔTBV (*e.g.*, -12.2 versus -14.0% in patient 7), whereas others displayed a substantial difference (*e.g.*, -8.1 versus -22.8% in patient 3). Notably, the divergence between ΔRBV and ΔTBV was of the same order of magnitude in patients with a relative low and high UF volume. In other words, the difference between ΔRBV and ΔTBV was also prominent in the patients with relatively low UF volumes (*e.g.*, -5.7 versus -14.3% with a UF volume of 1200 ml in patient 2).

Factors that may influence the extent of the intradialytic translocation of blood from the microcirculation to the macrocirculation (and thus the extent of the change in F cell ratio) are ambient and dialysate temperature, changes in position, administration of vasoactive drugs and exercise during HD. All of these factors are known to influence ΔRBV (reviewed in reference [12]). It remains to be studied whether these factors also influence ΔTBV and/or the divergence between ΔRBV and ΔTBV.

In this study, we investigated hemodynamically stable patients. Future studies should also address the issue of underestimation of ΔTBV by ΔRBV and the change in F cell ratio in hypotension-prone HD patients. For the moment, however, it is important for the dialysis staff to realize that the relative change in blood volume, as calculated by ΔRBV monitors, may mark-

edly underestimate the real decline in blood volume, especially in the second half of the HD session.

Conclusion

Δ RBV significantly underestimates Δ TBV during HD. The rise in F cell ratio strongly suggests that blood translocates from the microcirculation to the macrocirculation during HD, probably as a cardiovascular compensatory mechanism in response to intravascular volume depletion.

Acknowledgments

This study was supported by grants from Amgen BV and Gambro-Hospital BV.

Disclosures

None.

References

- Steuer RR, Leypoldt JK, Cheung AK, Senekjian HO, Conis JM: Reducing symptoms during hemodialysis by continuously monitoring the hematocrit. *Am J Kidney Dis* 27: 525–532, 1996
- Basile C: Should relative blood volume changes be routinely measured during the dialysis session? *Nephrol Dial Transplant* 16: 10–12, 2001
- Stiller S, Wirtz D, Waterbar F, Gladziwa U, Dakshinamurty KV, Mann H: Less symptomatic hypotension using blood volume controlled ultrafiltration. *ASAIO Trans* 37: M139–M141, 1991
- Johner C, Chamney PW, Schneditz D, Kramer M: Evaluation of an ultrasonic blood volume monitor. *Nephrol Dial Transplant* 13: 2098–2103, 1998
- Santoro A, Mancini E, Paolini F, Zucchelli P: Blood volume monitoring and control. *Nephrol Dial Transplant* 11: 42–47, 1996
- Steuer R, Harris D, Conis J: A new optical technique for monitoring hematocrit and circulating blood volume: Its application in renal dialysis. *Dial Transplant* 22: 260–265, 1993
- Harrison M, Graveney M, Cochrane L: Some sources of error in the calculation of relative change in plasma volume. *Eur J Appl Physiol* 50: 13–21, 1982
- Gibson JG, Seligman AM, Peacock WC, Aub JC, Fine J, Evans RD: The distribution of red cells and plasma in large and minute vessels of the normal dog, determined by radioactive isotopes of iron and iodine. *J Clin Invest* 25: 848–857, 1946
- Goldsmith H, Cokelet G, Gaetgens P: Robin Fahraeus: Evolution of his concepts in cardiovascular physiology. *Am J Physiol* 257: H1005–H1115, 1989
- Chaplin H Jr, Mollison P, Vetter H: The body/venous hematocrit ratio: Its constancy over a wide hematocrit range. *J Clin Invest* 32: 1309–1316, 1953
- Gregersen ML, Rawson RA: Blood volume. *Physiol Rev* 39: 307–342, 1959
- Dasselaar JJ, Huisman RM, de Jong PE, Franssen CFM: Measurement of relative blood volume changes during haemodialysis: Merits and limitations. *Nephrol Dial Transplant* 20: 2043–2049, 2005
- Mitra S, Chamney P, Greenwood R, Farrington K: The relationship between systemic and whole-body hematocrit is not constant during ultrafiltration on hemodialysis. *J Am Soc Nephrol* 15: 463–469, 2004
- Schneditz D, Roob J, Vaclavik M, Holzer H, Kenner T: Noninvasive measurement of blood volume in hemodialysis patients. *J Am Soc Nephrol* 7: 1241–1244, 1996
- Recommended methods for measurement of red-cell and plasma volume: International Committee for Standardization in Haematology. *J Nucl Med* 21: 793–800, 1980
- Standard techniques for the measurement of red-cell and plasma volume. A report by the International Committee for Standardization in Hematology (ICSH): Panel on Diagnostic Applications of Radioisotopes in Haematology. *Br J Haematol* 25: 801–814, 1973
- Hildebrandt P, Jensen HA, Henriksen JH: Studies on kinetics of albumin in uraemic patients on chronic haemodialysis: Evidence of interstitial albumin wash-down. *Clin Physiol* 3: 153–162, 1983
- Ookawara S, Suzuki M, Yahagi T, Saitou M, Tabei K: Effect of postural change on blood volume in long-term hemodialysis patients. *Nephron* 87: 27–34, 2001
- Lundvall J, Lindgren P: F-cell shift and protein loss strongly affect validity of PV reductions indicated by Hb/Hct and plasma proteins. *J Appl Physiol* 84: 822–829, 1998
- Lundvall J, Bjerkhoel P: Pronounced and rapid plasma volume reduction upon quiet standing as revealed by a novel approach to the determination of the intravascular volume change. *Acta Physiol Scand* 154: 131–142, 1995
- Shibagaki Y, Takaichi K: Significant reduction of the large-vessel blood volume by food intake during hemodialysis. *Clin Nephrol* 49: 49–54, 1998
- Paolini F, Mancini E, Bosetto A, Santoro A: Hemoscan: A dialysis machine-integrated blood volume monitor. *Int J Artif Organs* 18: 487–494, 1995
- Johansen LB, Videbaek R, Hammerum M, Norsk P: Underestimation of plasma volume changes in humans by hematocrit/hemoglobin method. *Am J Physiol* 274: R126–R130, 1998
- Fairbanks VF, Klee GG, Wiseman GA, Hoyer JD, Tefferi A, Pettitt RM, Silverstein MN: Measurement of blood volume and red cell mass: Re-examination of 51Cr and 125I methods. *Blood Cells Mol Dis* 22: 169–186, 1996
- Balga I, Solenthaler M, Furlan M: Should whole-body red cell mass be measured or calculated? *Blood Cells Mol Dis* 26: 25–31, 2000
- Geers AB, Koomans HA, Boer P, Dorhout Mees EJ: Plasma and blood volumes in patients with the nephrotic syndrome. *Nephron* 38: 170–173, 1984
- Evans TW: Review article: Albumin as a drug—Biological effects of albumin unrelated to oncotic pressure. *Aliment Pharmacol Ther* 16: 6–11, 2002
- Quinlan GJ, Martin GS, Evans TW: Albumin: Biochemical properties and therapeutic potential. *Hepatology* 41: 1211–1219, 2005