

# Role of Mycophenolate Mofetil in the Treatment of Lupus Nephritis

Gabriel Contreras and Jonathan Sosnov

Division of Nephrology and Hypertension, University of Miami Miller School of Medicine, Miami, Florida

*Clin J Am Soc Nephrol* 2: 879-882, 2007. doi: 10.2215/CJN.02740707

In patients with systemic lupus erythematosus (SLE), lupus nephritis is present in approximately 25% of patients at the time of diagnosis and eventually develops in up to 60% of adults and 80% of children (1). Selected observational studies have reported that patients with proliferative glomerulonephritis (class III, IV, and V with intracapillary proliferation) have an average absolute risk for the development of chronic kidney disease (CKD) and all-cause mortality of approximately 25 and 13%, respectively (2–5). Treatment of patients with SLE and proliferative lupus nephritis includes the use of immunosuppressive agents in combination with corticosteroids aiming to reduce the risk for the development of CKD and death.

Optimal management of proliferative lupus nephritis with immunosuppressive agents remains a challenge because of the need to balance the efficacy and safety of the therapeutic agents. In the last 30 yr of the last century, randomized clinical trials performed primarily at the National Institutes of Health demonstrated that regimens using cyclophosphamide with corticosteroids were superior to corticosteroids alone for the treatment of proliferative lupus nephritis (6–10). The incidence of CKD was significantly lower on average of 15% in patients who received long-term cyclophosphamide compared with 45% in patients who received corticosteroids alone. Two published meta-analyses indicated that regimens of cyclophosphamide were more efficacious than regimens of corticosteroids alone, reducing the risk for the development of CKD (11,12) and all-cause mortality (11). However, the success of cyclophosphamide regimens comes with the burden of adverse events. The incidence of amenorrhea is significantly increased, ranging from 45 to 71% in patients who receive cyclophosphamide for >6 mo. In addition, the incidence of herpes zoster infection is significantly increased, ranging from 25 to 33% with the use of cyclophosphamide. Hemorrhagic cystitis is seen primarily with the long-term use of oral cyclophosphamide with an incidence ranging from 14 to 17% (6–10). The safety concerns of cyclophosphamide regimens has led to the use of azathioprine (a nonselective inhibitor of purine synthesis) with corticosteroids, which reduces the risk for all-cause mortality (12). However,

azathioprine does not have a clear beneficial effect on the risk for important renal events (12,13) unless it is used after an induction regimen of cyclophosphamide (14,15).

In the past decade, the immunosuppressive agent mycophenolate mofetil (MMF) has been used in the treatment of lupus nephritis. The development of MMF, which is an ester prodrug of mycophenolic acid (MPA) with superior bioavailability (16), was based on the observations that patients with adenosine deaminase deficiency, a deficiency of the *de novo* pathway for purine synthesis, have a combined B and T cell immunodeficiency, whereas patients with hypoxanthine-guanine-phosphoribosyl transferase deficiency, a defect of the salvage pathway for purine synthesis, develop neurologic abnormalities and gout yet have essentially normal immune functions (17,18). Lymphocytes use preferentially the *de novo* pathway for synthesis of guanosine monophosphate. Therefore, inhibition of the *de novo* purine synthesis seemed to be an attractive option to modulate immune responses while limiting adverse effects of nonselective immunosuppressive agents. MPA, a fermentation product of *Penicillium brevicompactum* and related fungi, is an inhibitor of inosine monophosphate dehydrogenase, a key enzyme in the *de novo* pathway of purine synthesis. As expected, MPA *in vitro* and *in vivo* inhibits lymphocytes proliferation, modulates apoptosis in activated T lymphocytes, and attenuates the autoantibody production by B cells and the production of oxygen radicals and adhesion molecules, all essential mechanisms that propagate the autoimmune and inflammatory responses in SLE (19–21). The efficacy of MMF was demonstrated in rodent models of lupus nephritis (22,23).

The biologic plausibility for the use of MMF in the treatment of proliferative lupus nephritis has led to the study of this agent in five induction trials, aiming to achieve remission, and one maintenance trial, aiming to prevent relapse conducive to CKD.

In one randomized, controlled trial conducted by Chan *et al.* (24), an induction regimen of MMF with corticosteroids for 12 mo ( $n = 21$ ) was compared to an induction regimen of oral cyclophosphamide for 6 mo followed by azathioprine for 6 mo with corticosteroids ( $n = 21$ ). The study included primarily Asian patients with diffuse proliferative glomerulonephritis. In that study, similar remission rates were reported for both groups (combined partial and complete remission rates of 95 and 90% in the MMF and sequential groups, respectively). Recently, Chan *et al.* (25) published their 5-yr follow-up data of their clinical trial in which additional patients were randomly

---

Published online ahead of print. Publication date available at [www.cjasn.org](http://www.cjasn.org).

**Address correspondence to:** Dr. Gabriel Contreras, Division of Nephrology and Hypertension, University of Miami Miller School of Medicine, 1600 NW 10th Avenue, Suite 7168, Miami, FL 33133. Phone: 305-243-3583; Fax: 305-243-3506; E-mail: [gcontrer@med.miami.edu](mailto:gcontrer@med.miami.edu)

assigned to receive MMF ( $n = 33$ ) and sequential cyclophosphamide followed by azathioprine ( $n = 31$ ). Both short- and long-term efficacies in renal outcomes were again statistically similar with the incidences of leucopenia (0 versus 26%;  $P = 0.002$ ), infections (13 versus 40%;  $P = 0.013$ ), and amenorrhea (4 versus 36%;  $P = 0.004$ ) significantly lower in the MMF group compared to the sequential group. Although the studies of Chan *et al.* were not powered to detect a difference in hard clinical outcomes between the two arms, the encouraging results in other outcomes were mirrored in another Chinese nonrandomized, controlled trial by Hu *et al.* (26), who compared induction MMF with corticosteroids to pulses of intravenous cyclophosphamide with corticosteroids. In another randomized, controlled trial from Malaysia published by Ong *et al.* (27), an induction regimen of MMF with corticosteroids ( $n = 19$ ) was compared to monthly pulses of intravenous cyclophosphamide with corticosteroids ( $n = 25$ ) for 6 mo in patients with proliferative lupus nephritis. In that study, the induction regimen of MMF with corticosteroids was as effective as intravenous cyclophosphamide with corticosteroids (combined partial and complete remission rates of 58 and 52%, respectively) without difference between the two groups in the rate of adverse events. Recently, Ginzler *et al.* (28) reported that 6 mo of MMF with corticosteroids ( $n = 71$ ) was superior to monthly pulses of intravenous cyclophosphamide with corticosteroids ( $n = 69$ ) for 6 mo inducing complete (22.5 versus 5.8%;  $P < 0.05$ ) and combined partial and complete (52.1 versus 30.4;  $P < 0.05$ ) remissions of proliferative lupus nephritis in patients with SLE. The induction with MMF compared to intravenous cyclophosphamide had a more favorable safety profile with particularly fewer pyogenic infections (relative risk 0.36;  $P = 0.030$ ). The study by Ginzler *et al.* (28) included a large proportion of African-American patients (56%), who traditionally have been thought to have a more aggressive disease course with poor response to cyclophosphamide, giving additional credence to the notion that induction therapy with MMF is an alternative to cyclophosphamide. In a recent randomized controlled trial reported by Flores-Suarez *et al.* (29), published only in abstract form, an induction regimen of MMF ( $n = 10$ ) had similar efficacy and safety compared to intravenous cyclophosphamide ( $n = 10$ ).

In this issue of *CJASN*, Walsh *et al.* (30) report the results of a meta-analysis that pooled four randomized, controlled trials that compared MMF to cyclophosphamide as induction agents. The relative risk for failure to induce combined partial and complete remission was 30% lower ( $P = 0.004$ ) for patients who were treated with MMF compared to those who were treated with cyclophosphamide. At the end of the prespecified studies' duration, the relative risk for the composite outcome of ESRD or death was 65% lower for patients who were treated with MMF compared to those who were treated with cyclophosphamide; however, the 95% confidence interval (CI) of this estimate was wide and nonsignificant with a risk reduction to risk increase ranging from  $-90$  to  $+22\%$  ( $P = 0.10$ ). Analysis of this composite outcome using the extended-study follow-up showed a significant risk reduction of 54% ( $P = 0.02$ ) for patients who were treated with MMF compared to those who

were treated with cyclophosphamide. The relative risk for infections was 36% lower for patients who were treated with MMF compared to those who were treated with cyclophosphamide; however, the 95% CI of this estimate was wide and nonsignificant, ranging from  $-61$  to  $+6\%$  ( $P = 0.085$ ). Other adverse events were too few, resulting in inadequate power to compare both induction agents.

Another meta-analysis that evaluated MMF in lupus nephritis, pooling five induction trials, also showed that MMF was superior to cyclophosphamide (31). Combined partial and complete remission was significantly more frequent with MMF (66%) than with cyclophosphamide (54%), with a number needed to treat of eight (95% CI 4.3 to 69) to induce one additional combined partial and complete remission. Serious infections occurred less frequently with MMF (3.9%) than with cyclophosphamide (15%), with a number needed to treat of 8.7 (95% CI 5.5 to 21) to prevent one serious infectious event. Leucopenia also occurred less frequently with MMF (1.6%) than with cyclophosphamide (25%), with a number needed to treat of 4.3 (95% CI 2.9 to 8.3) to prevent one leucopenia event. Amenorrhea occurred less frequently with MMF (1.9%) than with cyclophosphamide (12%), with a number needed to treat of 9.5 (95% CI 6.2 to 20) to prevent one amenorrhea event. Diarrhea occurred more frequently with MMF (16%) than with cyclophosphamide (4%), with a number needed to treat of 8.5 (95% CI 5.3 to 21) to cause one additional diarrhea episode.

The meta-analysis by Walsh *et al.* (30) achieves its goals of pooling together four induction trials that were in most instances underpowered to detect superiority differences in outcomes of patients who were treated with MMF compared to cyclophosphamide. The consistent directionality across all studies of the superiority of MMF is reassuring. However, a meta-analysis is limited to what was studied and published in the combined studies. In most instances, the included studies were of limited quality by Jadad scores. The participants in the studies also varied with their risk for remission as shown by their varying rates of remission. With only four studies, a meta-regression exploring factors for why these studies differ cannot be done. Epidemiologic evidence suggests the importance of patient baseline characteristics (4,32), which may be associated with response to regimens such as MMF or cyclophosphamide. A patient-level meta-analysis to examine directly, for example, factors such as treatment allocation–race interaction and baseline activity of the disease, combining the actual data sets, might have helped to provide more information. The investigators mentioned other limitations worth highlighting. The participants included in those trials had a relatively preserved filtration function, and their results cannot be generalized to patients with moderately to severely reduced filtration function and rapidly progressive glomerulonephritis. Another limitation is that the result of the analysis for the composite outcome of ESRD or death using the extended-study follow-up may be related to factors other than the comparison of MMF to cyclophosphamide as induction agents. Such factors may be the actual maintenance regimen used (crossover or another immunosuppressive agent, such as azathioprine), the adherence to treatment, and the way patients were lost to

follow-up. The analysis of the extended follow-up should be considered only an observational phase, limiting the value of the treatments' comparison.

In summary, in patients with SLE, the development of proliferative lupus nephritis adds significant morbidity and mortality. The treatment options for proliferative lupus nephritis continue to evolve. In the past decade, clinical trials have defined better the role of MMF in the treatment of this disease. As highlighted by the clinical trials and the meta-analyses, MMF can be used as an induction agent for patients with active proliferative lupus nephritis and fairly preserved renal filtration function. The limitations of small, underpowered clinical trials and any meta-analysis may be helped by the ongoing large, multicenter, clinical trials that are comparing MMF as a prolonged induction/maintenance agent to cyclophosphamide or azathioprine, respectively (33,34).

## Disclosures

G.C. has received grant support from Roche and Aspreva to investigate the role of MMF in the treatment of lupus nephritis and serve as a speaker of conferences.

## References

- Cameron JS: Lupus nephritis. *J Am Soc Nephrol* 10: 413-424, 1999
- Appel GB, Cohen DJ, Pirani CL, Meltzer JL, Estes D: Long-term follow-up of patients with lupus nephritis: A study based on the classification of the World Health Organization. *Am J Med* 83: 877-885, 1987
- Contreras G, Pardo V, Cely C, Borja E, Hurtado A, De La Cuesta C, Iqbal K, Lenz O, Asif A, Nahar N, Leclercq B, Leon C, Schulman I, Ramirez-Seijas F, Paredes A, Cepero A, Khan T, Pachon F, Tozman E, Barreto G, Hoffman D, Almeida Suarez M, Busse JC, Esquenazi M, Esquenazi A, Garcia Mayol L, Garcia Estrada H: Factors associated with poor outcomes in patients with lupus nephritis. *Lupus* 14: 890-895, 2005
- Korbet SM, Schwartz MM, Evans J, Lewis EJ: Severe lupus nephritis: Racial differences in presentation and outcome. *J Am Soc Nephrol* 18: 244-254, 2007
- Mak A, Mok CC, Chu WP, To CH, Wong SN, Au TC: Renal damage in systemic lupus erythematosus: A comparative analysis of different age groups. *Lupus* 16: 28-34, 2007
- Austin HA 3rd, Klippel JH, Balow JE, le Riche NG, Steinberg AD, Plotz PH, Decker JL: Therapy of lupus nephritis: Controlled trial of prednisone and cytotoxic drugs. *N Engl J Med* 314: 614-619, 1986
- Boumpas DT, Austin HA 3rd, Vaughn EM, Klippel JH, Steinberg AD, Yarboro CH, Balow JE: Controlled trial of pulse methylprednisolone versus two regimens of pulse cyclophosphamide in severe lupus nephritis. *Lancet* 340: 741-745, 1992
- Gourley MF, Austin HA 3rd, Scott D, Yarboro CH, Vaughan EM, Muir J, Boumpas DT, Klippel JH, Balow JE, Steinberg AD: Methylprednisolone and cyclophosphamide, alone or in combination, in patients with lupus nephritis: A randomized, controlled trial. *Ann Intern Med* 125: 549-557, 1996
- Illei GG, Austin HA, Crane M, Collins L, Gourley MF, Yarboro CH, Vaughan EM, Kuroiwa T, Danning CL, Steinberg AD, Klippel JH, Balow JE, Boumpas DT: Combination therapy with pulse cyclophosphamide plus pulse methylprednisolone improves long-term renal outcome without adding toxicity in patients with lupus nephritis. *Ann Intern Med* 135: 248-257, 2001
- Steinberg AD, Steinberg SC: Long-term preservation of renal function in patients with lupus nephritis receiving treatment that includes cyclophosphamide versus those treated with prednisone only. *Arthritis Rheum* 34: 945-950, 1991
- Bansal VK, Beto JA: Treatment of lupus nephritis: A meta-analysis of clinical trials. *Am J Kidney Dis* 29: 193-199, 1997
- Flanc RS, Roberts MA, Strippoli GF, Chadban SJ, Kerr PG, Atkins RC: Treatment of diffuse proliferative lupus nephritis: A meta-analysis of randomized controlled trials. *Am J Kidney Dis* 43: 197-208, 2004
- Grootscholten C, Ligtenberg G, Hagen EC, van den Wall Bake AW, de Glas-Vos JW, Bijl M, Assmann KJ, Bruijn JA, Weening JJ, van Houwelingen HC, Derksen RH, Berden JH; Dutch Working Party on Systemic Lupus Erythematosus: Azathioprine/methylprednisolone versus cyclophosphamide in proliferative lupus nephritis: A randomized controlled trial. *Kidney Int* 70: 732-742, 2006
- Contreras G, Pardo V, Leclercq B, Lenz O, Tozman E, O'Nan P, Roth D: Sequential therapies for proliferative lupus nephritis. *N Engl J Med* 350: 971-980, 2004
- Houssiau FA, Vasconcelos C, D'Cruz D, Sebastiani GD, Garrido Ed Ede R, Danieli MG, Abramovicz D, Blockmans D, Mathieu A, Direskeneli H, Galeazzi M, Gul A, Levy Y, Petera P, Popovic R, Petrovic R, Sinico RA, Cattaneo R, Font J, Depresseux G, Cosyns JP, Cervera R: Immunosuppressive therapy in lupus nephritis: The Euro-Lupus Nephritis Trial, a randomized trial of low-dose versus high-dose intravenous cyclophosphamide. *Arthritis Rheum* 46: 2121-2131, 2002
- Lee WA, Gu L, Miksztal AR, Chu N, Leung K, Nelson PH: Bioavailability improvement of mycophenolic acid through amino ester derivatization. *Pharm Res* 7: 161-166, 1990
- Allison AC, Hovi T, Watts RW, Webster AD: Immunological observations on patients with Lesch-Nyhan syndrome, and on the role of de-novo purine synthesis in lymphocyte transformation. *Lancet* 2: 1179-1183, 1975
- Giblett ER, Anderson JE, Cohen F, Pollara B, Meuwissen HJ: Adenosine-deaminase deficiency in two patients with severely impaired cellular immunity. *Lancet* 2: 1067-1069, 1972
- Allison AC, Eugui EM: Mycophenolate mofetil and its mechanisms of action. *Immunopharmacology* 47: 85-118, 2000
- Eugui EM, Almquist SJ, Muller CD, Allison AC: Lymphocyte-selective cytostatic and immunosuppressive effects of mycophenolic acid in vitro: Role of deoxyguanosine nucleotide depletion. *Scand J Immunol* 33: 161-173, 1991
- Eugui EM, Mirkovich A, Allison AC: Lymphocyte-selective antiproliferative and immunosuppressive effects of mycophenolic acid in mice. *Scand J Immunol* 33: 175-183, 1991
- Corna D, Morigi M, Facchinetti D, Bertani T, Zoja C, Remuzzi G: Mycophenolate mofetil limits renal damage and

- prolongs life in murine lupus autoimmune disease. *Kidney Int* 51: 1583-1589, 1997
23. Van Bruggen MC, Walgreen B, Rijke TP, Berden JH: Attenuation of murine lupus nephritis by mycophenolate mofetil. *J Am Soc Nephrol* 9: 1407-1415, 1998
  24. Chan TM, Li FK, Tang CS, Wong RW, Fang GX, Ji YL, Lau CS, Wong AK, Tong MK, Chan KW, Lai KN: Efficacy of mycophenolate mofetil in patients with diffuse proliferative lupus nephritis. Hong Kong-Guangzhou Nephrology Study Group. *N Engl J Med* 343: 1156-1162, 2000
  25. Chan TM, Tse KC, Tang CS, Mok MY, Li FK: Long-term study of mycophenolate mofetil as continuous induction and maintenance treatment for diffuse proliferative lupus nephritis. *J Am Soc Nephrol* 16: 1076-1084, 2005
  26. Hu W, Liu Z, Chen H, Tang Z, Wang Q, Shen K, Li L: Mycophenolate mofetil vs cyclophosphamide therapy for patients with diffuse proliferative lupus nephritis. *Chin Med J (Engl)* 115: 705-709, 2002
  27. Ong LM, Hooi LS, Lim TO, Goh BL, Ahmad G, Ghazalli R, Teo SM, Wong HS, Tan SY, Shaariah W, Tan CC, Morad Z: Randomized controlled trial of pulse intravenous cyclophosphamide versus mycophenolate mofetil in the induction therapy of proliferative lupus nephritis. *Nephrology (Carlton)* 10: 504-510, 2005
  28. Ginzler EM, Dooley MA, Aranow C, Kim MY, Buyon J, Merrill JT, Petri M, Gilkeson GS, Wallace DJ, Weisman MH, Appel GB: Mycophenolate mofetil or intravenous cyclophosphamide for lupus nephritis. *N Engl J Med* 353: 2219-2228, 2005
  29. Flores-Suarez LF, Villa AR: Open randomized trial comparing mycophenolate mofetil versus intravenous cyclophosphamide as induction therapy for severe lupus nephritis. *J Am Soc Nephrol* 15: P0257, 2004
  30. Walsh M, James M, Jayne D, Tonelli M, Manns BJ, Hemmelgarn BR: Mycophenolate mofetil for induction therapy of lupus nephritis: A systematic review and meta-analysis. *Clin J Am Soc Nephrol* 2: 968-975, 2007
  31. Moore RA, Derry S: Systematic review and meta-analysis of randomised trials and cohort studies of mycophenolate mofetil in lupus nephritis. *Arthritis Res Ther* 8: R182, 2006
  32. Mok CC, Ying KY, Tang S, Leung CY, Lee KW, Ng WL, Wong RW, Lau CS: Predictors and outcome of renal flares after successful cyclophosphamide treatment for diffuse proliferative lupus glomerulonephritis. *Arthritis Rheum* 50: 2559-2568, 2004
  33. ALMS: Aspreva Lupus Management Study. Available at: <http://www.almstudy.com>. Accessed June 6, 2007
  34. Mycophenolate Mofetil Versus Azathioprine for Maintenance Therapy of Lupus Nephritis. Available at: <http://clinicaltrials.gov/ct/show/NCT00204022;jsessionid=95F7CD63B7CAE2A25DCD8D97CFE4E0F6?order=12>. Accessed June 6, 2007

See the related article, "Mycophenolate Mofetil for Induction Therapy of Lupus Nephritis: A Systematic Review and Meta-Analysis," on pages 968-975.