

BK Virus Nephropathy and Kidney Transplantation

Daniel L. Bohl and Daniel C. Brennan

Department of Internal Medicine, Washington University School of Medicine, St. Louis, Missouri

Nephropathy from BK virus (BKV) infection is an evolving challenge in kidney transplant recipients. It is the consequence of modern potent immunosuppression aimed at reducing acute rejection and improving allograft survival. Untreated BKV infections lead to kidney allograft dysfunction or loss. Decreased immunosuppression is the principle treatment but predisposes to acute and chronic rejection. Screening protocols for early detection and prevention of symptomatic BKV nephropathy have improved outcomes. Although no approved antiviral drug is available, leflunomide, cidofovir, quinolones, and intravenous Ig have been used. Retransplantation after BKV nephropathy has been successful.

Clin J Am Soc Nephrol 2: S36–S46, 2007. doi: 10.2215/CJN.00920207

Polyomavirus infection in kidney transplant recipients is of increasing interest and research. Although the two human polyomaviruses, BK virus (BKV) and JC virus (JCV), were reported in 1971 (1,2), their influence and importance were limited. The emergence of polyomavirus nephropathy has coincided with the use of new potent immunosuppressive medications (3,4). It is usually associated with BKV, affects up to 8% of recipients, and frequently results in allograft loss or permanent dysfunction (5). It presents as an asymptomatic gradual rise in creatinine with a tubulointerstitial nephritis that mimics rejection, producing a treatment dilemma. The decrease in immunosuppression that is needed to treat infection is opposite to the increases that are needed to treat rejection.

Two studies in kidney transplant recipients who were treated with prednisone and azathioprine in the early 1980s have provided the foundation for much of our current understanding of polyomaviruses in transplant recipients. Hogan *et al.* (6) and Gardner *et al.* (7) found that the pretransplantation seroprevalence was 80 to 88% for BKV and 54 to 55% for JCV. The posttransplantation rates of polyomavirus infection were 18 to 44% for BKV and 30 to 35% for JCV. Most polyomavirus infections were asymptomatic and occurred within the first 3 mo after transplantation. BKV infection was associated with a rising creatinine. More than 20 yr ago, Gardner *et al.* (7) warned, "The detection of polyomavirus infection is important as increased immunosuppression needs to be avoided to prevent possible complications."

Epidemiology

Three polyomaviruses—JCV, BKV, and SV40—cause disease in humans. Humans are the natural host for JCV and BKV. On the basis of serology, BKV is acquired during childhood, and seroprevalence stabilizes or wanes with increasing age (8,9). In contrast, JCV seroprevalence increases with age. The route of

the primary infection may be fecal-oral, respiratory, transplant, or from donor tissue (10–13). Presumably, during a viremic phase, the virus infects target tissues, including the uroepithelium, lymphoid tissue, and brain (13,14), establishing a latent or permissively lytic infection. SV40, a simian virus, was introduced into the human population through contaminated polio and adenovirus vaccines (15). It can be acquired through close contact with nonhuman primates and may spread at a low rate from person to person (10,16,17). Although SV40 has been identified in kidney transplant biopsies and associated with native kidney diseases (18–21), its importance in kidney transplantation is poorly defined and is not discussed further.

Virology

BKV and JCV are small, nonenveloped viruses with an icosahedral capsid and a core of circular double-stranded DNA in association with histones (22). The genome is transcribed bidirectionally. It encodes for the early regulatory proteins—small t antigen and large T antigen—and the late structural proteins—VP1, VP2, and VP3. The genome also contains a non-coding control region that contains the origin of replication and transcription factor binding sites. The agnogene and its protein product help regulate the virus replication and disrupt host cell processes (23–25). The capsid consists of 72 pentamers, each with five VP1 proteins and a central VP2 or VP3 protein. VP1 binds the sialic acid residues of its receptor onto permissive cells (26). The gangliosides GD1b and GT1b and $\alpha(2,3)$ -linked sialic acids on N-linked glycoproteins can act as the receptor for BKV (27,28), whereas $\alpha(2,6)$ -linked sialic acids and the serotonin receptor 5HT2A can act as the receptor for JCV (29,30). After attachment, BKV is internalized *via* caveolae-mediated endocytosis, whereas JCV enters through a clathrin-dependent endocytosis (31,32). Once inside the cell, the viruses traffic to the nucleus and establish a latent or lytic infection. Although JCV resides in the uroepithelium (33) and commonly reacts (6,7,34,35), it rarely causes nephropathy (36,37). Therefore, the remaining discussion focuses on BKV nephropathy.

Address correspondence to: Dr. Daniel C. Brennan, Department of Medicine, Transplant Nephrology, Washington University, Barnes-Jewish Hospital, 6107 Queeny Tower, One Barnes-Jewish Hospital Plaza, St. Louis, MO 63110. Phone: 314-362-8351; Fax: 314-361-4197; E-mail brennan@wudosis.wustl.edu

Pathogenesis

Replication of BKV occurs during states of immune suppression. BKV viremia occurs in pregnancy, cancer, HIV infection, diabetes, and transplantation (13). BKV viremia and BKV nephropathy, however, are rare outside of kidney transplantation. BKV viremia occurs in 13% and BKV nephropathy in 8% of kidney transplant recipients (5). In a kidney transplant recipient, BKV reactivation can come from the donor or the recipient. Recipients who had BKV infection and received a kidney from the same donor have been shown to have identical BKV genotypes, supporting donor transmission (12,38). Recipients whose donors had higher BKV antibody titers were more likely to develop BKV infection than those with lower titers, also supporting donor transmission (12,39). Injury is also believed to contribute to reactivation. In a mouse polyomavirus model, mechanical or chemical injury allowed for initiation of acute infection and also reactivation of latent polyomavirus (40). In humans, injury could come from ischemia or stent placement, or rejection could allow for new infection and reactivation of latent infection in either the donor or recipient.

Once the virus has reactivated, an ascending infection *via* cell-to-cell spread occurs (41–43). Without appropriate immunologic control, a progressive lytic infection ensues (44). This results in large nuclear and perinuclear virus-containing inclusions in the tubule cells. Lysis of these infected cells results in viral seepage into the tubule lumen and urine but also to the interstitium and propagation to surrounding cells. Subsequent tubular cell necrosis leads to cast formation and denudation of the basement membrane. Destruction of tubular capillary walls results in vascular spread of the virus. A heterogeneous interstitial infiltration of inflammatory cells as well as tubulitis may be absent, intermixed with the active infection, or noted in areas that lack cytopathic changes. Collateral damage with necrosis and apoptosis of noninfected tubule cells may occur. The resultant effect of continued intra-graft inflammation, tubular injury, and upregulation of profibrotic mediators is allograft dysfunction and loss.

Early retrospective studies identified tacrolimus and mycophenolate mofetil (MMF) as risk factors for BKV nephropathy (4,43,45–47). More recent retrospective studies found BKV nephropathy associated with the combination of tacrolimus levels (>8 ng/ml) and MMF dosages (1.5 to 2 g/d) (48,49). BKV nephropathy, however, has been reported with triple drug regimens that include a calcineurin inhibitor (tacrolimus or cyclosporine), an adjuvant agent (MMF, azathioprine, or sirolimus) and prednisone (45,49–53), calcineurin-free triple drug therapy (54), double therapy with a calcineurin inhibitor and sirolimus (50,55), tacrolimus monotherapy (56), and with or without use of an induction agent (53). A prospective, randomized study showed that BK viremia and viremia were not different among those who received tacrolimus compared with cyclosporine, azathioprine compared with MMF, and rabbit-antithymocyte globulin induction compared with no induction (57). The highest rates of viremia and viremia were among those who received the combination of cyclosporine and azathioprine or tacrolimus and MMF. Taken together, these studies suggest

that it is the net state of immunosuppression and not a specific drug that allows for development of progressive BKV infection.

Immunology

Although polyomavirus reactivation is common, clinically significant disease is unusual. This is because most recipients are able to control the viruses. Persistent viral infections, such as polyomaviruses, cannot be completely cleared and require continuous immune control (58–60). BKV replication typically begins early after transplantation and after treatment of rejection when immunosuppression is greater and immune control is reduced. The contribution of the humoral, cellular, and innate immune compartments to the control is not well known.

Although 60 to 80% of recipients are BKV seropositive before transplantation (5–7,39,57,61), the presence of these BKV-specific antibodies has not been shown to prevent development of BKV infection. However, BKV-specific antibodies can inhibit BKV infectivity (32,62), and a graded protective effect of the titer of recipient BKV-specific antibodies before transplantation has been suggested (63). BKV seronegativity is also a risk factor for BKV viremia (61) and nephropathy (64) in children. In adults, Shah (65) reported that seropositive donors and seronegative recipients (BKV D+/R–) developed a serologically defined BKV infection most frequently (43%). Bohl *et al.* (12) found that seropositive donor and recipients (BKV D+/R+) developed BKV viremia most frequently (50%). In both studies, only 10% of seronegative donors and recipients developed BKV infection. Thus, BKV antibodies may play a role in the immune response, but they also may indicate a risk for reactivation.

Reduction in immunosuppression results in a significant increase in BKV-specific IgG antibody titers (63,66,67), emergence of BKV-specific cellular immunity (66), clearance of viremia, and stabilization of graft function (57). The presence of BKV antibodies seems to have a limited role. Comoli *et al.* (66) found that despite persistently elevated BKV antibody titers, recurrent BKV viremia was associated with a low frequency of IFN- γ -producing cells. Chen *et al.* (68) found that viremia and an elevated creatinine persisted in most recipients who had BKV nephropathy and developed high BKV antibody titers but weak cytotoxic T lymphocyte responses. However, in recipients with a strong cytotoxic T lymphocyte response but low antibody titers, viremia cleared and creatinine returned to the pre-BKV nephropathy baseline.

The cellular immune response may also contribute to allograft dysfunction. Mannon *et al.* (50) found that the RNA transcriptional profiles that were associated with BKV nephropathy indicated a more intense CD8 functional response and more profibrotic response than acute cellular rejection. Hammer *et al.* (69) found that recipients with viral loads >250,000 copies/ml had detectable BKV-specific CD4⁺ T cells in peripheral blood, but only the two recipients with BKV-specific CD8⁺ T cells >0.1% lost their allografts. The specificity of the cellular response may also be detrimental. Recipients with greater donor and recipient HLA mismatching had an increased incidence of BKV nephropathy (70), possibly mediated by more episodes of rejection, intense immunosuppression, and impaired cytotoxicity in an allogeneic environment but less allograft loss (71). This

suggests that lysis of allogeneic BKV-infected target cells is less efficient with HLA-unrestricted T cells than with HLA-restricted T cells. T cells recognize epitopes that are shared by JCV and BKV that may produce a cross-protective effect (72–74).

Diagnosis

The diagnosis and the severity of BKV infection correspond to our understanding of the pathogenesis of BKV nephropathy. Viral replication begins early after transplantation and progresses through detectable stages: Viruria then viremia then nephropathy (5,57,75–77) (Figure 1). Viruria can be detected by PCR for BKV DNA, reverse transcription–PCR for BKV RNA, cytology for BKV inclusion-bearing epithelial cells termed “decoy cells,” or electron microscopy for viral particles (5,7,57,78). These tests are sensitive for detecting active BKV infections but lack specificity for nephropathy because the detected virus could originate anywhere along the urinary tract. Detection of BKV DNA in the plasma or of viremia may be a better indicator of nephropathy. As the infection intensifies, the markers of viral replication increase. Threshold values have been suggested to predict BKV nephropathy, but considerable overlap of these values exists among recipients without BKV nephropathy, active BKV nephropathy, and resolved BKV nephropathy (79) (Table 1). Therefore, a transplant kidney biopsy remains the gold standard for diagnosing BKV nephropathy. Importantly, the interstitial nephritis and tubular cytopathic changes of BKV nephropathy can be focal or isolated to the medulla and missed on one third of biopsies if only a single core is evaluated (80) (Figure 2). Therefore, at least two cores including medulla should be examined. If there are no cytopathic changes on routine histology but there is a high clinical suspicion, then adjunctive tests such as immunohistochemistry directed specifically against BKV or cross-reacting SV40 large T antigen should be performed because the histopathology of BKV infections may be misinterpreted (45). If the initial biopsy does not confirm BKV nephropathy, then preemptive treatment or repeat biopsy can be considered.

Histology

The characteristic findings on light microscopy are intranuclear basophilic and gelatinous-appearing viral inclusions in




Figure 1. Type and prevalence of BK virus (BKV) infections in kidney transplant recipients.

epithelial cells of the urothelium (46,47,80). These are found in the medulla or cortex and are multifocal with random distribution. Three histologic patterns (A, B, and C) have been described (80–82). In early disease (pattern A), the cytopathic changes are present with little to no inflammation or tubular atrophy. Pattern B consists of viral cytopathic changes with varying degrees of inflammation, tubular atrophy, and fibrosis. In late BKV nephropathy (pattern C), cytopathic changes often are less apparent as a result of a background of tubular atrophy, interstitial fibrosis, and chronic inflammatory infiltrate. The degree of damage corresponds to the degree of allograft dysfunction and allograft outcome (80) (Table 2). The distinction of BKV nephropathy from acute tubular necrosis, interstitial nephritis, and acute cellular rejection is difficult and aided by assessment of blood or urine PCR. Absence of definitive features of acute cellular rejection such as endotheliitis and absence of C4d deposits in peritubular capillaries are helpful. Other histopathologic changes include glomerular crescents (10 to 20%) (46,83), ischemic glomerulopathy (62%) (83), transplant glomerulopathy (62%) (83), abundant plasma cell infiltrates (up to 75%) (46,80,84), and tubular microcalcifications (25%) (81). Features of calcineurin inhibitor toxicity such as striped fibrosis (45,46,80,81), thrombotic microangiopathy (45,81), and tubular isometric vacuolization (81) may also be present.

Treatment

The principal treatment for BKV nephropathy is reduction in immunosuppression. Various strategies include reduction or discontinuation of the calcineurin inhibitor and/or adjuvant agent, changing from MMF to azathioprine, sirolimus, or leflunomide or from tacrolimus to cyclosporine (38,49,52,53,55,85–87). Importantly, BKV nephropathy seems to develop less frequently with maintenance protocols that involve steroid withdrawal (75,88). When BKV nephropathy is diagnosed early within the first 6 mo after transplantation and the creatinine is stable, survival is improved compared with when the diagnosis is made later and the creatinine is elevated.

Early or Presumptive BKV Nephropathy

Reduction in immunosuppression to clear the infection is balanced against the risk for precipitating acute or chronic rejection (Figure 3). Brennan *et al.* (57) showed that preemptive withdrawal of the antimetabolite upon detection of viremia prevented BKV nephropathy without significantly increasing the risk for rejection. Viremia cleared in 22 (96%) of 23 recipients with only one episode of acute rejection directly related to immunosuppression reduction. Of the 22 recipients whose viremia resolved, 32% cleared before protocol decreases in immunosuppression, 32% cleared after withdrawal of the antimetabolite, 9% cleared after reduction in the calcineurin inhibitor, and 27% required withdrawal of the antimetabolite followed by reduction in calcineurin inhibitor for persistent viremia. Prospectively screening pediatric recipients, Hymes and Warshaw (89) cleared viremia in 58% of recipients with presumptive nephropathy after a 50% reduction in the dosage of mycophenolate or sirolimus and targeting tacrolimus troughs of 3 to 5 $\mu\text{g}/\text{dl}$. With BKV nephropathy diagnosed on surveillance bi-

Table 1. Noninvasive tests for BKV nephropathy^a

| Diagnostic Test | Threshold Value | PPV (%) | NPV (%) |
|---|---------------------------|-----------------|-----------------|
| Plasma BKV DNA PCR (copies/ml) (5,43,79,89) | Presence to $\geq 10,000$ | 50 to 85 | 100 |
| Decoy cells (cells/cytospin)(5,43,81) | Presence to ≥ 10 | 27 to 90 | 99 to 100 |
| Urine BKV DNA PCR (copies/ml) (79) | $\geq 1 \times 10^7$ | 67 | 100 |
| Urine BKV mRNA RT-PCR (copies/ng total RNA) (78) | $\geq 6.54 \times 10^5$ | 75 ^b | 97 ^b |

^aBKV, BK virus; NPV, negative predictive value; PPV, positive predictive value; RT-PCR, reverse transcription–PCR.

^bBased on BKV nephropathy biopsy prevalence of 28.6%.




Figure 2. BKV nephropathy. Immunohistochemistry shows characteristic nuclear staining isolated to the medulla.

opsy before an elevation in creatinine, Buehrig *et al.* (51) found that creatinine remained stable and the number of BKV-positive tubules on follow-up biopsy significantly decreased after a step-wise reduction in MMF plus reduction in tacrolimus or conversion to cyclosporine. Conversion from tacrolimus to cyclosporine may lower MMF levels if dosages of MMF remain the same (90). It is interesting that cyclosporine *in vitro* but not tacrolimus *in vitro* has been shown to inhibit BKV reactivation (91). Although complete cessation of MMF may be necessary if viremia persists, MMF may limit proinflammatory and profibrotic cytokines (92,93).

Late BKV Nephropathy

The diagnosis of BKV nephropathy in the setting of allograft dysfunction often indicates more severe histologic changes, and renal function may only stabilize or may continue to progress despite treatment (48,49,51,53,79,80,86,94,95). Also, not treating or inadvertently treating with an antilymphocyte antibody often will lead to progression of disease (96). Whether to reduce or discontinue one or more components of the maintenance regimen is not clear (77,86,94). Ramos *et al.* (53) found no difference in graft survival whether immunosuppression was reduced or continued or between reduction and discontinuation of tacrolimus or MMF among 67 recipients with BKV nephropathy. Vasudev *et al.* (86) used an empiric immunosuppression scale based on drug dosage, not drug level, and found that improvement in renal function after diagnosis of BKV nephropathy correlated with reduction in the calcineurin inhibitor rather than total immunosuppression. Vasudev *et al.* (86) also found that renal function declined by 4.8 ml/min per mo

before the diagnosis of BKV nephropathy and slowed to 0.7 ml/min per mo after a 40% reduction in overall immunosuppression. Renal recovery or stabilization was delayed and occurred at a median of 112 d after diagnosis. Thirty-six percent progressed to allograft loss. This was more common when the creatinine was >2.2 mg/dl at diagnosis. Rocha *et al.* (49) reported no allograft loss and clearance of viremia in seven patients after discontinuation of MMF and reduction in calcineurin inhibitor dosage despite a mean creatinine of 3.2 mg/dl at diagnosis compared with persistent viremia and allograft loss in two recipients after only reduction in MMF and modification of calcineurin inhibitor. Dosage reduction of the adjuvant agent and calcineurin inhibitor in pediatric patients failed to clear viremia with 50% graft loss (89). Dosage reduction of the adjuvant agent and calcineurin inhibitor in adults failed to improve tubular BKV burden with a rising mean creatinine (51). Josephson *et al.* (55) reported only a 15% allograft loss in recipients who had BKV nephropathy and were treated with discontinuation of MMF, dosage reduction of tacrolimus, and addition of leflunomide, an immunosuppressive drug that inhibits BKV, *in vitro*. However, several factors have limited enthusiasm for the use of this drug: (1) High dosages of leflunomide (≥ 40 mg/d) are required to afford efficacy, (2) the relationship between the drug dosage and level is unpredictable, (3) drug levels are not available, and (4) the immunosuppressive potency of leflunomide is weak and the effect that is seen from its use may simply reflect lower immunosuppression.

Adjuvant Therapies

On the basis of *in vitro* activity against BKV, cidofovir, quinolones, and intravenous Ig (IVIG) have been reported as treatment options for BKV nephropathy. Although cidofovir, a cytosine analogue and viral DNA polymerase inhibitor, inhibits BKV replication, the mechanism is unclear because BKV lacks a viral polymerase gene (97–99). Rather than a direct effect on BKV replication, cidofovir may restore the function of p53 and pRB, targets of the large T antigen, and permit BKV-infected cells to undergo apoptosis (22,99). When used for treatment of BKV nephropathy, cidofovir has been given at dosages ranging from 0.25 to 1 mg/kg every 1 to 3 wk with generally favorable results (55,89,95,100–108). However, most studies were observational, and cidofovir was used in conjunction with immuno-

Table 2. Histologic patterns of BKV nephropathy

| Histologic Pattern | Biopsy Findings | Outcome (ESRD) ^a | Differential |
|--------------------|--|-----------------------------|---|
| A | Intranuclear viral inclusions Minimal inflammation tubular cell necrosis fibrosis | 13% | Normal Coexisting diagnosis |
| B | Intranuclear viral inclusions Moderate to severe interstitial inflammation Tubular cell necrosis Minimal tubular atrophy and fibrosis | 55% | Interstitial nephritis Acute tubular necrosis Acute rejection |
| C | Intranuclear viral inclusions Moderate to severe tubular atrophy and fibrosis | 100% | Chronic allograft nephropathy |

^aModified from Li *et al.* (74).




Figure 3. Impaired immune suppression balance. Inadequate immune suppression results in rejection, whereas excessive immune suppression results in BKV nephropathy. Both conditions present as allograft dysfunction with tubulointerstitial nephritis and progression to fibrosis.

suppression reduction. In a cohort of 21 recipients with BKV nephropathy, Kuypers *et al.* (102) reported no graft loss in eight recipients who agreed to treatment with cidofovir but a 70% graft loss in the 13 who did not receive cidofovir. Cidofovir should be used with caution, frequent monitoring, and informed consent because of the potential complications (109,110).

Quinolones, DNA gyrase inhibitors, may interfere with the large T antigen helicase activity (111) and have *in vitro* and *in vivo* activity against BKV (112–114). Two months after a 10-d course of gatifloxacin, seven of 10 recipients with active BKV replication had reduction in viremia or urinary decoy cells (115). Thamboo *et al.* (116), however, did not find improvement in viral clearance after a 10-d course of ciprofloxacin.

IVIG has been used for treatment for BKV nephropathy because of its immunomodulatory (117) as well as potential anti-BKV properties (118). Although IVIG contains BKV-specific antibodies, seropositive recipients as well as recipients

with active BKV infections may have high BKV-specific antibody titers, suggesting that antibody-mediated neutralization does not contribute to viral control. Nevertheless, in combination with immunosuppression reduction, IVIG (2 to 3.5 g/kg over 2 to 7 d) treatment was used as initial treatment for BKV nephropathy (119,120) and BKV nephropathy with concurrent acute rejection (95). Although Sener *et al.* (119) reported that only one (13%) of eight recipients who were treated with IVIG returned to dialysis, half had persistent viremia. Wadei *et al.* (95) found that compared with unmatched control subjects who were not treated with IVIG, IVIG treatment did not improve graft survival. Because of the cost (121), potential adverse effects (122), and unproven efficacy, IVIG use for BKV nephropathy should be limited until controlled studies suggest benefit.

Postinfection Monitoring

Failure to clear BKV leads to worse graft function and outcomes. Because histologic clearance of the virus (79) and disappearance of decoy cells (87) precede clearance from the blood, monitoring should be performed with quantitative assays, preferably BKV PCR, until the viral level is undetectable or at least falls below the threshold value that is associated with BKV nephropathy. On the basis of kinetic models (123) and prospective monitoring (57,79,87,123), viremia clears in 7 to 20 wk, but the initial decrease may be delayed by 4 to 10 wk after immunosuppression reduction. If viremia persists, then further reduction of current maintenance therapy, conversion to sirolimus, or addition of leflunomide can be considered. Twelve weeks after the initial immunosuppression reduction, Wali *et al.* (87) described three recipients who had persistent viremia and responded to further reduction in immunosuppression by conversion to sirolimus (target level 10 to 12 ng/ml) and low-dosage prednisone (2.5 mg every other day).

Retransplantation

BKV nephropathy shortens allograft survival. Tubulointerstitial damage from direct and indirect effects of the virus and rejection after immunosuppression reduction lead to early graft

loss or chronic dysfunction (46,48,51,53,80,86,94,95,124). In recipients who have advanced kidney disease or who have returned to dialysis from BKV nephropathy, retransplantation has been successful (125–130). In most cases, transplant nephrectomy and/or studies to confirm no active viral replication have been performed. However, in the setting of active viremia, viral levels become undetectable within 14 d after preemptive retransplantation with simultaneous allograft nephrectomy despite antibody induction (129). BKV viruria (131), viremia (129), nephropathy (128,132), and graft loss (132) can recur. Because of the long duration after transplantation, intervening negative studies for BKV, and genomic differences, these recurrent BKV infections likely reflected new BKV infections. Therefore, BKV nephropathy is not a contraindication for retransplantation but has recurred in two (12%) of 17 reported recipients. Allograft nephrectomy may not be necessary. In the setting of active viral replication, it seems prudent. Evidence of BKV-specific immunity can be evaluated before retransplantation (125) but in the setting of a resolved infection can be inferred. However, for preemptive transplantation in recipients of combined organ transplants, for whom reduction in immunosuppression is limited, evidence of no active viral replication or BKV-specific immunity should be determined.

BKV Nephropathy and Acute Rejection

The treatment of recipients whose biopsy shows rejection with concurrent BKV nephropathy or early after reduction of immunosuppression to treat BKV nephropathy remains problematic. More than half of biopsies may show tubulitis (5,43,47,80), and reduction in immunosuppression can precipitate rejection in 10 to 30% of recipients (51,53,57,86,108,124). The infiltrating mononuclear cells may represent a BKV-specific and/or allospecific response, and treatment is debatable (133,134). Studies that compared BKV nephropathy with acute rejection have identified differences in the proportion and type of infiltrating cell (80,84,85), protein expression (43,80) and proteomic profiles (135), and gene expression profiles (50). However, these differences have not been characterized serially after modification in immunosuppression. Clinically, reports have described improved, stable, and worse graft function after steroid pulses (5,45,51,94). Comparing recipients who initially received increased immunosuppression with those whose immunosuppression was decreased, Celik *et al.* (94) found no significant short- or long-term improvement in tubulitis or creatinine with brief steroid therapy. On biopsies that were performed within the first 8 wk, the histologic viral load had improved significantly with initially decreased compared with increased immunosuppression but was similar on later biopsies. Therefore, in contrast to others (82), we favor initial reduction in immunosuppression without a steroid pulse. However, the presence of atypical features such as strong peritubular capillary C4d staining, vasculitis, glomerulitis, or interstitial hemorrhage would support rejection (84,136) and require an individualized approach. The delayed improvement in creatinine after reduction in immunosuppression likely reflects the slow resolution of the cellular infiltrate. On follow-up biopsy, we consider a cellular infiltrate with or without tubulitis in the

setting of persistent BKV viremia consistent with resolving BKV nephropathy and continued monitoring. McGilvray *et al.* (137) found severe tubulitis, mild intimal arteritis, and no viral inclusions on a biopsy 1 mo after reduction in immunosuppression. Despite no adjustment to immunosuppression, the patient's creatinine gradually improved. Once BKV nephropathy and viremia have cleared, the benefit of uptitrating immunosuppression to prevent chronic rejection or late acute rejection remains unknown.

Screening

It has been recommended that screening for BKV should be performed every 3 mo for the first 2 yr after transplantation, then annually through the fifth year, when allograft dysfunction occurs and when a transplant kidney biopsy is performed (82). Screening should be based on a urinary assay for decoy cells, BKV DNA, or BKV RNA. A positive screening test should be confirmed within 4 wk along with a quantitative assay. Recipients with persistent high viral levels for >3 wk should undergo biopsy and intervention. Monitoring should continue every 2 to 4 wk until the viral level falls below threshold values and preferably to undetectable levels.

These recommendations are guidelines and should vary on the basis of assay availability and cost, recipient risk, and local incidence of BKV nephropathy. With a low incidence of BKV nephropathy, high false-positive testing rates, or high occurrence of acute rejection or chronic allograft dysfunction after reduction in immunosuppression, screening could produce greater cost and harm than not screening (138). Screening protocols should be used in centers with higher incidences of BKV nephropathy, using triple-drug therapy including tacrolimus and MMF, and with clinical trials to evaluate new therapeutic agents.

At our institution, we screen plasma monthly for the first 6 mo and at months 9 and 12 after transplantation, at the time of a transplant kidney biopsy, and after augmentation in immunosuppression (Figure 4). BKV viremia with stable allograft function triggers empiric immunosuppression reduction and continued monitoring, with the realization that viral levels should decline and clear within 1 to 6 mo (57). An allograft biopsy is performed for allograft dysfunction or persistent high-level viremia. This strategy has resulted in only one case




Figure 4. Screening protocol based on plasma BKV DNA PCR.

of BKV nephropathy in >700 new transplants in the past 5 yr, with an overall acute rejection rate of <10% at 1 yr.

Conclusion

BKV nephropathy remains a significant posttransplantation complication. Modern potent immunosuppressive medications have reduced acute rejection rates, improved early kidney allograft survival, and expanded the potential transplant population. For an individual recipient, however, the increased immunosuppression may be excessive and predispose to BKV nephropathy. No clinical risk factors clearly identify the recipients who will develop BKV nephropathy. Screening all recipients for replication of BKV and targeted reduction of immunosuppression can resolve the infection with stable renal function. However, this method of early detection and treatment is costly, compliance dependent, and potentially detrimental. Once a tubulointerstitial nephritis has developed, progression or resolution to tubular atrophy and interstitial fibrosis results in permanent allograft dysfunction. Treatment of BKV nephropathy is problematic. No antiviral medication is approved, and none has been appropriately studied in a randomized manner. Decreasing immunosuppression to allow a BKV-specific immune response to control the infection is the principle treatment. One potential consequence is the development of an allospecific immune response and rejection. Because monitoring for BKV-specific immunity is not widely available, monitoring for declining viral loads is used as a surrogate. With diffuse parenchymal involvement, resolution of viremia and improvement in creatinine may not occur for months. Although BKV infection is an area of active investigation in kidney transplantation, much of the data are derived from retrospective case-control studies. Prospective, randomized studies to address immunosuppression protocols, immune monitoring for BKV-specific and allospecific responses, and treatment protocols are needed. Until reliable measures of immunosuppression or novel agents that specifically target BKV are available, BKV infections in kidney transplantation will remain a challenge.

Acknowledgments

D.C.B. is supported by a grant from the National Institute of Diabetes and Digestive and Kidney Diseases (K24-DK-002886).

Disclosures

None.

References

- Gardner SD, Field AM, Coleman DV, Hulme B: New human papovavirus (B.K.) isolated from urine after renal transplantation. *Lancet* 1: 1253–1257, 1971
- Padgett BL, Walker DL, ZuRhein GM, Eckroade RJ, Dessel BH: Cultivation of papova-like virus from human brain with progressive multifocal leucoencephalopathy. *Lancet* 1: 1257–1260, 1971
- Purighalla R, Shapiro R, McCauley J, Randhawa P: BK virus infection in a kidney allograft diagnosed by needle biopsy. *Am J Kidney Dis* 26: 671–673, 1995
- Binet I, Nickeleit V, Hirsch HH, Prince O, Dalquen P, Gudat F, Mihatsch MJ, Thiel G: Polyomavirus disease under new immunosuppressive drugs: A cause of renal graft dysfunction and graft loss. *Transplantation* 67: 918–922, 1999
- Hirsch HH, Knowles W, Dickenmann M, Passweg J, Klimkait T, Mihatsch MJ, Steiger J: Prospective study of polyomavirus type BK replication and nephropathy in renal-transplant recipients. *N Engl J Med* 347: 488–496, 2002
- Hogan TF, Borden EC, McBain JA, Padgett BL, Walker DL: Human polyomavirus infections with JC virus and BK virus in renal transplant patients. *Ann Intern Med* 92: 373–378, 1980
- Gardner SD, MacKenzie EF, Smith C, Porter AA: Prospective study of the human polyomaviruses BK and JC and cytomegalovirus in renal transplant recipients. *J Clin Pathol* 37: 578–586, 1984
- Knowles WA, Pipkin P, Andrews N, Vyse A, Minor P, Brown DW, Miller E: Population-based study of antibody to the human polyomaviruses BKV and JCV and the simian polyomavirus SV40. *J Med Virol* 71: 115–123, 2003
- Stolt A, Sasnauskas K, Koskela P, Lehtinen M, Dillner J: Seroepidemiology of the human polyomaviruses. *J Gen Virol* 84: 1499–1504, 2003
- Vanchiere JA, Nicome RK, Greer JM, Demmler GJ, Butel JS: Frequent detection of polyomaviruses in stool samples from hospitalized children. *J Infect Dis* 192: 658–664, 2005
- Bofill-Mas S, Formiga-Cruz M, Clemente-Casares P, Calafell F, Girones R: Potential transmission of human polyomaviruses through the gastrointestinal tract after exposure to virions or viral DNA. *J Virol* 75: 10290–10299, 2001
- Bohl DL, Storch GA, Ryschkewitsch C, Gaudreault-Keener M, Schnitzler MA, Major EO, Brennan DC: Donor origin of BK virus in renal transplantation and role of HLA C7 in susceptibility to sustained BK viremia. *Am J Transplant* 5: 2213–2221, 2005
- Repløeg MD, Storch GA, Clifford DB: BK virus: A clinical review. *Clin Infect Dis* 33: 191–202, 2001
- Eash S, Tavares R, Stopa EG, Robbins SH, Brossay L, Atwood WJ: Differential distribution of the JC virus receptor-type sialic acid in normal human tissues. *Am J Pathol* 164: 419–428, 2004
- Shah KV: Simian virus 40 and human disease. *J Infect Dis* 190: 2061–2064, 2004
- Paracchini V, Garte S, Pedotti P, Poli F, Frison S, Taioli E: Molecular identification of simian virus 40 infection in healthy Italian subjects by birth cohort. *Mol Med* 11: 48–51, 2005
- Engels EA, Switzer WM, Heneine W, Viscidi RP: Serologic evidence for exposure to simian virus 40 in North American zoo workers. *J Infect Dis* 190: 2065–2069, 2004
- Li RM, Branton MH, Tanawattanacharoen S, Falk RA, Jenette JC, Kopp JB: Molecular identification of SV40 infection in human subjects and possible association with kidney disease. *J Am Soc Nephrol* 13: 2320–2330, 2002
- Milstone A, Vilchez RA, Geiger X, Fogo AB, Butel JS, Dummer S: Polyomavirus simian virus 40 infection associated with nephropathy in a lung-transplant recipient. *Transplantation* 77: 1019–1024, 2004
- Li RM, Mannon RB, Kleiner D, Tsokos M, Bynum M, Kirk AD, Kopp JB: BK virus and SV40 co-infection in polyomavirus nephropathy. *Transplantation* 74: 1497–1504, 2002

21. Butel JS, Arrington AS, Wong C, Lednicky JA, Finegold MJ: Molecular evidence of simian virus 40 infections in children. *J Infect Dis* 180: 884–887, 1999
22. Imperiale MJ: The human polyomaviruses: An overview. In: *Human Polyomaviruses: Molecular and Clinical Perspectives*, edited by Khalili K, Stoner GL, New York, Wiley-Liss, 2001, pp 53–71
23. Akan I, Sariyer IK, Biffi R, Palermo V, Woolridge S, White MK, Amini S, Khalili K, Safak M: Human polyomavirus JCV late leader peptide region contains important regulatory elements. *Virology* 349: 66–78, 2006
24. Khalili K, White MK, Sawa H, Nagashima K, Safak M: The agnoprotein of polyomaviruses: A multifunctional auxiliary protein. *J Cell Physiol* 204: 1–7, 2005
25. Suzuki T, Okada Y, Semba S, Orba Y, Yamanouchi S, Endo S, Tanaka S, Fujita T, Kuroda S, Nagashima K, Sawa H: Identification of FEZ1 as a protein that interacts with JC virus agnoprotein and microtubules: Role of agnoprotein-induced dissociation of FEZ1 from microtubules in viral propagation. *J Biol Chem* 280: 24948–24956, 2005
26. Gee GV, Tsomaia N, Mierke DF, Atwood WJ: Modeling a sialic acid binding pocket in the external loops of JC virus VP1. *J Biol Chem* 279: 49172–49176, 2004
27. Low JA, Magnuson B, Tsai B, Imperiale MJ: Identification of gangliosides GD1b and GT1b as receptors for BK virus. *J Virol* 80: 1361–1366, 2006
28. Dugan AS, Eash S, Atwood WJ: An N-linked glycoprotein with alpha(2,3)-linked sialic acid is a receptor for BK virus. *J Virol* 79: 14442–14445, 2005
29. Elphick GF, Querbes W, Jordan JA, Gee GV, Eash S, Manley K, Dugan A, Stanifer M, Bhatnagar A, Kroeze WK, Roth BL, Atwood WJ: The human polyomavirus JCV, uses serotonin receptors to infect cells. *Science* 306: 1380–1383, 2004
30. Komagome R, Sawa H, Suzuki T, Suzuki Y, Tanaka S, Atwood WJ, Nagashima K: Oligosaccharides as receptors for JC virus. *J Virol* 76: 12992–13000, 2002
31. Pho MT, Ashok A, Atwood WJ: JC virus enters human glial cells by clathrin-dependent receptor-mediated endocytosis. *J Virol* 74: 2288–2292, 2000
32. Eash S, Querbes W, Atwood WJ: Infection of vero cells by BK virus is dependent on caveolae. *J Virol* 78: 11583–11590, 2004
33. Boldorini R, Veggiani C, Barco D, Monga G: Kidney and urinary tract polyomavirus infection and distribution: Molecular biology investigation of 10 consecutive autopsies. *Arch Pathol Lab Med* 129: 69–73, 2005
34. Muller A, Beck B, Theilemann K, Stapenhorst L, Licht C, Michalk D, Franzen C, Hoppe B: Detection of polyomavirus BK and JC in children with kidney diseases and renal transplant recipients. *Pediatr Infect Dis J* 24: 778–781, 2005
35. Randhawa P, Uhrmacher J, Pasculle W, Vats A, Shapiro R, Eghtsead B, Weck K: A comparative study of BK and JC virus infections in organ transplant recipients. *J Med Virol* 77: 238–243, 2005
36. Wen MC, Wang CL, Wang M, Cheng CH, Wu MJ, Chen CH, Shu KH, Chang D: Association of JC virus with tubulointerstitial nephritis in a renal allograft recipient. *J Med Virol* 72: 675–678, 2004
37. Kazory A, Ducloux D, Chalopin JM, Angonin R, Fontaniere B, Moret H: The first case of JC virus allograft nephropathy. *Transplantation* 76: 1653–1655, 2003
38. Vera-Sempere FJ, Rubio L, Felipe-Ponce V, Garcia A, Sanahuja MJ, Zamora I, Ramos D, Beneyto I, Sanchez-Plumed J: Renal donor implication in the origin of BK infection: Analysis of genomic viral subtypes. *Transplant Proc* 38: 2378–2381, 2006
39. Andrews CA, Shah KV, Daniel RW, Hirsch MS, Rubin RH: A serological investigation of BK virus and JC virus infections in recipients of renal allografts. *J Infect Dis* 158: 176–181, 1988
40. Atencio IA, Shadan FF, Zhou XJ, Vaziri ND, Villarreal LP: Adult mouse kidneys become permissive to acute polyomavirus infection and reactivate persistent infections in response to cellular damage and regeneration. *J Virol* 67: 1424–1432, 1993
41. Meehan SM, Kraus MD, Kadambi PV, Chang A: Nephron segment localization of polyoma virus large T antigen in renal allografts. *Hum Pathol* 37: 1400–1406, 2006
42. Drachenberg CB, Papadimitriou JC, Wali R, Cubitt CL, Ramos E: BK polyoma virus allograft nephropathy: Ultrastructural features from viral cell entry to lysis. *Am J Transplant* 3: 1383–1392, 2003
43. Nicleleit V, Hirsch HH, Zeiler M, Gudat F, Prince O, Thiel G, Mihatsch MJ: BK-virus nephropathy in renal transplants-tubular necrosis, MHC-class II expression and rejection in a puzzling game. *Nephrol Dial Transplant* 15: 324–332, 2000
44. Low J, Humes HD, Szczypka M, Imperiale M: BKV and SV40 infection of human kidney tubular epithelial cells in vitro. *Virology* 323: 182–188, 2004
45. Howell DN, Smith SR, Butterly DW, Klassen PS, Krigman HR, Burchette JL Jr, Miller SE: Diagnosis and management of BK polyomavirus interstitial nephritis in renal transplant recipients. *Transplantation* 68: 1279–1288, 1999
46. Nicleleit V, Hirsch HH, Binet IF, Gudat F, Prince O, Dalquen P, Thiel G, Mihatsch MJ: Polyomavirus infection of renal allograft recipients: From latent infection to manifest disease. *J Am Soc Nephrol* 10: 1080–1089, 1999
47. Randhawa PS, Finkelstein S, Scantlebury V, Shapiro R, Vivas C, Jordan M, Picken MM, Demetris AJ: Human polyoma virus-associated interstitial nephritis in the allograft kidney. *Transplantation* 67: 103–109, 1999
48. Mengel M, Marwedel M, Radermacher J, Eden G, Schwarz A, Haller H, Kreipe H: Incidence of polyomavirus-nephropathy in renal allografts: Influence of modern immunosuppressive drugs. *Nephrol Dial Transplant* 18: 1190–1196, 2003
49. Rocha PN, Plumb TJ, Miller SE, Howell DN, Smith SR: Risk factors for BK polyomavirus nephritis in renal allograft recipients. *Clin Transplant* 18: 456–462, 2004
50. Mannon RB, Hoffmann SC, Kampen RL, Cheng OC, Kleiner DE, Ryschkewitsch C, Curfman B, Major E, Hale DA, Kirk AD: Molecular evaluation of BK polyomavirus nephropathy. *Am J Transplant* 5: 2883–2893, 2005
51. Buehrig CK, Lager DJ, Stegall MD, Kreps MA, Kremers WK, Gloor JM, Schwab TR, Velosa JA, Fidler ME, Larson TS, Griffin MD: Influence of surveillance renal allograft biopsy on diagnosis and prognosis of polyomavirus-associated nephropathy. *Kidney Int* 64: 665–673, 2003
52. Hirsch HH, Mohaupt M, Klimkait T: Prospective monitoring of BK virus load after discontinuing sirolimus treatment in a renal transplant patient with BK virus nephropathy. *J Infect Dis* 184: 1494–1495, 2001

53. Ramos E, Drachenberg CB, Papadimitriou JC, Hamze O, Fink JC, Klassen DK, Drachenberg RC, Wiland A, Wali R, Cangro CB, Schweitzer E, Bartlett ST, Weir MR: Clinical course of polyoma virus nephropathy in 67 renal transplant patients. *J Am Soc Nephrol* 13: 2145–2151, 2002
54. Lipshutz GS, Flechner SM, Govani MV, Vincenti F: BK nephropathy in kidney transplant recipients treated with a calcineurin inhibitor-free immunosuppression regimen. *Am J Transplant* 4: 2132–2134, 2004
55. Josephson MA, Gillen D, Javaid B, Kadambi P, Meehan S, Foster P, Harland R, Thistlethwaite RJ, Garfinkel M, Atwood W, Jordan J, Sadhu M, Millis MJ, Williams J: Treatment of renal allograft polyoma BK virus infection with leflunomide. *Transplantation* 81: 704–710, 2006
56. Randhawa PS, Gupta G, Vats A, Shapiro R, Viscidi RP: Immunoglobulin G, A, and M responses to BK virus in renal transplantation. *Clin Vaccine Immunol* 13: 1057–1063, 2006
57. Brennan DC, Agha I, Bohl DL, Schnitzler MA, Hardinger KL, Lockwood M, Torrence S, Schuessler R, Roby T, Gaudreault-Keener M, Storch GA: Incidence of BK with tacrolimus versus cyclosporine and impact of preemptive immunosuppression reduction. *Am J Transplant* 5: 582–594, 2005
58. Dorries K: Molecular biology and pathogenesis of human polyomavirus infections. *Dev Biol Stand* 94: 71–79, 1998
59. Kemball CC, Lee ED, Vezys V, Pearson TC, Larsen CP, Lukacher AE: Late priming and variability of epitope-specific CD8⁺ T cell responses during a persistent virus infection. *J Immunol* 174: 7950–7960, 2005
60. Randhawa PS, Popescu I, Macedo C, Zeevi A, Shapiro R, Vats AN, Metes D: Detection of CD8⁺ T cells sensitized to BK virus large T antigen in healthy volunteers and kidney transplant recipients. *Hum Immunol* 67: 298–302, 2006
61. Ginevri F, De Santis R, Comoli P, Pastorino N, Rossi C, Botti G, Fontana I, Nocera A, Cardillo M, Ciardi MR, Locatelli F, Maccario R, Perfumo F, Azzi A: Polyomavirus BK infection in pediatric kidney-allograft recipients: A single-center analysis of incidence, risk factors, and novel therapeutic approaches. *Transplantation* 75: 1266–1270, 2003
62. Flaegstad T, Traavik T, Christie KE, Joergensen J: Neutralization test for BK virus: Plaque reduction detected by immunoperoxidase staining. *J Med Virol* 19: 287–296, 1986
63. Bohl DL, Ryschkewitsch C, Major EO, Storch GA, Brennan DC: BK virus antibody titers markedly increase with viremia [Abstract]. *Am J Transplant* 5: 273, 2005
64. Smith JM, McDonald RA, Finn LS, Healey PJ, Davis CL, Limaye AP: Polyomavirus nephropathy in pediatric kidney transplant recipients. *Am J Transplant* 4: 2109–2117, 2004
65. Shah KV: Human polyomavirus BKV and renal disease. *Nephrol Dial Transplant* 15: 754–755, 2000
66. Comoli P, Azzi A, Maccario R, Basso S, Botti G, Basile G, Fontana I, Labirio M, Cometa A, Poli F, Perfumo F, Locatelli F, Ginevri F: Polyomavirus BK-specific immunity after kidney transplantation. *Transplantation* 78: 1229–1232, 2004
67. Hariharan S, Cohen EP, Vasudev B, Orentas R, Viscidi RP, Kakela J, DuChateau B: BK virus-specific antibodies and BKV DNA in renal transplant recipients with BKV nephritis. *Am J Transplant* 5: 2719–2724, 2005
68. Chen Y, Trofe J, Gordon J, Du Pasquier RA, Roy-Chaudhury P, Kuroda MJ, Woodle ES, Khalili K, Koralknik IJ: Interplay of cellular and humoral immune responses against BK virus in kidney transplant recipients with polyomavirus nephropathy. *J Virol* 80: 3495–3505, 2006
69. Hammer MH, Brestrich G, Andree H, Engelmann E, Rosenberger C, Tillmann H, Zwinger S, Babel N, Nickel P, Volk HD, Reinke P: HLA type-independent method to monitor polyoma BK virus-specific CD4 and CD8 T-cell immunity. *Am J Transplant* 6: 625–631, 2006
70. Awadalla Y, Randhawa P, Ruppert K, Zeevi A, Duquesnoy RJ: HLA mismatching increases the risk of BK virus nephropathy in renal transplant recipients. *Am J Transplant* 4: 1691–1696, 2004
71. Drachenberg CB, Papadimitriou JC, Mann D, Hirsch HH, Wali R, Ramos E: Negative impact of human leukocyte antigen matching in the outcome of polyomavirus nephropathy. *Transplantation* 80: 276–278, 2005
72. Sharma MC, Zhou W, Martinez J, Krymskaya L, Srivastava T, Haq W, Diamond DJ, Lacey SF: Cross-reactive CTL recognizing two HLA-A*02-restricted epitopes within the BK virus and JC virus VP1 polypeptides are frequent in immunocompetent individuals. *Virology* 350: 128–136, 2006
73. Krymskaya L, Sharma MC, Martinez J, Haq W, Huang EC, Limaye AP, Diamond DJ, Lacey SF: Cross-reactivity of T lymphocytes recognizing a human cytotoxic T-lymphocyte epitope within BK and JC virus VP1 polypeptides. *J Virol* 79: 11170–11178, 2005
74. Li J, Melenhorst J, Hensel N, Rezvani K, Sconocchia G, Kilical Y, Hou J, Curfman B, Major E, Barrett AJ: T-cell responses to peptide fragments of the BK virus T antigen: Implications for cross-reactivity of immune response to JC virus. *J Gen Virol* 87: 2951–2960, 2006
75. Bressollette-Bodin C, Coste-Burel M, Hourmant M, Sebillé V, Andre-Garnier E, Imbert-Marcille BM: A prospective longitudinal study of BK virus infection in 104 renal transplant recipients. *Am J Transplant* 5: 1926–1933, 2005
76. Limaye AP, Jerome KR, Kuhr CS, Ferrenberg J, Huang ML, Davis CL, Corey L, Marsh CL: Quantitation of BK virus load in serum for the diagnosis of BK virus-associated nephropathy in renal transplant recipients. *J Infect Dis* 183: 1669–1672, 2001
77. Nicleleit V, Klimkait T, Binet IF, Dalquen P, Del Zenero V, Thiel G, Mihatsch MJ, Hirsch HH: Testing for polyomavirus type BK DNA in plasma to identify renal-allograft recipients with viral nephropathy. *N Engl J Med* 342: 1309–1315, 2000
78. Ding R, Medeiros M, Dadhania D, Muthukumar T, Kracker D, Kong JM, Epstein SR, Sharma VK, Seshan SV, Li B, Suthanthiran M: Noninvasive diagnosis of BK virus nephritis by measurement of messenger RNA for BK virus VP1 in urine. *Transplantation* 74: 987–994, 2002
79. Randhawa P, Ho A, Shapiro R, Vats A, Swalsky P, Finkelstein S, Uhrmacher J, Weck K: Correlates of quantitative measurement of BK polyomavirus (BKV) DNA with clinical course of BKV infection in renal transplant patients. *J Clin Microbiol* 42: 1176–1180, 2004
80. Drachenberg CB, Papadimitriou JC, Hirsch HH, Wali R, Crowder C, Nogueira J, Cangro CB, Mendley S, Mian A, Ramos E: Histological patterns of polyomavirus nephropathy: Correlation with graft outcome and viral load. *Am J Transplant* 4: 2082–2092, 2004

81. Drachenberg RC, Drachenberg CB, Papadimitriou JC, Ramos E, Fink JC, Wali R, Weir MR, Cangro CB, Klassen DK, Khaled A, Cunningham R, Bartlett ST: Morphological spectrum of polyoma virus disease in renal allografts—Diagnostic accuracy of urine cytology. *Am J Transplant* 1: 373–381, 2001
82. Hirsch HH, Brennan DC, Drachenberg CB, Ginevri F, Gordon J, Limaye AP, Mihatsch MJ, Nickleit V, Ramos E, Randhawa P, Shapiro R, Steiger J, Suthanthiran M, Trofe J: Polyomavirus-associated nephropathy in renal transplantation: Interdisciplinary analyses and recommendations. *Transplantation* 79: 1277–1286, 2005
83. Celik B, Randhawa PS: Glomerular changes in BK virus nephropathy. *Hum Pathol* 35: 367–370, 2004
84. Meehan SM, Kadambi PV, Manaligod JR, Williams JW, Josephson MA, Javaid B: Polyoma virus infection of renal allografts: Relationships of the distribution of viral infection, tubulointerstitial inflammation, and fibrosis suggesting viral interstitial nephritis in untreated disease. *Hum Pathol* 36: 1256–1264, 2005
85. Ahuja M, Cohen EP, Dayer AM, Kampalath B, Chang CC, Bresnahan BA, Hariharan S: Polyoma virus infection after renal transplantation. Use of immunostaining as a guide to diagnosis. *Transplantation* 71: 896–899, 2001
86. Vasudev B, Hariharan S, Hussain SA, Zhu YR, Bresnahan BA, Cohen EP: BK virus nephritis: Risk factors, timing, and outcome in renal transplant recipients. *Kidney Int* 68: 1834–1839, 2005
87. Wali RK, Drachenberg C, Hirsch HH, Papadimitriou J, Nahar A, Mohanlal V, Brisco MA, Bartlett ST, Weir MR, Ramos E: BK virus-associated nephropathy in renal allograft recipients: Rescue therapy by sirolimus-based immunosuppression. *Transplantation* 78: 1069–1073, 2004
88. Matas AJ, Kandaswamy R, Humar A, Payne WD, Dunn DL, Najarian JS, Gruessner RW, Gillingham KJ, McHugh LE, Sutherland DE: Long-term immunosuppression, without maintenance prednisone, after kidney transplantation. *Ann Surg* 240: 510–516, 2004
89. Hymes LC, Warshaw BL: Polyomavirus (BK) in pediatric renal transplants: Evaluation of viremic patients with and without BK associated nephritis. *Pediatr Transplant* 10: 920–922, 2006
90. Filler G, Zimmering M, Mai I: Pharmacokinetics of mycophenolate mofetil are influenced by concomitant immunosuppression. *Pediatr Nephrol* 14: 100–104, 2000
91. Acott PD, O'Regan PA, Lee SH, Crocker JF: Antiviral properties of immunosuppressant drugs in primary and latent BK virus infection in kidney cells [Abstract]. *Transplantation* 82: 683, 2006
92. Morath C, Schwenger V, Beimler J, Mehrabi A, Schmidt J, Zeier M, Muranyi W: Antifibrotic actions of mycophenolic acid. *Clin Transplant* 20[Suppl 17]: 25–29, 2006
93. Allison AC, Eugui EM: Mechanisms of action of mycophenolate mofetil in preventing acute and chronic allograft rejection. *Transplantation* 80[Suppl]: S181–S190, 2005
94. Celik B, Shapiro R, Vats A, Randhawa PS: Polyomavirus allograft nephropathy: Sequential assessment of histologic viral load, tubulitis, and graft function following changes in immunosuppression. *Am J Transplant* 3: 1378–1382, 2003
95. Wadei HM, Rule AD, Lewin M, Mahale AS, Khamash HA, Schwab TR, Gloor JM, Textor SC, Fidler ME, Lager DJ, Larson TS, Stegall MD, Cosio FG, Griffin MD: Kidney transplant function and histological clearance of virus following diagnosis of polyomavirus-associated nephropathy (PVAN). *Am J Transplant* 6: 1025–1032, 2006
96. Hussain S, Bresnahan BA, Cohen EP, Hariharan S: Rapid kidney allograft failure in patients with polyoma virus nephritis with prior treatment with antilymphocyte agents. *Clin Transplant* 16: 43–47, 2002
97. Farasati NA, Shapiro R, Vats A, Randhawa P: Effect of leflunomide and cidofovir on replication of BK virus in an in vitro culture system. *Transplantation* 79: 116–118, 2005
98. Andrei G, Snoeck R, Vandeputte M, De Clercq E: Activities of various compounds against murine and primate polyomaviruses. *Antimicrob Agents Chemother* 41: 587–593, 1997
99. De Clercq E: Clinical potential of the acyclic nucleoside phosphonates cidofovir, adefovir, and tenofovir in treatment of DNA virus and retrovirus infections. *Clin Microbiol Rev* 16: 569–596, 2003
100. Tong CY, Hilton R, MacMahon EM, Brown L, Pantelidis P, Chrystie IL, Kidd IM, Tunekar MF, Pattison JM: Monitoring the progress of BK virus associated nephropathy in renal transplant recipients. *Nephrol Dial Transplant* 19: 2598–2605, 2004
101. Araya CE, Lew JF, Fennell RS 3rd, Neiberger RE, Dharnidharka VR: Intermediate-dose cidofovir without probenecid in the treatment of BK virus allograft nephropathy. *Pediatr Transplant* 10: 32–37, 2006
102. Kuypers DR, Vandooren AK, Lerut E, Evenepoel P, Claes K, Snoeck R, Naesens L, Vanrenterghem Y: Adjuvant low-dose cidofovir therapy for BK polyomavirus interstitial nephritis in renal transplant recipients. *Am J Transplant* 5: 1997–2004, 2005
103. Lim WH, Mathew TH, Cooper JE, Bowden S, Russ GR: Use of cidofovir in polyomavirus BK viral nephropathy in two renal allograft recipients. *Nephrology (Carlton)* 8: 318–323, 2003
104. Keller LS, Peh CA, Nolan J, Bannister KM, Clarkson AR, Faull RJ: BK transplant nephropathy successfully treated with cidofovir. *Nephrol Dial Transplant* 18: 1013–1014, 2003
105. Kadambi PV, Josephson MA, Williams J, Corey L, Jerome KR, Meehan SM, Limaye AP: Treatment of refractory BK virus-associated nephropathy with cidofovir. *Am J Transplant* 3: 186–191, 2003
106. Vats A, Shapiro R, Singh RP, Scantlebury V, Tuzuner A, Saxena M, Moritz ML, Beattie TJ, Gonwa T, Green MD, Ellis D: Quantitative viral load monitoring and cidofovir therapy for the management of BK virus-associated nephropathy in children and adults. *Transplantation* 75: 105–112, 2003
107. BJORANG O, Tveitan H, Midtvedt K, Broch LU, Scott H, Andresen PA: Treatment of polyomavirus infection with cidofovir in a renal-transplant recipient. *Nephrol Dial Transplant* 17: 2023–2025, 2002
108. Burgos D, Lopez V, Cabello M, Sola E, Gutierrez C, Garcia I, Siles J, Florez P, Gonzalez MM: Polyomavirus BK nephropathy: The effect of an early diagnosis on renal function or graft loss. *Transplant Proc* 38: 2409–2411, 2006
109. Lopez V, Sola E, Gutierrez C, Burgos D, Cabello M, Garcia I, Florez P, Lopez J, Gonzalez-Molina M: Anterior uveitis associated with treatment with intravenous cidofovir in kidney transplant patients with BK virus nephropathy. *Transplant Proc* 38: 2412–2413, 2006
110. Lalezari JP, Stagg RJ, Kuppermann BD, Holland GN,

- Kramer F, Ives DV, Youle M, Robinson MR, Drew WL, Jaffe HS: Intravenous cidofovir for peripheral cytomegalovirus retinitis in patients with AIDS. A randomized, controlled trial. *Ann Intern Med* 126: 257–263, 1997
111. Stenlund A: Initiation of DNA replication: Lessons from viral initiator proteins. *Nat Rev Mol Cell Biol* 4: 777–785, 2003
112. Leung AY, Chan MT, Yuen KY, Cheng VC, Chan KH, Wong CL, Liang R, Lie AK, Kwong YL: Ciprofloxacin decreased polyoma BK virus load in patients who underwent allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis* 40: 528–537, 2005
113. Randhawa PS: Anti-BK virus activity of ciprofloxacin and related antibiotics. *Clin Infect Dis* 41: 1366–1367, 2005
114. Portolani M, Pietrosevoli P, Cermelli C, Mannini-Palenzona A, Grossi MP, Paolini L, Barbanti-Brodano G: Suppression of BK virus replication and cytopathic effect by inhibitors of prokaryotic DNA gyrase. *Antiviral Res* 9: 205–218, 1988
115. Chandraker A, Ali S, Drachenberg CB, Wali R, Hirsch HH, DeCaprio J, Ramos E: Use of fluoroquinolones to treat BK infection in renal transplant recipients [Abstract]. *Am J Transplant* 4: 587, 2004
116. Thamboo TP, Jeffery KJ, Friend PJ, Turner GD, Roberts IS: Urine cytology screening for polyoma virus infection following renal transplantation: The Oxford experience. *J Clin Pathol* December 8, 2006 [epub ahead of print]
117. Kazatchkine MD, Kaveri SV: Immunomodulation of autoimmune and inflammatory diseases with intravenous immune globulin. *N Engl J Med* 345: 747–755, 2001
118. Puliyaanda D, Radha RK, Amet N, Hilo L, Jordan SC: IVIG contains antibodies reactive with polyoma BK virus and may represent a therapeutic option for BK nephropathy [Abstract]. *Am J Transplant* 3: 393A, 2003
119. Sener A, House AA, Jevnikar AM, Boudville N, McAlister VC, Muirhead N, Rehman F, Luke PP: Intravenous immunoglobulin as a treatment for BK virus associated nephropathy: One-year follow-up of renal allograft recipients. *Transplantation* 81: 117–120, 2006
120. Cibrik DM, O'Toole JF, Norman SP, Shah R, Johnson K, Arenas J, Meier-Kriesche HU: IVIG for the treatment of transplant BK nephropathy [Abstract]. *Am J Transplant* 3: 370, 2007
121. Jordan SC, Vo AA, Peng A, Toyoda M, Tyan D: Intravenous gammaglobulin (IVIG): A novel approach to improve transplant rates and outcomes in highly HLA-sensitized patients. *Am J Transplant* 6: 459–466, 2006
122. Orbach H, Katz U, Sherer Y, Shoenfeld Y: Intravenous immunoglobulin: Adverse effects and safe administration. *Clin Rev Allergy Immunol* 29: 173–184, 2005
123. Funk GA, Steiger J, Hirsch HH: Rapid dynamics of polyomavirus type BK in renal transplant recipients. *J Infect Dis* 193: 80–87, 2006
124. Trofe J, Roy-Chaudhury P, Gordon J, Wadih G, Maru D, Cardi MA, Succop P, Alloway RR, Khalili K, Woodle ES: Outcomes of patients with rejection post-polyomavirus nephropathy. *Transplant Proc* 37: 942–944, 2005
125. Ginevri F, Pastorino N, De Santis R, Fontana I, Sementa A, Losurdo G, Santopietro A, Perfumo F, Locatelli F, Maccario R, Azzi A, Comoli P: Retransplantation after kidney graft loss due to polyoma BK virus nephropathy: Successful outcome without original allograft nephrectomy. *Am J Kidney Dis* 42: 821–825, 2003
126. Lipshutz GS, Mahanty H, Feng S, Hirose R, Stock PG, Kang SM, Posselt AM, Freise CE: BKV in simultaneous pancreas-kidney transplant recipients: A leading cause of renal graft loss in first 2 years post-transplant. *Am J Transplant* 5: 366–373, 2005
127. Poduval RD, Meehan SM, Woodle ES, Thistlethwaite JR, Haas M, Cronin DC, Vats A, Josephson MA: Successful retransplantation after renal allograft loss to polyoma virus interstitial nephritis. *Transplantation* 73: 1166–1169, 2002
128. Ramos E, Vincenti F, Lu WX, Shapiro R, Trofe J, Stratta RJ, Jonsson J, Randhawa PS, Drachenberg CB, Papadimitriou JC, Weir MR, Wali RK: Retransplantation in patients with graft loss caused by polyoma virus nephropathy. *Transplantation* 77: 131–133, 2004
129. Womer KL, Meier-Kriesche HU, Patton PR, Dibadj K, Bucci CM, Foley D, Fujita S, Croker BP, Howard RJ, Srinivas TR, Kaplan B: Preemptive retransplantation for BK virus nephropathy: Successful outcome despite active viremia. *Am J Transplant* 6: 209–213, 2006
130. Al Jedai AH, Honaker MR, Trofe J, Egidi MF, Gaber LW, Gaber AO, Stratta RJ: Renal allograft loss as the result of polyomavirus interstitial nephritis after simultaneous kidney-pancreas transplantation: Results with kidney retransplantation. *Transplantation* 75: 490–494, 2003
131. Azzi A, De Santis R, Salotti V, Di Pietro N, Ginevri F, Comoli P: BK virus regulatory region sequence deletions in a case of human polyomavirus associated nephropathy (PVAN) after kidney transplantation. *J Clin Virol* 35: 106–108, 2006
132. Boucek P, Voska L, Saudek F: Successful retransplantation after renal allograft loss to polyoma virus interstitial nephritis. *Transplantation* 74: 1478, 2002
133. Nickleit V, Mihatsch MJ: Polyomavirus allograft nephropathy and concurrent acute rejection: A diagnostic and therapeutic challenge. *Am J Transplant* 4: 838–839, 2004
134. Randhawa P, Shapiro R: Conceptual problems in the diagnosis and therapy of acute rejection in patients with polyomavirus nephropathy. *Am J Transplant* 4: 840, 2004
135. Jahnukainen T, Malehorn D, Sun M, Lyons-Weiler J, Bigbee W, Gupta G, Shapiro R, Randhawa PS, Pelikan R, Hauskrecht M, Vats A: Proteomic analysis of urine in kidney transplant patients with BK virus nephropathy. *J Am Soc Nephrol* 17: 3248–3256, 2006
136. Drachenberg CB, Papadimitriou JC: Polyomavirus-associated nephropathy: Update in diagnosis. *Transpl Infect Dis* 8: 68–75, 2006
137. McGilvray ID, Lajoie G, Humar A, Cattral MS: Polyomavirus infection and acute vascular rejection in a kidney allograft: Coincidence or mimicry. *Am J Transplant* 3: 501–504, 2003
138. Kiberd BA: Screening to prevent polyoma virus nephropathy: A medical decision analysis. *Am J Transplant* 5: 2410–2416, 2005