

Role of Vasopressin Antagonists

Vicente E. Torres

Mayo Clinic College of Medicine and Division of Nephrology, Mayo Clinic College of Medicine, Rochester, Minnesota

Alterations in intracellular calcium homeostasis and cyclic adenosine 3',5'-phosphate likely underlie the increased cell proliferation and fluid secretion in polycystic kidney disease. Hormone receptors that affect cyclic adenosine 3',5'-phosphate and are preferentially expressed in affected tissues are logical treatment targets. There is a sound rationale for considering the arginine vasopressin V2 receptor as a target. The arginine vasopressin V2 receptor antagonists OPC-31260 and tolvaptan inhibit the development of polycystic kidney disease in cpk mice and in three animal orthologs to human autosomal recessive polycystic kidney disease (PCK rat), autosomal dominant polycystic kidney disease (Pkd2-/WS25 mice), and nephronophthisis (pcy mouse). PCK rats that are homozygous for an arginine vasopressin mutation and lack circulating vasopressin are markedly protected. Administration of V2 receptor agonist 1-deamino-8-D-arginine vasopressin to these animals completely recovers the cystic phenotype. Administration of 1-deamino-8-D-arginine vasopressin to PCK rats with normal arginine vasopressin aggravates the disease. Suppression of arginine vasopressin release by high water intake is protective. V2 receptor antagonists may have additional beneficial effects on hypertension and chronic kidney disease progression. A number of clinical studies in polycystic kidney disease have been performed or are currently active. The results of phase 2 and 2-3 studies indicate that tolvaptan seems to be safe and well tolerated in autosomal dominant polycystic kidney disease. A phase 3, placebo-controlled, double-blind study in 18- to 50-yr-old patients with autosomal dominant polycystic kidney disease and preserved renal function but relatively rapid progression, as indicated by a total kidney volume >750 ml, has been initiated.

Clin J Am Soc Nephrol 3: 1212-1218, 2008. doi: 10.2215/CJN.05281107

The polycystic kidney disease (PKD) proteins are multi-functional interacting proteins that are essential to maintain the differentiated phenotype of the tubular epithelium. Reduction in one or more of these proteins below a critical threshold results in a phenotypic switch characterized by altered protein trafficking and targeting, increased cell-matrix and reduced cell-cell adhesiveness, increased rates of proliferation and apoptosis, loss of planar polarity, and expression of a secretory phenotype. Increased cell proliferation and loss of planar polarity are responsible for cyst initiation, whereas increased fluid secretion and the persistent effects of factors that initiate cyst formation contribute to cyst growth (reviewed in reference [1]).

Molecular Mechanisms of Polycystic Kidney Disease

The proteins encoded by PKD1 (polycystin-1 [PC1]), PKD2 (polycystin-2 [PC2]), and PKHD1 (fibrocystin/polyductin [FC]) are membrane-associated proteins. PC2 is a transient receptor potential channel with high permeability to calcium (also known

as TRPP2). PC1 (also known as TRPP1) directly and FC indirectly interact with PC2 and modulate its channel activity (2-4). PC1 and FC have other functions that in turn may be modulated by PC2 and calcium. PC1, PC2, and FC are located in primary cilia (5-7). PC2 is also present in the endoplasmic reticulum, where it interacts with inositol triphosphate and ryanodine receptors (8-10). Together, these receptors are responsible for calcium release from intracellular stores. In primary cilia, the polycystin complex translates mechanical stimulation of the cilia into calcium entry, which triggers calcium-induced calcium release from the endoplasmic reticulum (Figure 1A). PC2 interacts with TRPV4, a likely component of the mechanosensory apparatus in primary cilia (11), and TRPC1, the strongest candidate component of the store-operated calcium channel (12). Reductions in the levels of PC or FC below a critical threshold disrupt intracellular calcium homeostasis (13-17). In renal tubular epithelial cells, intracellular calcium limits cAMP accumulation by inhibiting adenylyl cyclase 6 and possibly by activating phosphodiesterase 1 (18,19). In animal models of autosomal dominant (ADPKD) and recessive (ARPKD) PKD, the renal levels of cAMP are increased, likely as a result of the disruption of intracellular calcium homeostasis (20-24). cAMP stimulates chloride-driven fluid secretion (25) (Figure 1B). Whereas under normal conditions cAMP inhibits mitogen-activated protein kinase signaling and cell proliferation, in PKD or in conditions of calcium deprivation it stimulates cell proliferation in an src-, Ras-, and B-Raf-dependent manner (26-28). Cell proliferation may be further enhanced by stimulation of Erb-B by EGF-like

Published online ahead of print. Publication date available at www.cjasn.org.

Correspondence: Dr. Vicente E. Torres, Division of Nephrology and Hypertension, Mayo Clinic, 200 First Street SW, Rochester, MN 55905. Phone: 507-284-7572; Fax: 507-266-9315; E-mail: torres.vicente@mayo.edu




Figure 1. (A) Polycystin-1 (PC1), polycystin-2 (PC2), and fibrocystin/polyductin (FC) are located in primary cilia. PC2 is also present in the endoplasmic reticulum, where it interacts with inositol triphosphate (IP₃R) and ryanodine (RR) receptors. Together, these receptors are responsible for calcium release from intracellular stores. In primary cilia, the polycystin complex translates mechanical stimulation of the cilia into calcium entry, which triggers calcium-induced calcium release from the endoplasmic reticulum (ER). (B) Reductions in the levels of PC or FC below a critical threshold disrupt intracellular calcium homeostasis. Reduced calcium level at certain cellular domains enhances cAMP accumulation by increasing the activity of adenylyl cyclase 6 and possibly decreasing the activity of phosphodiesterase 1. cAMP stimulates chloride-driven fluid secretion. (C) Whereas under normal conditions cAMP inhibits mitogen-activated protein kinase signaling and cell proliferation, in PKD or in conditions of calcium deprivation it stimulates cell proliferation in an src-, Ras-, and B-raf–dependent manner. The proliferative effect of cAMP may be further enhanced by the stimulation of ErbB receptors by EGF-like factors present in cyst fluid. (D) Mammalian target of rapamycin (mTOR) is also activated in cystic epithelium, likely as a result of a disrupted tuberlin-PC1 interaction or to extracellular signal-regulated kinase- or Akt-dependent phosphorylation of tuberlin that prevents its association with hamartin and inhibits its GTPase activating function for Rheb. AC-VI, adenylyl cyclase type VI; ATP, adenosine triphosphate; CAML, calcium modulating cyclophilin ligand; CFTR, cystic fibrosis transmembrane conductance regulator; ER, endoplasmic reticulum; ERK, extracellular signal-regulated kinase; G_i, G_i-type protein; G_q, G_q-type protein; IP₃, inositol triphosphate; MEK, mitogen-activated protein kinase kinase; PDE, phosphodiesterase; PKA, protein kinase A; PKD, polycystic kidney disease; PLC, phospholipase C; R, receptor.

factors that are present in cyst fluid (Figure 1C) (29). Mammalian target of rapamycin is also activated in cystic epithelium, likely as a result of a disrupted tuberlin-polycystin-1 interaction or to extracellular signal-regulated kinase- or Akt-dependent phosphorylation of tuberlin that prevents its association with hamartin and inhibits its GTPase activating function for Rheb (30) (Figure 1D).

Hormone Receptor–Targeted Interventions

The understanding of molecular mechanisms operative in PKD has identified a number of targets for therapeutic interventions that have been effective in animal models of the disease (31). A strategy to target hormone receptors that affect cAMP is attractive, given the restricted expression of particular

receptors in certain tissues and the central role of cAMP. There are a number of reasons for considering the use of arginine vasopressin (AVP) V2 receptor antagonists. The localization of the V2 receptors in the distal nephron and collecting duct (32) corresponds to the main site of cystogenesis in ARPKD and arguably in ADPKD (33). As a result of terrestrial adaptation, tetrapods are constantly subjected to the tonic action of vasopressin on V2 receptors. AVP is the main agonist of adenylyl cyclase in freshly dissociated collecting ducts (34). Patients with PKD have increased circulating levels of AVP (35–37). Animal models of PKD also have increased circulating levels of AVP as well as upregulation of AVP- and cAMP-dependent genes such as the V2 receptor and aquaporin-2 (20–22). In the past decade, several nonpeptide, orally bioavailable V2 receptors have been extensively investigated for the treatment of euvolemic and hypervolemic hyponatremia and found to be reasonably safe (38–42).

Preclinical Studies

In 1999, Gattone *et al.* (43) reported that the V2 receptor antagonist OPC-31260 had a marked protective effect on the development of PKD in the *cpk* mouse, a model of rapidly progressive cystic disease. To extend this observation, OPC-31260 was then used in three animal models orthologous to human ARPKD (PCK rat), ADPKD (*Pkd2*^{WS25/-} mouse), and adolescent nephronophthisis (*pcy* mouse) (20,21). Renal concentrations of cAMP are significantly increased in the three models, compared with wild-type animals. In PCK rats, the administration of OPC-31260 between 3 and 10 wk or between 10 and 18 wk of age significantly reduced the renal levels of cAMP, the activation of Ras and extracellular signal-regulated kinase, and the expression of the pro-proliferative isoform of B-Raf. This was accompanied by a marked inhibition of disease development, when administered between 3 and 10 wk of age, or of disease progression, when administered between 10 and 18 wk of age, as reflected by significant reductions in kidney volume, cyst and fibrosis volumes, plasma blood urea nitrogen (BUN), and mitotic and apoptotic indices. In *Pkd2*^{WS25/-} mice, the administration of OPC-31260 lowered the renal levels of cAMP, downregulated the expression of V2 receptor- and cAMP-dependent genes (V2 receptor and aquaporin 2), and markedly inhibited the development of PKD, as reflected by lower kidney weights, cyst and fibrosis volumes, plasma BUN levels, and mitotic and apoptotic indices. OPC-31260 was also protective in the *pcy* mouse (Figure 2). Because OPC-31260 is a weak antagonist for the human V2 receptor, a derivative with a higher affinity for the human V2 receptor (tolvaptan) has been under commercial development for the treatment of hyponatremia in humans (44). This antagonist was also effective in animal models of ARPKD, ADPKD, and nephronophthisis (22,45,46). Neither OPC-31260 nor tolvaptan had a beneficial effect on the development of fibropolycystic liver disease, which is consistent with the absence of V2 receptor expression in the liver.

To determine whether the protective effect of these drugs is indeed due to V2 receptor antagonism, we decided to generate PKD rats that lack circulating AVP and determine the effect of administering the exogenous V2 receptor agonist 1-deamino-8-D-arginine vasopressin (dDAVP) (47). For this purpose, we crossed Brattleboro




Figure 2. Kidney sections from PCK rats, *Pkd2*^{-/WS25} mice, and *pcy* mice untreated (top) or treated (bottom) with OPC-31260. Sections from PCK rats and *pcy* mice are representative kidney sections. Sections from *Pkd2*^{-/WS25} mice illustrate the most severe cystic disease in the control and treatment groups.

rats, which are homozygous for a mutation in the *AVP* gene and lack circulating AVP, and PCK rats and intercrossed the F1 animals to create PCK rats with normal *AVP*, heterozygous or homozygous for an *AVP* mutation, as well as wild-type and Brattleboro controls. At 10 and 20 wk of age, PCK rats homozygous for an *AVP* mutation were almost completely protected, whereas *AVP* heterozygosity had no detectable protective effect. The administration of dDAVP between 12 and 20 wk of age completely recovered the cystic phenotype. The administration of dDAVP to PCK rats with normal *AVP* aggravated the development of the cystic disease (Figure 3).

Nagao *et al.* (48) offered PKD rats 5% sucrose in drinking water to increase water intake and suppress AVP levels. This strategy caused a marked amelioration of the renal cystic dis-




Figure 3. Representative kidney sections from 20-wk-old male (top) and female (bottom) PCK *AVP*^{+/+}, PCK *AVP*^{+/-}, and PCK *AVP*^{-/-} rats, and PCK *AVP*^{-/-} rats treated with 1-deamino-8-D-arginine vasopressin (dDAVP; 10 ng/h per 100 g body wt) using osmotic minipumps from 12 to 20 wk of age.

ease, supporting a central role for AVP and cAMP in cystic disease.

In addition to an effect on cystogenesis, V2 receptor antagonists may have beneficial effects on hypertension and chronic kidney disease (CKD) progression. In regard to hypertension, V2 receptor antagonists may have an antihypertensive effect by functionally downregulating amiloride-sensitive epithelial Na^+ channel, Na^+K^+ -ATPase, $\text{Na}^+\text{K}^+-2\text{Cl}^-$ co-transporter, and thiazide-sensitive Na^+Cl^- co-transporter, thus reducing sodium reabsorption (49–53). Consistent with this is the observation that Brattleboro rats are resistant to the development of DOCA-salt hypertension and that chronic stimulation of vasopressin V2 receptors increases BP in the healthy rat and worsens hypertension in DOCA-salt-treated animals (54,55). Conversely, an argument could be made that V2 receptor antagonists could have a prohypertensive effect by inhibiting nitric oxide production in the collecting ducts and interfering with medullary vasodilation (56).

In regard to CKD progression, Bankir and colleagues (57,58) argued that V2 receptor activation, by increasing urea recycling from the collecting duct into the loop of Henle, reduces sodium concentration at the macula densa; inhibits tubuloglomerular feedback; stimulates renin release; and results in glomerular hyperfiltration, proteinuria, and renal damage. By interfering with this chain of events, V2 receptor antagonists might have a renoprotective effect similar to that of angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers. In this regard, it is of interest that AVP deficiency and water loading inhibit CKD progression in five-sixths nephrectomized rats (59,60).

Clinical Studies

A number of clinical studies on the effect of tolvaptan in ADPKD have been completed or are currently active under the Tolvaptan Efficacy and Safety in Management of PKD and Outcomes (TEMPO) program. Phase 2a studies (248 and 249) have been completed, whereas two multicenter studies (250 and 251) and a single-center mechanistic study (260) are currently active.

In TEMPO 248, ascending single doses of tolvaptan (15, 30, 60, and 120 mg) were administered every 3 d and showed a dosage-dependent increase in urine output (61–63). Hypostenuria, a surrogate marker for vasopressin suppression, was sustained during 16 h after the administration of 30, 60, or 120 mg, but urine osmolality increased above 300 mOsm/kg in most patients 16 to 24 h after receiving 30 or 60 mg of tolvaptan. In TEMPO 249, different split doses of tolvaptan (15/15, 30/placebo, 30/15, 30/30 mg) were administered for 5 d at 8 a.m. and 4 p.m. to groups of patients (parallel-arm design) (61–63). Split-dose administration was more effective than a single dose in achieving sustained hypostenuria, and the mean urine output on the fifth day was lower than that on the first day of treatment, 4 to 6 compared with 6 to 7 L per 24 h.

Study 250 is an open-label study that consists of a titration phase and a fixed-dose phase (64). The initial dose of tolvaptan in the titration phase was 30/15 mg at 8 a.m. and 4 p.m., with the option to down-titrate to 15/15 if not tolerated. The dosage

was increased at weekly intervals to 45/15, 60/30, and 90/30 mg when tolerated. During the titration phase 96, 61, and 46% of the patients said that they could tolerate 45/15, 60/30, and 90/30 mg for the rest of their life. Urine osmolalities before the morning dose of tolvaptan were >300 mOsm/kg in 20 to 30% of patients who were taking 45/15, 60/30, or 90/30 mg/d, pointing to the difficulty in achieving sustained hypostenuria in all of the patients. On the basis of the results of the titration phase, participants were randomly assigned to high (60/30 mg) and low (45/15 mg) dosages of tolvaptan for the fixed phase that is planned to last 3 yr. At the time of a published interim report, at approximately half point into the trial, the mean premorning dose urine osmolality was <300 mOsm/kg. There was an initial, slight increase in serum creatinine that later declined toward baseline. Serum BUN significantly decreased, whereas there was a slight but significant increase in uric acid. Both systolic and diastolic BP tended to decrease with time, but this change is difficult to interpret in the absence of a control group. Five patients withdrew from the study, in four cases because of adverse events and one case because of noncompliance. One of the adverse events that led to discontinuation was deemed to be related to the drug, an increase in serum creatinine from 1.4 to 1.7 mg/dl that was rapidly reversible after discontinuation of the drug. The other adverse events that led to discontinuation included periorbital swelling, transient ischemic attack, and a benign pituitary microadenoma.

TEMPO 251 is a phase 3, placebo-controlled, double-blind study in 18- to 50-yr-old patients with preserved renal function but relatively rapid progression, as indicated by a total kidney volume >750 ml. The primary end point is renal volume change by magnetic resonance, and the secondary end point is time to multiple progression events. The duration of treatment is 3 yr. After a screening and a randomization visit, the masked medication is increased at weekly intervals from 45/15 to 90/30 mg/d if tolerated. After the titration phase, visits occur every 4 mo and magnetic resonance scans are obtained yearly.

Conclusions

Extensive animal studies suggest that AVP is a powerful modulator of cystogenesis, that inhibition of renal cAMP production accounts for the protective effect of V2 receptor antagonists, and that these drugs may afford additional benefits on hypertension and CKD progression. These studies provide a strong rationale for clinical trials using V2 receptor antagonists in ADPKD. Phase 2a studies have been completed. Two multicenter studies (an open-label study and a phase 3, placebo-controlled, double-blind study) are currently active.

Disclosures

None.

References

1. Torres VE, Harris PC: Mechanisms of disease: Autosomal

- dominant and recessive polycystic kidney diseases. *Nat Clin Pract Nephrol* 2: 40–54, 2006
2. Qian F, Germino FJ, Cai Y, Zhang X, Somlo S, Germino GG: PKD1 interacts with PKD2 through a probable coiled-coil domain. *Nat Genet* 16: 179–183, 1997
 3. Tsiokas L, Kim E, Arnould T, Sukhatme VP, Walz G: Homo- and heterodimeric interactions between the gene products of PKD1 and PKD2. *Proc Natl Acad Sci U S A* 94: 6965–6970, 1997
 4. Wu Y, Dai XQ, Li Q, Chen CX, Mai W, Hussain Z, Long W, Montalbetti N, Li G, Glynne R, Wang S, Cantiello HF, Wu G, Chen XZ: Kinesin-2 mediates physical and functional interactions between polycystin-2 and fibrocystin. *Hum Mol Genet* 15: 3280–3292, 2006
 5. Pazour GJ, San Agustin JT, Follit JA, Rosenbaum JL, Witman GB: Polycystin-2 localizes to kidney cilia and the ciliary level is elevated in orpk mice with polycystic kidney disease. *Curr Biol* 12: R378–R380, 2002
 6. Ward CJ, Yuan D, Masyuk TV, Wang X, Punyashtithi R, Whelan S, Bacallao R, Torra R, LaRusso NF, Torres VE, Harris PC: Cellular and subcellular localization of the ARPKD protein: Fibrocystin is expressed on primary cilia. *Hum Mol Genet* 12: 2703–2710, 2003
 7. Yoder BK, Hou X, Guay-Woodford LM: The polycystic kidney disease proteins, polycystin-1, polycystin-2, polaris, and cystin, are co-localized in renal cilia. *J Am Soc Nephrol* 13: 2508–2516, 2002
 8. Anyatonwu GI, Estrada M, Tian X, Somlo S, Ehrlich BE: Regulation of ryanodine receptor-dependent calcium signaling by polycystin-2. *Proc Natl Acad Sci U S A* 104: 6454–6459, 2007
 9. Koulen P, Cai Y, Geng L, Maeda Y, Nishimura S, Witzgall R, Ehrlich BE, Somlo S: Polycystin-2 is an intracellular calcium release channel. *Nat Cell Biol* 4: 191–197, 2002
 10. Li Y, Wright JM, Qian F, Germino GG, Guggino WB: Polycystin 2 interacts with type I inositol 1,4,5-trisphosphate receptor to modulate intracellular Ca²⁺ signaling. *J Biol Chem* 280: 41298–41306, 2005
 11. Doerken M, Kotsis F, Kottgen M, Braeg S, Walz G, Kuehn W: Suppression of ciliary flow sensing through inducible knock-down of the PKD2 interactor TRPV4 [Abstract]. *J Am Soc Nephrol* 18: 13A, 2007
 12. Tsiokas L, Arnould T, Shu C, Kim E, Walz G, Sukhatme VP: Specific association of the gene product of PKD₂ with the TRPC1 channel. *Proc Natl Acad Sci U S A* 96: 3934–3939, 1999
 13. Nauli SM, Alenghat FJ, Luo Y, Williams E, Vassilev P, Li X, Elia AE, Lu W, Brown EM, Quinn SJ, Ingber DE, Zhou J: Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet* 33: 129–137, 2003
 14. Praetorius HA, Frokiaer J, Nielsen S, Spring KR: Bending the primary cilium opens Ca²⁺-sensitive intermediate-conductance K⁺ channels in MDCK cells. *J Membr Biol* 191: 193–200, 2003
 15. Qian Q, Hunter LW, Li M, Marin-Padilla M, Prakash YS, Harris PC, Somlo S, Torres VE, Sieck GC: Pkd2 haploinsufficiency alters intracellular calcium in vascular smooth muscle cells. *Hum Mol Genet* 12: 1875–1880, 2003
 16. Ahrabi AK, Terryn S, Valenti G, Caron N, Serradeil-Le Gal C, Raufaste D, Nielsen S, Horie S, Verbavatz JM, Devuyst O: PKD1 haploinsufficiency causes a syndrome of inappropriate antidiuresis in mice. *J Am Soc Nephrol* 18: 1740–1753, 2007
 17. Yang J, Zhang S, Zhou Q, Guo H, Zhang K, Zheng R, Xiao C: PKHD1 gene silencing may cause cell abnormal proliferation through modulation of intracellular calcium in autosomal recessive polycystic kidney disease. *J Biochem Mol Biol* 40: 467–474, 2007
 18. Chabardes D, Firsov D, Aarab L, Clabecq A, Bellanger AC, Siaume-Perez S, Elalouf JM: Localization of mRNAs encoding Ca²⁺-inhibitable adenylyl cyclases along the renal tubule. Functional consequences for regulation of the cAMP content. *J Biol Chem* 271: 19264–19271, 1996
 19. Dousa TP: Cyclic-3',5'-nucleotide phosphodiesterase isozymes in cell biology and pathophysiology of the kidney. *Kidney Int* 55: 29–62, 1999
 20. Gattone VH, Wang X, Harris PC, Torres VE: Inhibition of renal cystic disease development and progression by a vasopressin V2 receptor antagonist. *Nat Med* 9: 1323–1326, 2003
 21. Torres VE, Wang X, Qian Q, Somlo S, Harris PC, Gattone VH: Effective treatment of an orthologous model of autosomal dominant polycystic kidney disease. *Nat Med* 10: 363–364, 2004
 22. Wang X, Gattone II, VH, Harris PC, Torres VE: Effectiveness of vasopressin V2 receptor antagonists OPC-31260 and OPC-41061 on polycystic kidney disease development in the PCK rat. *J Am Soc Nephrol* 16: 846–851, 2005
 23. Yamaguchi T, Nagao S, Kasahara M, Takahashi H, Grantham J: Renal accumulation and excretion of cyclic adenosine monophosphate in a murine model of slowly progressive polycystic kidney disease. *Am J Kidney Dis* 30: 703–709, 1997
 24. Smith LA, Bukanov NO, Husson H, Russo RJ, Barry TC, Taylor AL, Beier DR, Ibraghimov-Beskrovnaya O: Development of polycystic kidney disease in juvenile cystic kidney mice: insights into pathogenesis, ciliary abnormalities, and common features with human disease. *J Am Soc Nephrol* 17: 2821–2831, 2006
 25. Grantham JJ: Lillian Jean Kaplan International Prize for advancement in the understanding of polycystic kidney disease. Understanding polycystic kidney disease: A systems biology approach. *Kidney Int* 64: 1157–1162, 2003
 26. Yamaguchi T, Nagao S, Wallace DP, Belibi FA, Cowley BD, Pelling JC, Grantham JJ: Cyclic AMP activates B-Raf and ERK in cyst epithelial cells from autosomal-dominant polycystic kidneys. *Kidney Int* 63: 1983–1994, 2003
 27. Hanaoka K, Guggino W: cAMP regulates cell proliferation and cyst formation in autosomal polycystic kidney disease cells. *J Am Soc Nephrol* 11: 1179–1187, 2000
 28. Yamaguchi T, Wallace DP, Magenheimer BS, Hempson SJ, Grantham JJ, Calvet JP: Calcium restriction allows cAMP activation of the B-Raf/ERK pathway, switching cells to a cAMP-dependent growth-stimulated phenotype. *J Biol Chem* 279: 40419–40430, 2004
 29. Sweeney WE Jr, Avner ED: Molecular and cellular pathophysiology of autosomal recessive polycystic kidney disease (ARPKD). *Cell Tissue Res* 326: 671–685, 2006
 30. Shillingford JM, Murcia NS, Larson CH, Low SH, Hedgepeth R, Brown N, Flask CA, Novick AC, Goldfarb DA, Kramer-Zucker A, Walz G, Piontek KB, Germino GG, Weimbs T: The mTOR pathway is regulated by polycystin-1, and its inhibition reverses renal cystogenesis in poly-

- cystic kidney disease. *Proc Natl Acad Sci U S A* 103: 5466–5471, 2006
31. Torres VE, Harris PC, Pirson Y: Autosomal dominant polycystic kidney disease. *Lancet* 369: 1287–1301, 2007
 32. Mutig K, Paliège A, Kahl T, Jons T, Muller-Esterl WP, Bachmann S: Vasopressin V2 receptor expression along rat, mouse, and human renal epithelia with focus on TAL. *Am J Physiol Renal Physiol* 293: F1166–F1177, 2007
 33. Torres VE: Vasopressin antagonists in polycystic kidney disease. *Kidney Int* 68: 2405–2418, 2005
 34. Yasuda G, Jeffries WB: Regulation of cAMP production in initial and terminal inner medullary collecting ducts. *Kidney Int* 54: 80–86, 1998
 35. Danielsen H, Pedersen EB, Nielsen AH, Herlevsen P, Kornerup HJ, Posborg V: Expansion of extracellular volume in early polycystic kidney disease. *Acta Med Scand* 219: 399–405, 1986
 36. Michalski A, Grzeszczak W: The effect of hypervolemia on electrolyte level and level of volume regulating hormones in patients with autosomal dominant polycystic kidney disease [in Polish]. *Pol Arch Med Wewn* 96: 329–343, 1996
 37. Seeman T, Dusek J, Vondrak K, Blahova K, Simkova E, Kreisinger J, Dvorak P, Kyncl M, Hribal Z, Janda J: Renal concentrating capacity is linked to blood pressure in children with autosomal dominant polycystic kidney disease. *Physiol Res* 53: 629–634, 2004
 38. Ali F, Guglin M, Vaitkevicius P, Ghali JK: Therapeutic potential of vasopressin receptor antagonists. *Drugs* 67: 847–858, 2007
 39. Gheorghide M, Konstam MA, Burnett JC Jr, Grinfeld L, Maggioni AP, Swedberg K, Udelson JE, Zannad F, Cook T, Ouyang J, Zimmer C, Orlandi C: Short-term clinical effects of tolvaptan, an oral vasopressin antagonist, in patients hospitalized for heart failure: The EVEREST Clinical Status Trials. *JAMA* 297: 1332–1343, 2007
 40. Greenberg A, Verbalis JG: Vasopressin receptor antagonists. *Kidney Int* 69: 2124–2130, 2006
 41. Konstam MA, Gheorghide M, Burnett JC Jr, Grinfeld L, Maggioni AP, Swedberg K, Udelson JE, Zannad F, Cook T, Ouyang J, Zimmer C, Orlandi C: Effects of oral tolvaptan in patients hospitalized for worsening heart failure: The EVEREST Outcome Trial. *JAMA* 297: 1319–1331, 2007
 42. Schrier RW, Gross P, Gheorghide M, Berl T, Verbalis JG, Czerwiec FS, Orlandi C: Tolvaptan, a selective oral vasopressin V₂-receptor antagonist, for hyponatremia. *N Engl J Med* 355: 2099–2112, 2006
 43. Gattone VH, Maser RL, Tian C, Rosenberg JM, Branden MG: Developmental expression of urine concentration-associated genes and their altered expression in murine infantile-type polycystic kidney disease. *Dev Genet* 24: 309–318, 1999
 44. Yamamura Y, Nakamura S, Itoh S, Hirano T, Onogawa T, Yamashita T, Yamada Y, Tsujimae K, Aoyama M, Kotosai K, Ogawa H, Yamashita H, Kondo K, Tominaga M, Tsujimoto G, Mori T: OPC-41061, a highly potent human vasopressin V₂-receptor antagonist: Pharmacological profile and aquaretic effect by single and multiple oral dosing in rats. *J Pharmacol Exp Ther* 287: 860–867, 1998
 45. Gattone VH, Kinne Q, Torres VE: Efficacy of OPC-41061 in the treatment of murine nephronophthisis [Abstract]. *J Am Soc Nephrol* 16: 138A, 2005
 46. Wang S, Gattone VH, Somlo S, Harris PC, Torres VE: Effectiveness of OPC-41061 on polycystic kidney disease development in Pkd2WS25/– [Abstract]. *J Am Soc Nephrol* 16: 361A, 2005
 47. Wang X, Wu Y, Ward CJ, Harris PC, Torres VE: Vasopressin directly regulates cyst growth in the PCK rat. *J Am Soc Nephrol* 19: 102–108, 2008
 48. Nagao S, Kazuhiro N, Katsuyama M, Kurahashi H, Marunouchi T, Takahashi H, Wallace DP: Increased water intake decreases progression of polycystic kidney disease in the PCK rat. *J Am Soc Nephrol* 17: 228–235, 2006
 49. Kim GH, Ecelbarger CA, Mitchell C, Packer RK, Wade JB, Knepper MA: Vasopressin increases Na-K-2Cl cotransporter expression in thick ascending limb of Henle's loop. *Am J Physiol* 276: F96–F103, 1999
 50. Nicco C, Wittner M, DiStefano A, Jounier S, Bankir L, Bouby N: Chronic exposure to vasopressin upregulates ENaC and sodium transport in the rat renal collecting duct and lung. *Hypertension* 38: 1143–1149, 2001
 51. Sauter D, Fernandes S, Goncalves-Mendes N, Boulkroun S, Bankir L, Loffing J, Bouby N: Long-term effects of vasopressin on the subcellular localization of ENaC in the renal collecting system. *Kidney Int* 69: 1024–1032, 2006
 52. Gonin S, Deschenes G, Roger F, Bens M, Martin PY, Carpentier JL, Vandewalle A, Doucet A, Feraille E: Cyclic AMP increases cell surface expression of functional Na,K-ATPase units in mammalian cortical collecting duct principal cells. *Mol Biol Cell* 12: 255–264, 2001
 53. Ortiz PA: cAMP increases surface expression of NKCC2 in rat thick ascending limbs: Role of VAMP. *Am J Physiol Renal Physiol* 290: F608–F616, 2006
 54. Fernandes S, Bruneval P, Hagege A, Heudes D, Ghostine S, Bouby N: Chronic V2 vasopressin receptor stimulation increases basal blood pressure and exacerbates deoxycorticosterone acetate-salt hypertension. *Endocrinology* 143: 2759–2766, 2002
 55. Intengan HD, He G, Schiffrin EL: Effect of vasopressin antagonism on structure and mechanics of small arteries and vascular expression of endothelin-1 in deoxycorticosterone acetate salt hypertensive rats. *Hypertension* 32: 770–777, 1998
 56. Cowley AW Jr, Skelton MM, Kurth TM: Effects of long-term vasopressin receptor stimulation on medullary blood flow and arterial pressure. *Am J Physiol* 275: R1420–R1424, 1998
 57. Bankir L: Antidiuretic action of vasopressin: Quantitative aspects and interaction between V1a and V2 receptor-mediated effects. *Cardiovasc Res* 51: 372–390, 2001
 58. Bardoux P, Bichet DG, Martin H, Gallois Y, Marre M, Arthus MF, Lonergan M, Ruel N, Bouby N, Bankir L: Vasopressin increases urinary albumin excretion in rats and humans: involvement of V2 receptors and the renin-angiotensin system. *Nephrol Dial Transplant* 18: 497–506, 2003
 59. Bouby N, Bachmann S, Bichet D, Bankir L: Effect of water intake on the progression of chronic renal failure in the 5/6 nephrectomized rat. *Am J Physiol* 258: F973–F979, 1990
 60. Bouby N, Hassler C, Bankir L: Contribution of vasopressin to progression of chronic renal failure: Study in Brattleboro rats. *Life Sci* 65: 991–1004, 1999
 61. Chapman AB, Torres VE, Grantham JJ, Shoaf SS, Ouyang JJ, Czerwiec FS: A phase IIB pilot study of the safety and efficacy of tolvaptan, a vasopressin V2 receptor antagonist

- (V2RA) in patients with ADPKD [Abstract]. *J Am Soc Nephrol* 16: 68A, 2005
62. Grantham JJ, Chapman AB, Torres VE, Ouyang J, Shoaf SE, Czerwiec FS: Acute and chronic osmotic stress after vasopressin V2 receptor inhibition with tolvaptan in ADPKD [Abstract]. *J Am Soc Nephrol* 16: 361A, 2005
63. Torres VE, Wang X, Ward CJ, Grantham JJ, Chapman AB, Ouyang J, Shoaf SE, Czerwiec FS: Urine aquaporin 2 and cyclic AMP responses to tolvaptan administration in autosomal dominant polycystic kidney disease [Abstract]. *J Am Soc Nephrol* 16: 361A, 2005
64. Torres VE, Grantham JJ, Chapman AB, Watnick T, Kedzierski K, Ouyang J, Orlandi C, Czerwiec FS, Investigators, T: Phase 2 open-label study to determine safety, tolerability and efficacy of split-dose tolvaptan in ADPKD. *J Am Soc Nephrol* 18: 361–362A, 2007