

# Inflammation in Renal Transplantation

Sadollah Abedini,\* Ingar Holme,<sup>†</sup> Winfried März,<sup>‡</sup> Gisela Weihrauch,<sup>§</sup> Bengt Fellström,<sup>||</sup> Alan Jardine,<sup>¶</sup> Edward Cole,\*\* Bart Maes,<sup>††</sup> Hans-Hellmut Neumayer,<sup>‡‡</sup> Carola Grönhagen-Riska,<sup>§§</sup> Patrice Ambühl,<sup>|||</sup> and Hallvard Holdaas<sup>¶¶</sup> on behalf of the ALERT study group.

\*Renal Section, Department of Medicine, Toensberg County Hospital, Toensberg, Norway; <sup>†</sup>Department of Preventive Medicine and Centre for Clinical Research, Oslo University Hospital Ullevaal, Oslo, Norway; <sup>‡</sup>Synlab for Medizinisches Versorgungszentrum für Labordiagnostik Heidelberg, Heidelberg, Germany; <sup>§</sup>Clinical Institute of Medical and Chemical Laboratory Diagnosis, Medical University of Graz, Graz, Austria; <sup>||</sup>Department of Medical Science, Renal Unit, University Hospital, Uppsala, Sweden; <sup>¶</sup>Department of Medicine and Therapeutics, Western Infirmary Hospital, Glasgow, United Kingdom; \*\*University Health Network, University of Toronto, Ontario, Canada; <sup>††</sup>Medical Department, Heilig Hartziekenhuis, Roeselare, Belgium; <sup>‡‡</sup>Universitätsklinikum Charité, Berlin, Germany; <sup>§§</sup>University Hospital, Helsinki, Finland; <sup>|||</sup>University Hospital, Zürich, Switzerland; and <sup>¶¶</sup>Department of Medicine, Oslo University Hospital Rikshospitalet, Oslo, Norway

**Background and objectives:** Renal transplant recipients experience premature cardiovascular disease and death. The association of inflammation, all-cause mortality, and cardiovascular events in renal transplant recipients has not been examined in a large prospective controlled trial.

**Design, setting, participants, & measurements:** ALERT was a randomized, double-blind, placebo-controlled study of the effect of fluvastatin on cardiovascular and renal outcomes in 2102 renal transplant recipients. Patients initially randomized to fluvastatin or placebo in the 5- to 6-yr trial were offered open-label fluvastatin in a 2-yr extension to the original study. The association between inflammation markers, high-sensitivity C-reactive protein (hsCRP), and IL-6 on cardiovascular events and all-cause mortality was investigated.

**Results:** The baseline IL-6 value was  $2.9 \pm 1.9$  pg/ml ( $n = 1751$ ) and that of hsCRP was  $3.8 \pm 6.7$  mg/L ( $n = 1910$ ). After adjustment for baseline values for established risk factors, the hazard ratios for a major cardiac event and all-cause mortality for IL-6 were 1.08 [95% confidence interval (CI), 1.01 to 1.15,  $P = 0.018$ ] and 1.11 (95% CI, 1.05 to 1.18,  $P < 0.001$ ), respectively. The adjusted hazard ratio for hsCRP for a cardiovascular event was 1.10 (95% CI, 1.01 to 1.20,  $P = 0.027$ ) and for all-cause mortality was 1.15 (95% CI, 1.06 to 1.1.25,  $P = 0.049$ ).

**Conclusions:** The inflammation markers IL-6 and hsCRP are independently associated with major cardiovascular events and all-cause mortality in renal transplant recipients.

*Clin J Am Soc Nephrol* 4: 1246–1254, 2009. doi: 10.2215/CJN.00930209

**A**lthough cholesterol lowering therapy with statins has reduced the incidence of cardiac events in renal transplant recipients, premature cardiovascular events and premature death is still a major concern in this population (1,2). The prevalence of traditional cardiovascular risk factors cannot fully explain the increased incidence of cardiovascular events, and several reports have emphasized the role of non-traditional cardiovascular risk factors (3,4).

Inflammation and activation of the immune system may play important roles in atherogenesis (5,6). Cytokines induce production of IL-6 from various tissues, including the liver, and increase downstream mediators such as C-reactive protein

(CRP) (7). Renal transplant recipients, by the process of receiving an allograft, have an additional activation of their immune system (8–11).

In the general population, CRP is regarded as a risk factor for cardiovascular events and all-cause mortality (12,13). In patients without hyperlipidemia but with elevated high-sensitivity CRP (hsCRP), it was recently demonstrated that rosuvastatin significantly reduced the incidence of major cardiovascular events (14).

In patients with chronic renal failure, an association between malnutrition, inflammation, and atherosclerosis has been demonstrated (15). In dialysis patients CRP has been shown to be a marker of all-cause and cardiovascular death (16). It has also been speculated that the better graft survival achieved with preemptive transplantation might be due to less exposure to inflammation markers associated with dialysis (17,18). In renal transplant recipients there are few studies examining the association between inflammatory markers and all-cause mortality,

Received February 10, 2009. Accepted May 6, 2009.

Published online ahead of print. Publication date available at [www.cjasn.org](http://www.cjasn.org).

**Correspondence:** Sadollah Abedini, Toensberg County Hospital, Medical Department, Renal Section, Post Box 2168, 3103 Toensberg, Norway. Phone: +47-3334-2000; Fax: +47-3334-3993; E-mail: [Sadollah.abedini@gmail.com](mailto:Sadollah.abedini@gmail.com)

cardiac events, or cerebrovascular events. A small cross-sectional study indicated that renal transplant patients with cardiovascular disease (CVD) had higher CRP than patients without CVD (19). Another retrospective study indicated that elevated pretransplant CRP was associated with all-cause and cardiovascular mortality (20). Ducloux *et al.* followed a small cohort of renal transplant patients and reported that elevated CRP predicted coronary events (21). In a prospective registry-based study, Winkelmayr and colleagues showed that kidney transplant patients with a CRP above 5 mg/dl had an increased mortality compared with patients below that threshold (22), and in a subsequent follow-up paper the authors demonstrated a J-shaped association between hsCRP (below 5 mg/L) and mortality (23). There are no reports of assessment of an association between hsCRP and/or IL-6 and major cardiovascular events or all-cause mortality in prospectively controlled trials in renal transplant patients. We therefore performed a *post hoc* analysis of the ALERT trial examining the predictive value of the inflammation markers hsCRP and IL-6 for cardiovascular events and all-cause mortality in renal transplant recipients.

## Materials and Methods

### Study Design

The ALERT study design and baseline data have been described previously (1). Briefly, ALERT was a randomized, double-blind, placebo-controlled study of the effect of fluvastatin, 40 to 80 mg daily, on cardiovascular and renal outcomes in renal transplant recipients over a follow-up period of 5 to 6 yr. Eligible patients were men and women aged 30 to 75 yr who had received a renal transplant more than 6 mo before enrolment and with a total serum cholesterol concentration between 4.0 and 9.0 mmol/L (155 to 348 mg/dl). Patients with a history of myocardial infarction more than 6 mo before randomization could be enrolled if their total cholesterol levels ranged from 4.0 to 7.0 mmol/L (155 to 270 mg/dl). Of 1787 patients who completed ALERT, 1652 (92%) accepted open-label fluvastatin, 80 mg daily, in a 2-yr extension of the original study. Mean total follow-up time in the extension study was 6.7 yr. The study adhered to the International Conference on Harmonization guidelines for Good Clinical Practice and in accordance with the Declaration of Helsinki. All participants provided written informed consent, and the ethics committee at each participating center approved the trial.

Study patients were seen at 1.5 mo after randomization and at 6-mo intervals thereafter. Clinical status was assessed at each visit. Laboratory measurements of lipids, serum creatinine, creatine kinase, and hepatic enzymes were performed at a central laboratory (CRL, Medinet, Breda, The Netherlands).

IL-6 was measured by a human IL-6 immunoassay (R&D Systems, Inc., Minneapolis, MN, USA) and hsCRP by using an immunoturbometric analysis (Roche Diagnostics GmbH, Mannheim, Germany). Measurements of IL-6 and hsCRP were performed on blood samples at baseline and were not followed longitudinally during the study.

An independent critical events committee blinded to treatment assignment reviewed all primary and secondary endpoints for adjudication. All analyses were based on the committee's classification of endpoints, which was agreed on by consensus or by majority.

### Statistical Analyses

SPSS version 16.0 (SPSS, Inc.) was used for statistical analyses. For normally distributed variables, mean, and SD are presented. hsCRP and IL-6 values were not normally distributed and showed a marked

tail to the right. Logarithmic transformation showed a strictly bell shaped distribution of both variables and these transformed values were used for the statistical analysis. A Cox proportional hazard model was used to analyze the relationship between risk factors and time to event for all endpoints. All covariates were carefully examined and fulfilled assumptions of proportionality of hazard ratios in the Cox hazard models. Univariate and multivariate analysis were then carried out to study how risk factors associated with the risk of all-cause mortality, major cardiac events (MACE), and cerebrovascular (CBV) events. Corresponding hazard ratios for group comparisons were calculated with 95% confidence limits.

## Results

### Baseline Characteristics

Basic patient and demographic data in ALERT and the ALERT extension have been previously published (2). Patients were transplanted at a mean of 4.5 yr before randomization into the trial. Both treatments groups in the ALERT trial were well balanced with regard to baseline demographics and renal characteristics.

In this *post hoc* analysis, we separated the patients into those with available baseline hsCRP and IL-6 values. The patients' demographic characteristics based on this separation are summarized in Table 1.

The number of patients with hsCRP ( $n = 1910$ ) and IL-6 ( $n = 1751$ ) at baseline differed, and baseline data are therefore shown separately for each cohort. However, baseline characteristics between the two cohorts were practically identical. There were no differences in proportion of patients with diabetes, coronary heart disease, current smokers, and transplant type between the groups (Table 1). hsCRP and IL-6 were correlated ( $r = 0.332$ ,  $P < 0.001$ ).

### Endpoints

hsCRP and IL-6 increased with increasing body mass index (BMI) and both were significantly correlated with BMI ( $r = 0.18$ ,  $P < 0.001$ ). However, BMI was not a significant risk factor for all-cause death and MACE in univariate analysis.

Further analysis by Kaplan–Meier survival method did not show any significant differences in occurrence of all-cause death or MACE by stratifying patients by BMI (normal BMI 18.5 to 25,  $n = 890$ , 46%; overweight BMI 25 to 30,  $n = 684$ , 35%; obese BMI  $\geq 30$ ,  $n = 268$ , 14%).

In the hsCRP cohorts, there were 176 CBV events, 287 cardiac events, and 354 deaths. Corresponding numbers for the IL-6 cohort were 166, 255, and 308, respectively. The prevalence of the study endpoints within the quartiles of hsCRP and IL-6 are summarized in Table 2 a and b, and demonstrate a comparable incidence of all-cause mortality and MACE.

In both cohorts of patients there were significantly more cardiovascular events and all-cause death in the fourth quartile compared with the first quartile, with the exception of CBV events in the hsCRP cohort. We also performed separate analyses for deaths in patients with a baseline hsCRP below 5 mg/L (23), and we divided this population in quartiles (Figure 1d). Also in this subset of patients there was a successive increase in deaths by increasing hsCRP quartiles. There was a consistent increase in all endpoints from first to fourth quartile.

Table 1. Patients demographic and baseline characteristics in patients with available baseline hsCRP and IL-6<sup>a</sup>

Baseline Variables <sup>b</sup>	Patients with hsCRP Data (n = 1910)	Patients with IL-6 Data (n = 1751)
Cholesterol (mmol/L)	6.48 (1.13)	6.52 (1.13)
HDL cholesterol (mmol/L)	1.34 (0.45)	1.34 (0.45)
LDL cholesterol (mmol/L)	4.18 (1.02)	4.19 (1.01)
Apolipoprotein B (mg/dl)	117.97 (25.88)	117.08 (25.63)
Triglycerides (mmol/L)	2.22 (1.39)	2.21 (1.39)
Age at baseline (yr)	49.7 (10.0)	49.5 (11.0)
Total time on RRT (mo)	89.0 (57.9)	89.2 (57.8)
Systolic BP (mmHg)	144.0 (19.0)	145.0 (19.0)
Diastolic BP (mmHg)	86.0 (10.0)	86.0 (10.0)
Diabetes mellitus	361 (19.0)	327 (18.7)
Body mass index (kg/m <sup>2</sup> )	25.7 (4.3)	25.6 (4.18)
Serum creatinine (μmol/L)	145.0 (53.0)	145.0 (53.0)
Serum glucose (mg/dl)	93.4 (51.5)	93.0 (51.0)
PTH (pg/ml)	51.65 (66.6)	50.72 (65.07)
hsCRP (mg/L)	3.82 (6.71)	–
IL-6 pg/ml	–	2.92 (1.87)
Coronary heart disease N (%)	175 (9.2)	145 (8.3)
Diabetes mellitus N (%)	361 (19)	327 (18.7)
Smoking (current) N (%)	358 (18.8)	326 (18.6)
Living donor transplant N (%)	427 (22.4)	400 (22.8)
Cadaveric donor transplant N (%)	1477 (77.5)	1351 (77.2)
MACE N (%)	287 (15.1)	256 (14.6)
All-cause death N (%)	354 (18.6)	309 (17.6)

<sup>a</sup>Data are expressed as mean (SD) unless otherwise indicated.

<sup>b</sup>hsCRP, high-sensitivity C-reactive protein; RRT, renal replacement therapy; BP, blood pressure; PTH, parathyroid hormone; MACE, major adverse cardiovascular events.

Table 2a. Occurrence of all-cause death, MACE, and CBV events by quartiles of hsCRP<sup>a</sup>

Occurrence	Quartile 1 (<0.63)	Quartile 2 (0.63 to 1.53)	Quartile 3 (1.54 to 3.68)	Quartile 4 (≥3.69)
All-cause death	476/60 (12.3)	479/79 (16.5)	479/93 (19.4)	476/122 (25.6)
patients/events (%)	–	P = 0.088	P = 0.006	P < 0.001
MACE	476/52 (10.9)	479/71 (14.8)	479/74 (15.4)	476/90 (18.9)
patients/events (%)	–	P = 0.056	P = 0.035	P < 0.001
CBV events	476/33 (6.9)	479/49 (10.2)	479/52 (10.9)	476/42 (8.8)
patients/events (%)	–	P = 0.052	P = 0.025	P = 0.153

<sup>a</sup>Data expressed as number of patients in quartile/number of events (%). Significance versus first quartile. CBV events, cerebrovascular events.

Table 2b. Occurrence of all-cause death, MACE, and CBV events by quartiles of IL-6<sup>a</sup>

Occurrence	Quartile 1 (<1.57)	Quartile 2 (1.57 to 2.38)	Quartile 3 (2.39 to 3.74)	Quartile 4 (≥3.75)
All-cause death	444/46 (10.4)	428/60 (14.0)	437/70 (16.0)	440/132 (30.0)
patients/events (%)	–	P = 0.117	P = 0.012	P < 0.001
MACE	444/45 (10.1)	428/58 (13.6)	437/65 (14.9)	440/87 (19.8)
patients/events (%)	–	P = 0.134	P = 0.028	P < 0.001
CBV events	444/25 (5.6)	428/34 (7.9)	437/46 (10.5)	440/58 (13.2)
patients/events	–	P = 0.172	P = 0.005	P < 0.000

<sup>a</sup>Data expressed as number of patients in quartile/number of events (%). Significance versus first quartile.




Figure 1. Association between (A) all-cause death, (B) major cardiovascular events, and (C) cerebrovascular events for quartiles of high-sensitivity C-reactive protein (hsCRP) over the whole range of hsCRP. (D) Association between all-cause death, (E) major cardiovascular events, and (F) cerebrovascular events for quartiles of hsCRP below 5 mg/dl. *x*-axis represents follow-up time in years. *y*-axis represents cumulative survival.

### Risk Factor Analyses

Exploratory risk factor analysis was carried out on many potentially important risk factors for all-cause death, MACE, and CBV events (Tables 3 and 4).

On the basis of univariate analysis using Cox regression, diabetes mellitus, smoking, previous coronary heart disease, hsCRP, age, serum creatinine, and systolic blood pressure (BP), were significant risk factors for all causes of death and MACE in the hsCRP cohort. Gender and LDL cholesterol were significant univariate risk factors for MACE but not for all-cause death in the hsCRP cohort.

For all types of CBV events, diabetes mellitus, previous coronary heart disease, age, and systolic BP were significant univariate risk factors.

On the basis of multivariate Cox regression including all of the above risk factors, diabetes mellitus, smoking, age, previous coronary heart disease, hsCRP, and serum creatinine were significant for all causes of death and MACE in the hsCRP cohort, but LDL cholesterol was only a significant risk factor for MACE in the hsCRP cohort. Smoking was a significant risk factor for CBV events in multivariate analysis, but IL-6 and systolic BP were no longer significant risk factors for CBV events. Risk factors for all-cause death, MACE, and CBV events in the IL-6

cohort were very much the same as in the hsCRP cohort, with small differences in the size of the hazard ratios.

Hazard ratios for the different univariate and multivariate risk factors for all-cause death, MACE, and CBV events did not differ substantially for the two cohorts (Tables 3 and 4).

Kaplan–Meier survival curves show associations between all-cause death, MACE, and CBV events and hsCRP values within the hsCRP quartiles (Figures 1A through 1C). Risk of death, MACE, and CBV events increased significantly with increasing hsCRP quartiles not only in the continuous range of hsCRP values but also in the increasing quartile for hsCRP values below 5.0 mg/L (Figures 1D through 1F).

The Kaplan–Meier survival curves by quartiles of IL-6 (Figures 2A through 2C) were comparable to the hsCRP curves in Figures 1A through 1C.

### Discussion

In this *post hoc* analysis, we have shown that the inflammatory markers hsCRP and IL-6, adjusted for traditional risk factors, are predictive for future cardiovascular events and all-cause mortality in renal graft recipients. Although hsCRP and IL-6 are correlated, the observation that both markers were predictive for outcome parameters strengthens the notion that inflammation is an important risk factor for cardiovascular events and all-cause mortality in renal transplant patients.

Our data support previous observational and/or registry data that have indicated that CRP is associated with cardiovascular events and/or all-cause mortality in renal transplant recipients (19–23) and in the general population (5,6,12,13)

In hemodialysis and peritoneal dialysis patients (24–26) and also in the general population, an association between IL-6 and cardiovascular morbidity and mortality has been established (13,27). No trial has examined the association of IL-6 with all-cause mortality or cardiovascular events in renal transplant patients. The data from this analysis demonstrate that the association of IL-6 with cardiovascular events and all-cause mortality also extends to renal transplant patients.

Vascular inflammation is an important factor in the pathogenesis of atherosclerosis in renal transplant recipients (8–11), whereas genetically based elevation in CRP is not associated with posttransplant morbidity and mortality (28), as also shown for the general population (29,30).

In the adult nontransplanted population, the American Heart Association and the Centers for Disease Control and Prevention have endorsed three cutoff points for CRP: <1 mg/dl, low risk; 1 to 3 mg/dl, average risk; and >3 mg/L, high risk (31). By this definition, the overall population in ALERT is in the moderate to high-risk end. The IL-6 data are more difficult to assess across diverse populations because of lack of standardized methods (31). However, in the posttransplant recipients, the inflammatory markers CRP and IL-6 have been reported to be elevated in most studies on renal transplant recipients (19,21–23,32–36), although they are lower than pretransplant values (37).

By dichotomizing the CRP at 0.5 mg/dl, Winkelmayr *et al.* showed that renal transplant patients above that level had a 53% increased mortality risk compared with patients with a

Table 3a. Hazard ratios for all-cause mortality by univariate and multivariate Cox regression in the hsCRP cohort<sup>a</sup>

Parameter	Univariate Hazard Ratio (95% CI) <sup>b</sup>	P Value	Multivariate Hazard Ratio (95% CI)	P Value
Diabetes mellitus	2.12 (1.69 to 2.65)	0.000	2.03 (1.60 to 2.58)	0.000
Smoking current	1.87 (1.43 to 2.46)	0.000	1.98 (1.49 to 2.64)	0.000
CHD	2.49 (1.90 to 3.27)	0.000	1.48 (1.11 to 1.99)	0.008
hsCRP (mg/L)	1.36 (1.21 to 1.52)	0.000	1.15 (1.06 to 1.25)	0.001
Age (yr)	1.07 (1.06 to 1.07)	0.000	1.07 (1.06 to 1.07)	0.000
Creatinine ( $\mu$ mol/L)	1.00 (1.00 to 1.01)	0.000	1.01 (1.00 to 1.01)	0.000
Systolic BP (mmHg)	1.02 (1.01 to 1.02)	0.000	1.00 (0.99 to 1.01)	0.149
Gender	1.08 (1.86 to 1.34)	0.512	1.04 (0.82 to 1.33)	0.740
LDL cholesterol (mmol/L)	1.01 (0.98 to 1.21)	0.980	1.02 (0.91 to 1.13)	0.788

<sup>a</sup>n = 1910.<sup>b</sup>CHD, coronary heart disease; CI, confidence interval.Table 3b. Hazard ratios for MACE by univariate and multivariate Cox regression in the hsCRP cohort<sup>a</sup>

Parameter	Univariate Hazard Ratio (95% CI)	P Value	Multivariate Hazard Ratio (95% CI)	P Value
CHD	3.35 (2.53 to 4.44)	0.000	2.28 (1.69 to 3.08)	0.000
Diabetes mellitus	2.06 (1.59 to 2.64)	0.000	2.08 (1.59 to 2.73)	0.000
LDL cholesterol (mmol/L)	1.38 (1.23 to 1.54)	0.000	1.35 (1.20 to 1.51)	0.000
Smoking current	1.54 (1.11 to 2.14)	0.010	1.43 (1.02 to 2.02)	0.041
hsCRP (mg/L)	1.25 (1.10 to 1.43)	0.001	1.10 (1.01 to 1.20)	0.027
Age (yr)	1.04 (1.03 to 1.05)	0.000	1.03 (1.02 to 1.04)	0.000
Systolic BP (mmHg)	1.01 (1.01 to 1.08)	0.000	1.00 (0.99 to 1.00)	0.560
Creatinine ( $\mu$ mol/L)	1.00 (1.00 to 1.01)	0.000	1.01 (1.00 to 1.00)	0.000
Gender	0.72 (0.55 to 0.93)	0.012	0.86 (0.66 to 1.13)	0.287

<sup>a</sup>n = 1910.Table 3c. Hazard ratios for CBV events by univariate and multivariate Cox regression in the hsCRP cohort<sup>a</sup>

Parameter	Univariate Hazard Ratio (95% CI)	P Value	Multivariate Hazard Ratio (95% CI)	P Value
CHD	2.35 (1.58 to 3.51)	0.000	1.46 (0.96 to 2.22)	0.080
Diabetes mellitus	3.36 (2.49 to 4.54)	0.000	3.17 (2.30 to 4.37)	0.000
LDL cholesterol (mmol/L)	1.10 (0.95 to 1.28)	0.215	1.07 (0.92 to 1.25)	0.401
Smoking current	1.44 (0.97 to 2.14)	0.071	1.61 (1.06 to 2.43)	0.024
hsCRP (mg/L)	1.09 (0.98 to 1.20)	0.109	1.02 (0.92 to 1.14)	0.727
Age (yr)	1.06 (1.04 to 1.07)	0.000	1.06 (1.04 to 1.08)	0.000
Systolic BP (mmHg)	1.02 (1.02 to 1.03)	0.000	1.01 (1.00 to 1.01)	0.059
Creatinine ( $\mu$ mol/L)	1.00 (1.00 to 1.00)	0.314	1.00 (1.00 to 1.01)	0.057
Gender	1.05 (0.90 to 1.23)	0.515	0.49 (0.63 to 1.24)	0.489

<sup>a</sup>n = 1910.

CRP below the threshold (22). In a subsequent paper, Winkel-mayer and colleagues demonstrated a J-shaped association between hsCRP and mortality in kidney transplant recipients when dividing hsCRP values into quartiles for values below 5 mg/L. Patients in the lowest quartile (quartile 1) had an increased mortality risk of about two-fold compared with the chosen reference of quartile 2. There is a linear relationship

between CRP and mortality in the general population (12,38) and in the dialysis population (15,16). No J-shaped curves for mortality have been demonstrated in nonrenal (39) or in dialysis patients (40) for the companion inflammation marker IL-6.

We performed detailed statistical analyses over the whole range of hsCRP values and a separate analysis for hsCRP below 5 mg/L. None of these analyses revealed a J-shaped curve for

Table 4a. Hazard ratios for all-cause death by univariate and multivariate Cox regression in the IL-6 cohort<sup>a</sup>

Parameter	Univariate Hazard Ratio (95% CI)	P Value	Multivariate Hazard Ratio 95% (95% CI)	P Value
Diabetes mellitus	2.11 (1.65 to 2.70)	0.000	1.96 (1.52 to 2.54)	0.000
Smoking current	1.85 (1.38 to 2.45)	0.000	1.96 (1.44 to 2.67)	0.000
CHD	2.56 (1.89 to 3.46)	0.000	1.52 (1.11 to 2.10)	0.010
IL-6 (pg/ml)	1.24 (1.18 to 1.30)	0.000	1.11 (1.05 to 1.18)	0.000
Age (yr)	1.07 (1.05 to 1.08)	0.000	1.07 (1.05 to 1.08)	0.000
Creatinine ( $\mu\text{mol/L}$ )	1.00 (1.00 to 1.01)	0.000	1.01 (1.00 to 1.01)	0.000
Systolic BP (mmHg)	1.02 (1.01 to 1.03)	0.000	1.01 (0.99 to 1.01)	0.080
LDL cholesterol (mmol/L)	1.11 (0.99 to 1.24)	0.075	1.03 (0.92 to 1.16)	0.621
Gender	0.94 (0.75 to 1.20)	0.637	1.07 (0.82 to 1.39)	0.615

<sup>a</sup>*n* = 1751.Table 4b. Hazard ratios for MACE by univariate and multivariate Cox regression in the IL-6 cohort<sup>a</sup>

Parameter	Univariate Hazard Ratio (95% CI)	P Value	Multivariate Hazard Ratio 95% (95% CI)	P Value
Diabetes mellitus	2.15 (1.65 to 2.80)	0.000	2.22 (1.66 to 2.95)	0.000
CHD	3.10 (2.27 to 4.25)	0.000	2.04 (1.45 to 2.85)	0.000
Smoking current	1.55 (1.09 to 2.20)	0.014	1.45 (1.00 to 2.08)	0.048
LDL cholesterol (mmol/L)	1.34 (1.24 to 1.58)	0.000	1.38 (1.22 to 1.55)	0.000
IL-6 (pg/ml)	1.15 (1.08 to 1.22)	0.000	1.08 (1.01 to 1.15)	0.018
Age (yr)	1.04 (1.03 to 1.02)	0.000	1.03 (1.01 to 1.04)	0.000
Creatinine ( $\mu\text{mol/L}$ )	1.00 (1.00 to 1.01)	0.000	1.01 (1.00 to 1.01)	0.000
Systolic BP (mmHg)	1.01 (1.00 to 1.02)	0.000	1.00 (0.99 to 1.01)	0.385
Gender	0.75 (0.57 to 0.99)	0.038	0.94 (0.73 to 1.25)	0.653

<sup>a</sup>*n* = 1751.Table 4c. Hazard ratios for all type of CBV events by univariate and multivariate Cox regression in the IL-6 cohort<sup>a</sup>

Parameter	Univariate Hazard Ratio (95% CI)	P Value	Multivariate Hazard Ratio 95% (95% CI)	P Value
CHD	2.67 (1.77 to 4.04)	0.000	1.65 (1.06 to 2.55)	0.026
Diabetes mellitus	3.34 (2.44 to 4.56)	0.000	3.10 (2.23 to 4.33)	0.000
LDL cholesterol (mmol/L)	1.08 (0.92 to 1.26)	0.345	1.05 (0.89 to 1.23)	0.572
Smoking current	1.48 (0.98 to 2.22)	0.061	1.62 (1.05 to 2.45)	0.028
IL-6 (pg/ml)	1.81 (1.41 to 2.33)	0.000	1.24 (0.94 to 1.63)	0.123
Age (yr)	1.06 (1.04 to 1.07)	0.000	1.06 (1.04 to 1.07)	0.000
Systolic BP (mmHg)	1.02 (1.02 to 1.03)	0.000	1.01 (1.00 to 1.02)	0.075
Creatinine ( $\mu\text{mol/L}$ )	1.00 (1.00 to 1.00)	0.632	1.00 (1.00 to 1.01)	0.288
Gender	0.87 (0.63 to 1.19)	0.369	0.88 (0.62 to 1.24)	0.464

<sup>a</sup>*n* = 1751.

the association of mortality and hsCRP. Repeating these analyses using IL-6 also indicated no J-shaped curve for mortality. The reason for the difference in results is unclear; however, we believe that the data from our trial are more robust because we do have 354 fatal events with a follow-up time of 6.7 yr within the hsCRP cohort.

Inflammatory markers CRP and IL-6 are associated with

diabetes mellitus, insulin resistance, obesity, and adipose tissue (41). Our findings also suggest that higher BMI is associated with increased values of inflammatory markers in transplant patients, but BMI was not a significant risk factor for all-cause death or MACE.

The dialysis population is characterized by a status of high inflammation, and an inverse relationship between cholesterol




Figure 2. Association between (A) all-cause death, (B) major cardiovascular events, and (C) cerebrovascular events with IL-6 for quartiles of IL-6. *x*-axis represents follow-up time in years. *y*-axis represents cumulative survival.

and all-cause mortality has been shown (15,42). However, when corrected for inflammation the usual association for cholesterol and cardiovascular outcome is evident (43). In two recent trials in hemodialysis patients, 4D and Aurora, the inflammation marker hsCRP was significantly associated with increased risk for cardiovascular endpoints (44,45). However, although statin treatment in both trials significantly reduced in LDL cholesterol and hsCRP, the trials failed to show a reduction in the primary cardiovascular endpoints. The findings from these two trials might indicate that inflammation markers in patients receiving hemodialysis differ from other populations, including the renal transplant patients (44,46)

Potential limitations of our study merit consideration. First, this analysis was a *post hoc* analysis of the ALERT extension trial. IL-6 and hsCRP were not available for all patients at baseline, but baseline demographic data of those with or without inflammation analyzed were not different. The strengths of the study are the many patients followed for an extended period and the independent adjudication of predefined outcome parameters.

Renal transplant recipients are at high risk of for premature CVD, which is the leading cause of death in patients with a functional renal graft (3,47). We have previously shown that long-term statin treatment reduces MACE in this population (1,2). However, despite lipid-lowering there is considerable residual risk for cardiovascular events in renal transplant recipients. This might indicate that current management of kidney transplant recipients fails to utilize optimal cardiovascular risk reduction strategies. There is a pressing need to explore the role of other classical and novel cardiovascular risk factors in this population. Adding markers of inflammation such as hsCRP might be helpful to identify high-risk patients.

In conclusion, the inflammatory markers hsCRP and IL-6 may be predictive for future cardiovascular events and all-

cause mortality in renal transplant recipients. These parameters should be considered in the design of future intervention trials.

## Disclosures

Novartis Pharma supported the ALERT core (*Lancet* 36: 2024–2031, 2003) and the extension study (*Am J Transplant* 5: 2929–2936, 2005) by a research grant. This analysis was not supported

Dr. Fellström reports receiving consulting fees from AstraZeneca, Novartis, Roche, and Wyeth; lecture fees from Astellas, Novartis, and Roche; grant support from Novartis, Roche, Merck-Schering-Plough, and Wyeth; and serving as a national coordinator for the Study of Heart and Renal Protection (SHARP) study at Oxford University's Trial Service Unit. Dr. Jardine has received consulting fees from Novartis, AstraZeneca, and Wyeth, and lecture fees from Novartis and Astellas. Dr. Holdaas has received consulting fees from Novartis, AstraZeneca, and Roche; lecture fees from Novartis, AstraZeneca, and Roche; and serves as a regional coordinator for the SHARP study. Ingar Holme reports receiving steering committee honorarium from Merck, Roche, and Pfizer, and consultancy honorariums from AstraZeneca and Merck. Dr. März reports receiving consulting fees from AstraZeneca and Pfizer; lecture fees from Astellas, Astra-Zeneca, Merck-Schering-Plough, Pfizer, and Roche; and grant support from Aventis, Pfizer, Roche, and Siemens. Drs Abedini, Cole, Maes, Weihrach, Neumayer, Ambüh, and Grønhaugen-Riska report no conflict of interest.

## References

1. Holdaas H, Fellstrom B, Jardine AG, Holme I, Nyberg G, Fauchald P, Gronhagen-Riska C, Madsen S, Neumayer HH, Cole E, Maes B, Ambuhl P, Olsson AG, Hartmann A, Solbu DO, Pedersen TR: Effect of fluvastatin on cardiac outcomes in renal transplant recipients: A multicentre, randomised, placebo-controlled trial. *Lancet* 361: 2024–2031, 2003
2. Holdaas H, Fellstrom B, Cole E, Nyberg G, Olsson AG, Pedersen TR, Madsen S, Gronhagen-Riska C, Neumayer HH, Maes B, Ambuhl P, Hartmann A, Staffler B, Jardine AG: Long-term cardiac outcomes in renal transplant recipients receiving fluvastatin: The ALERT extension study. *Am J Transplant* 5: 2929–2936, 2005
3. Kasiske BL, Chakkeri HA, Roel J: Explained and unexplained ischemic heart disease risk after renal transplantation. *J Am Soc Nephrol* 11: 1735–1743, 2000
4. Rigatto C, Parfrey P: Therapy insight: Management of cardiovascular disease in the renal transplant recipient. *Nat Clin Pract Nephrol* 2: 514–526, 2006
5. Kaysen GA: The microinflammatory state in uremia: Causes and potential consequences. *J Am Soc Nephrol* 12: 1549–1557, 2001
6. Willerson JT, Ridker PM: Inflammation as a cardiovascular risk factor. *Circulation* 109: II2–II10, 2004
7. Hansson GK: Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 352: 1685–1695, 2005
8. Vazquez MA, Jeyarajah DR, Kielar ML, Lu CY: Long-term outcomes of renal transplantation: A result of the original endowment of the donor kidney and the inflammatory response to both alloantigens and injury. *Curr Opin Nephrol Hypertens* 9: 643–648, 2000
9. Cottone S, Palermo A, Vaccaro F, Mule G, Guarneri M, Arsena R, Vadala A, Cerasola G: Inflammation and endo-

- thelial activation are linked to renal function in long-term kidney transplantation. *Transpl Int* 20: 82–87, 2007
10. Gotsman I, Grabie N, Gupta R, Dacosta R, MacConmara M, Lederer J, Sukhova G, Witztum JL, Sharpe AH, Lichtman AH: Impaired regulatory T-cell response and enhanced atherosclerosis in the absence of inducible costimulatory molecule. *Circulation* 114: 2047–2055, 2006
  11. Martinez-Castelao A, Sarrias X, Bestard O, Gil-Vernet S, Seron D, Cruzado JM, Moreso F, Diez-Noguera A, Grinyo JM: Arterial elasticity measurement in renal transplant patients under anticalcineurin immunosuppression. *Transplant Proc* 37: 3788–3790, 2005
  12. Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH: Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 336: 973–979, 1997
  13. Pai JK, Pischon T, Ma J, Manson JE, Hankinson SE, Joshipura K, Curhan GC, Rifai N, Cannuscio CC, Stampfer MJ, Rimm EB: Inflammatory markers and the risk of coronary heart disease in men and women. *N Engl J Med* 351: 2599–2610, 2004
  14. Ridker PM, Danielson E, Fonseca FA, Genest J, Gotto AM Jr, Kastelein JJ, Koenig W, Libby P, Lorenzatti AJ, MacFadyen JG, Nordestgaard BG, Shepherd J, Willerson JT, Glynn RJ: Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med* 359: 2195–2207, 2008
  15. Stenvinkel P, Heimbürger O, Paulter F, Diczfalusy U, Wang T, Berglund L, Jogestrand T: Strong association between malnutrition, inflammation, and atherosclerosis in chronic renal failure. *Kidney Int* 55: 1899–1911, 1999
  16. Wanner C, Zimmermann J, Schwedler S, Metzger T: Inflammation and cardiovascular risk in dialysis patients. *Kidney Int* 61: S99–S102, 2002
  17. Lauzurica R, Pastor MC, Bayes B, Hernandez JM, Bonet J, Dolade M, Navarro M, Romero R: Pretransplant inflammation: A risk factor for delayed graft function? *J Nephrol* 21: 221–228, 2008
  18. Mange KC, Joffe MM, Feldman HI: Effect of the use or nonuse of long-term dialysis on the subsequent survival of renal transplants from living donors. *N Engl J Med* 344: 726–731, 2001
  19. Bakri RS, Afzali B, Covic A, Sriskantharan R, Bharmariza P, Park WH, Sriharan M, Dalton N, Wierzbicki AS, Crook MA, Goldsmith DJ: Cardiovascular disease in renal allograft recipients is associated with elevated sialic acid or markers of inflammation. *Clin Transplant* 18: 201–204, 2004
  20. Varaganam M, Finney H, Trevitt R, Sharples E, McCloskey DJ, Sinnott PJ, Raftery MJ, Yaqoob MM: Pretransplantation levels of C-reactive protein predict all-cause and cardiovascular mortality, but not graft outcome, in kidney transplant recipients. *Am J Kidney Dis* 43: 502–507, 2004
  21. Ducloux D, Kazory A, Chalopin JM: Predicting coronary heart disease in renal transplant recipients: A prospective study. *Kidney Int* 66: 441–447, 2004
  22. Winkelmayer WC, Lorenz M, Kramar R, Fodinger M, Horl WH, Sunder-Plassmann G: C-reactive protein and body mass index independently predict mortality in kidney transplant recipients. *Am J Transplant* 4: 1148–1154, 2004
  23. Winkelmayer WC, Schaeffner ES, Chandraker A, Kramar R, Rumpold H, Sunder-Plassmann G, Fodinger M: A J-shaped association between high-sensitivity C-reactive protein and mortality in kidney transplant recipients. *Transpl Int* 20: 505–511, 2007
  24. Noh H, Lee SW, Kang SW, Shin SK, Choi KH, Lee HY, Han DS: Serum C-reactive protein: A predictor of mortality in continuous ambulatory peritoneal dialysis patients. *Perit Dial Int* 18: 387–394, 1998
  25. Stenvinkel P, Barany P, Heimbürger O, Pecoits-Filho R, Lindholm B: Mortality, malnutrition, and atherosclerosis in ESRD: What is the role of interleukin-6? *Kidney Int* 61: S103–S108, 2002
  26. Zimmermann J, Herrlinger S, Pruy A, Metzger T, Wanner C: Inflammation enhances cardiovascular risk and mortality in hemodialysis patients. *Kidney Int* 55: 648–658, 1999
  27. Ridker PM, Rifai N, Rose L, Buring JE, Cook NR: Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med* 347: 1557–1565, 2002
  28. Kruger B, Boger CA, Schroppe B, Farkas S, Schnitzbauer AA, Hoffmann U, Obed A, Murphy BT, Banas B, Kramer BK: No effect of C-reactive protein (CRP) haplotypes on CRP levels and post-transplant morbidity and mortality in renal transplantation. *Transpl Int* 21: 452–458, 2008
  29. Zacho J, Tybjaerg-Hansen A, Jensen JS, Grande P, Silleesen H, Nordestgaard BG: Genetically Elevated C-reactive protein and ischemic vascular disease. *N Engl J Med* 359: 1897–1908, 2008
  30. Timpson NJ, Lawlor DA, Harbord RM, Gaunt TR, Day IN, Palmer LJ, Hattersley AT, Ebrahim S, Lowe GD, Rumley A, Davey SG: C-reactive protein and its role in metabolic syndrome: Mendelian randomisation study. *Lancet* 366: 1954–1959, 2005
  31. Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO, III, Criqui M, Fadl YY, Fortmann SP, Hong Y, Myers GL, Rifai N, Smith SC Jr, Taubert K, Tracy RP, Vinicor F: Markers of inflammation and cardiovascular disease: Application to clinical and public health practice: A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 107: 499–511, 2003
  32. Kusec V, Smalcelj R, Puretic Z, Szekeres T: Interleukin-6, transforming growth factor-beta 1, and bone markers after kidney transplantation. *Calcif Tissue Int* 75: 1–6, 2004
  33. Ioannidou E, Kao D, Chang N, Burleson J, Dongari-Bagtzoglou A: Elevated serum interleukin-6 (IL-6) in solid-organ transplant recipients is positively associated with tissue destruction and IL-6 gene expression in the periodontium. *J Periodontol* 77: 1871–1878, 2006
  34. Sezer S, Akcay A, Ozdemir FN, Kulah E, Arat Z, Haberal M: Post-transplant C-reactive protein monitoring can predict chronic allograft nephropathy. *Clin Transplant* 18: 722–725, 2004
  35. van Ree RM, Oterdoom LH, de Vries AP, Gansevoort RT, van der Heide JJ, van Son WJ, Ploeg RJ, de Jong PE, Gans RO, Bakker SJ: Elevated levels of C-reactive protein independently predict accelerated deterioration of graft function in renal transplant recipients. *Nephrol Dial Transplant* 22: 246–253, 2007
  36. Cueto-Manzano AM, Morales-Buenrostro LE, Gonzalez-Espinoza L, Gonzalez-Tableros N, Martin-del-Campo F, Correa-Rotter R, Valera I, Alberu J: Markers of inflammation before and after renal transplantation. *Transplantation* 80: 47–51, 2005



37. Simmons EM, Langone A, Sezer MT, Vella JP, Recupero P, Morrow JD, Ikizler TA, Himmelfarb J: Effect of renal transplantation on biomarkers of inflammation and oxidative stress in end-stage renal disease patients. *Transplantation* 79: 914–919, 2005
38. Ridker PM, Cook N: Clinical usefulness of very high and very low levels of C-reactive protein across the full range of Framingham risk scores. *Circulation* 109: 1955–1959, 2004
39. Harris TB, Ferrucci L, Tracy RP, Corti MC, Wacholder S, Ettinger WH Jr, Heimovitz H, Cohen HJ, Wallace R: Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. *Am J Med* 106: 506–512, 1999
40. Bologa RM, Levine DM, Parker TS, Cheigh JS, Serur D, Stenzel KH, Rubin AL: Interleukin-6 predicts hypoalbuminemia, hypocholesterolemia, and mortality in hemodialysis patients. *Am J Kidney Dis* 32: 107–114, 1998
41. Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW: C-reactive protein in healthy subjects: Associations with obesity, insulin resistance, and endothelial dysfunction: A potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 19: 972–978, 1999
42. Lowrie EG, Lew NL: Commonly measured laboratory variables in hemodialysis patients: Relationships among them and to death risk. *Semin Nephrol* 12: 276–283, 1992
43. Liu Y, Coresh J, Eustace JA, Longenecker JC, Jaar B, Fink NE, Tracy RP, Powe NR, Klag MJ: Association between cholesterol level and mortality in dialysis patients: Role of inflammation and malnutrition. *JAMA* 291: 451–459, 2004
44. Fellstrom BC, Jardine AG, Schmieder RE, Holdaas H, Bannister K, Beutler J, Chae DW, Chevaile A, Cobbe SM, Gronhagen-Riska C, De Lima JJ, Lins R, Mayer G, McMahon AW, Parving HH, Remuzzi G, Samuelsson O, Sonkodi S, Sci D, Suleymanlar G, Tsakiris D, Tesar V, Todorov V, Wiecek A, Wuthrich RP, Gottlow M, Johnsson E, Zannad F: Rosuvastatin and cardiovascular events in patients undergoing hemodialysis. *N Engl J Med* 360: 1395–1407, 2009
45. Wanner C, Krane V, Marz W, Olschewski M, Mann JF, Ruf G, Ritz E: Atorvastatin in patients with type 2 diabetes mellitus undergoing hemodialysis. *N Engl J Med* 353: 238–248, 2005
46. Krane V, Winkler K, Drechsler C, Lilienthal J, Marz W, Wanner C: Effect of atorvastatin on inflammation and outcome in patients with type 2 diabetes mellitus on hemodialysis. *Kidney Int* 74: 1461–1467, 2008
47. Lindholm A, Albrechtsen D, Frodin L, Tufveson G, Persson NH, Lundgren G: Ischemic heart disease—Major cause of death and graft loss after renal transplantation in Scandinavia. *Transplantation* 60: 451–457, 1995

**Access to UpToDate on-line is available for additional clinical information  
at <http://www.cjasn.org/>**