

Physical Activity in ESRD: Time to Get Moving

Stephen L. Seliger

Clin J Am Soc Nephrol 7: 1927–1929, 2012. doi: 10.2215/CJN.11041012

Studies over the last 3 decades among ESRD patients on maintenance dialysis have demonstrated greatly impaired physical performance and cardiorespiratory fitness compared with the general population (1–4). Levels of habitual physical activity are also markedly lower in these patients. For example, Johansen *et al.* reported that among 34 maintenance hemodialysis patients, directly measured physical activity was nearly a third lower compared with healthy “sedentary” controls (5). Using data from the US Renal Data System (USRDS), Stack *et al.* reported that 56% of incident dialysis patients did not engage in any exercise, even once weekly (6). Accurate measurement of physical activity can be challenging in both ESRD patients and the general population. In large samples of participants, the most feasible technique is the use of questionnaires that ask about a range of potential physical activities that individuals may have engaged in within a fixed time period. “Gold-standard” measurement of physical activity relies on the use of a monitoring device worn at the waist or the leg that detects acceleration and records the duration and intensity of movement. The correlation between the self-reported questionnaires and directly measured activity varies widely but is often modest, including in renal disease patients (7).

In this issue of *CJASN*, Mastsuzawa and colleagues performed a single-center prospective observational cohort study among 202 maintenance hemodialysis patients (8). Physical activity on 4 sequential nondialysis days was measured directly using uniaxial accelerometry. The authors defined “habitual physical activity” as the average daily activity consistent with at least “gentle walking,” corresponding to at least an intensity of 1 on a 0–9 semi-quantitative scale. On the basis of prior studies in healthy adults cited by the authors, this corresponds to at least approximately 1.9 times the baseline energy expenditure occurring at rest (9). The authors considered a cut-point of at least 50 minutes average daily activity on nondialysis days as indicating a high level of activity, based on prior research suggesting that activity above this threshold is associated with maintenance of physical function in hemodialysis patients. Among the 202 participants, 37% met this threshold of activity. As expected, those hemodialysis patients who were more active were younger and had lower burden of overall comorbidity but surprisingly were not significantly less likely to have diabetes or heart disease. Over 7 years of follow-up, a total of 34 deaths occurred, 19 attributed to

cardiovascular disease. The estimated cumulative survival was markedly different between the two groups, at 93.3% in the high-activity group versus 77.2% in the low-activity group. After adjustment for comorbidity, each 10-min/d increment in activity was associated with a 12% lower risk of all-cause mortality, a difference that was highly statistically significant.

It is perhaps not surprising that physical activity should predict longevity and mortality in ESRD. A number of prospective cohort studies have identified physical activity as a strong predictor of mortality after accounting for demographics and comorbid conditions (10). These studies have generally been conducted among the general population, and the few studies conducted among renal disease patients relied on self-reported rather than directly measured activity. Increasing physical activity has a number of well described beneficial physiologic effects on cardiorespiratory fitness, endothelial function (11), BP (12), dyslipidemia (13), muscle strength, and inflammation (14), among others. These effects offer plausible biologic pathways to explain the relationship of greater physical activity with lower risk of mortality. The major contribution of the study by Mastsuzawa *et al.* is to demonstrate that the apparent survival advantage of those with higher levels of physical activity extends to ESRD patients on maintenance dialysis, as determined by objective measures of physical activity.

The results of this study should be placed in appropriate context. At first glance, it might seem surprising to nephrologists caring for maintenance hemodialysis patients in the United States that nearly 40% of such patients engage in high levels of physical activity, as was the case in the Japanese center in this study. Comparison of the overall levels of activity in this study with the results of other ESRD studies is made challenging by the different methods used to quantify activity. A potential frame of reference might be the consensus guidelines from the US Centers for Disease Control and Prevention (CDC), which suggest that adults engage in at least 150 minutes of “moderate to vigorous” physical activity per week (15). With mean daily activity duration of 42 minutes on each of the 5 nondialysis days in this study, one might expect the average weekly activity to well surpass these CDC recommendations. However, it is unclear to what extent the activity performed by the ESRD patients in this study would constitute moderate to vigorous activity.

Division of Nephrology, University of Maryland School of Medicine, Baltimore, Maryland; and Department of Medicine, Maryland Veterans Affairs Healthcare System, Baltimore, Maryland

Correspondence:

Dr. Stephen L. Seliger, Division of Nephrology, University of Maryland School of Medicine, 22 S. Greene Street, N3W143, Baltimore, MD 21201. Email: sseliger@medicine.umaryland.edu

It seems unlikely that the gentle walking (a score of 1 on the 0–9 semi-quantitative scale) included in the authors' definition of habitual activity would coincide with the intensity recommended by the CDC. In non-ESRD populations, the intensity of physical activity has prognostic significance independent of duration and total energy expenditure (10,16). This study unfortunately does not offer information on the important question of how greater duration and/or greater intensity of physical activity relate to mortality and morbidity risk in ESRD.

The major limitation of this study, however, is the problem of unmeasured confounding, a limitation common to observational studies of physical activity. It is uncertain whether physical activity itself or factors such as comorbidity, which determine both physical activity and mortality risk, are to account for the observed associations in this study. The authors attempt to adjust for comorbidity through the use of a previously devised ESRD-specific comorbidity index, which has been shown to have independent prognostic benefit in predicting all-cause mortality (17). However, this score was derived using administrative USRDS data rather than comorbidity reported by patients or verified through medical records, and has not been validated in non-US ESRD populations. Even if the comorbidity score was accurate in reflecting the presence of chronic medical conditions, it is highly likely that residual confounding would remain, due to differences in the severity and/or chronicity of these conditions (18). For example, although the presence of congestive heart failure may be appropriately captured in this score, differences between patients in systolic function and dyspnea severity would not be measured. Such differences could plausibly affect both physical activity and risk for mortality. Ultimately, it is not possible to know with certainty from these results whether low-activity patients are incapable of being more active—because of comorbidity such as cardiac disease, pulmonary disease, neurologic impairment, orthopedic conditions, or a host of other factors—or whether the level of physical activity is primarily determined by patients' lifestyle and is therefore modifiable.

Even with these limitations, the study raises intriguing questions regarding the potential of increasing physical activity in ESRD patients as an intervention to reduce mortality and morbidity. Over the past 3 decades, a number of interventional studies have examined the clinical and physiologic effects of increased physical activity through structured exercise training interventions in maintenance dialysis patients. As summarized in a recent Cochrane meta-analysis of 32 reported studies (19), exercise interventions resulted in improvements in aerobic capacity, walking capacity, BP, and health-related quality of life. Unfortunately, most of the studies included in the meta-analysis were individually quite small (<30 patients) and suffered from a high risk of bias due to methodologic limitations. Furthermore, the studies varied widely in their approach to exercise training, including aerobic exercise, resistance exercise, yoga, intradialytic exercise, and home-based training. Even if the observed benefits in physiologic measures and quality of life are reliable, it is unclear whether these benefits translate to reduced mortality, which was not examined as an outcome in any of the prior studies. The recent results of the Look AHEAD (Action for Health in Diabetes) lifestyle intervention study in obese patients with type 2 diabetes

suggest that these physiologic measures may not be reliable surrogate outcomes; the diet and physical activity intervention in that study resulted in favorable changes in cardiorespiratory fitness, BP, dyslipidemia, and glycemic control (20), but no difference in risk of cardiovascular morbidity and mortality (21).

Undoubtedly, there would be considerable challenges to the design and conduct of a large, adequately powered, interventional study of increasing physical activity in ESRD. As demonstrated by prior smaller single-center studies, there are many different approaches to exercise that could be considered in this population. For example, resistance versus aerobic exercise training, in-center versus at-home exercise, and intradialysis versus nondialysis exercise are among a few of many possible designs of exercise regimens. Difficulties with retention and treatment adherence are common among lifestyle intervention studies, and choice of the control intervention can influence the observed results. On the other hand, one should consider that a variety of pharmacologic interventions to reduce morbidity and mortality in ESRD—including statins (22), noncalcium phosphate binders (23), and anemia correction (24), among others—have already been evaluated in large multicenter trials with largely underwhelming results. These studies were motivated in part based on earlier observational studies suggesting associations of biologic risk factors with morbidity and mortality in ESRD. Perhaps it is time to consider inactivity and impaired physical function as “risk factors” for morbidity and mortality in this population, and to design appropriately powered investigations of increasing physical activity.

Disclosures

None.

References

- Johansen KL, Chertow GM, da Silva M, Carey S, Painter P: Determinants of physical performance in ambulatory patients on hemodialysis. *Kidney Int* 60: 1586–1591, 2001
- Sterky E, Stegmayr BG: Elderly patients on haemodialysis have 50% less functional capacity than gender- and age-matched healthy subjects. *Scand J Urol Nephrol* 39: 423–430, 2005
- Goldberg AP, Geltman EM, Gavin JR 3rd, Carney RM, Hagberg JM, Delmez JA, Naumovich A, Oldfield MH, Harter HR: Exercise training reduces coronary risk and effectively rehabilitates hemodialysis patients. *Nephron* 42: 311–316, 1986
- Painter P, Messer-Rehak D, Hanson P, Zimmerman SW, Glass NR: Exercise capacity in hemodialysis, CAPD, and renal transplant patients. *Nephron* 42: 47–51, 1986
- Johansen KL, Chertow GM, Ng AV, Mulligan K, Carey S, Schoenfeld PY, Kent-Braun JA: Physical activity levels in patients on hemodialysis and healthy sedentary controls. *Kidney Int* 57: 2564–2570, 2000
- Stack AG, Molony DA, Rives T, Tyson J, Murthy BV: Association of physical activity with mortality in the US dialysis population. *Am J Kidney Dis* 45: 690–701, 2005
- Robinson-Cohen C, Littman AJ, Duncan GE, Roshanravan B, Izkizler TA, Himmelfarb J, Kestenbaum BR: Assessment of physical activity in chronic kidney disease [published online ahead of print June 26, 2012]. *J Ren Nutr* doi:10.1053/j.jrn.2012.04.008
- Matsuzawa R, Matsunaga A, Wang G, Kutsuna T, Ishii A, Abe Y, Takagi Y, Yoshida A, Takahira N: Habitual physical activity measured by accelerometer and survival in maintenance hemodialysis patients. *Clin J Am Soc Nephrol* 7: 2010–2016, 2012

9. Kumahara H, Schutz Y, Ayabe M, Yoshioka M, Yoshitake Y, Shindo M, Ishii K, Tanaka H: The use of uniaxial accelerometry for the assessment of physical-activity-related energy expenditure: A validation study against whole-body indirect calorimetry. *Br J Nutr* 91: 235–243, 2004
10. Lee IM, Hsieh CC, Paffenbarger RS Jr: Exercise intensity and longevity in men. The Harvard Alumni Health Study. *JAMA* 273: 1179–1184, 1995
11. McDermott MM, Ades P, Guralnik JM, Dyer A, Ferrucci L, Liu K, Nelson M, Lloyd-Jones D, Van Horn L, Garside D, Kibbe M, Domanchuk K, Stein JH, Liao Y, Tao H, Green D, Pearce WH, Schneider JR, McPherson D, Laing ST, McCarthy WJ, Shroff A, Criqui MH: Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: A randomized controlled trial. *JAMA* 301: 165–174, 2009
12. Kelley GA, Kelley KA, Tran ZV: Aerobic exercise and resting blood pressure: A meta-analytic review of randomized, controlled trials. *Prev Cardiol* 4: 73–80, 2001
13. Goldberg AP, Hagberg JM, Delmez JA, Haynes ME, Harter HR: Metabolic effects of exercise training in hemodialysis patients. *Kidney Int* 18: 754–761, 1980
14. Nicklas BJ, Hsu FC, Brinkley TJ, Church T, Goodpaster BH, Kritchevsky SB, Pahor M: Exercise training and plasma C-reactive protein and interleukin-6 in elderly people. *J Am Geriatr Soc* 56: 2045–2052, 2008
15. Physical Activity Guidelines Advisory Committee: Physical Activity Guidelines Advisory Committee Report, 2008. US Department of Health and Human Services, Washington, DC, 2008
16. Laursen AH, Kristiansen OP, Marott JL, Schnohr P, Prescott E: Intensity versus duration of physical activity: Implications for the metabolic syndrome. A prospective cohort study. *BMJ Open* 2: pii: e001711, 2012
17. Liu J, Huang Z, Gilbertson DT, Foley RN, Collins AJ: An improved comorbidity index for outcome analyses among dialysis patients. *Kidney Int* 77: 141–151, 2010
18. Seliger SL: Comorbidity and confounding in end-stage renal disease. *Kidney Int* 77: 83–85, 2010
19. Heiwe S, Jacobson SH: Exercise training for adults with chronic kidney disease. *Cochrane Database Syst Rev* (10): CD003236, 2011
20. Wing RR; Look AHEAD Research Group: Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: Four-year results of the Look AHEAD trial. *Arch Intern Med* 170: 1566–1575, 2010
21. National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health: Weight loss does not lower heart disease risk from type 2 diabetes. Available at: <http://www.nih.gov/news/health/oct2012/niddk-19.htm>. Accessed October 30, 2012
22. Wanner C, Krane V, März W, Olschewski M, Mann JF, Ruf G, Ritz E; German Diabetes and Dialysis Study Investigators: Atorvastatin in patients with type 2 diabetes mellitus undergoing hemodialysis. *N Engl J Med* 353: 238–248, 2005
23. Suki WN, Zabaneh R, Cangiano JL, Reed J, Fischer D, Garrett L, Ling BN, Chasan-Taber S, Dillon MA, Blair AT, Burke SK: Effects of sevelamer and calcium-based phosphate binders on mortality in hemodialysis patients. *Kidney Int* 72: 1130–1137, 2007
24. Besarab A, Bolton WK, Browne JK, Egrie JC, Nissenson AR, Okamoto DM, Schwab SJ, Goodkin DA: The effects of normal as compared with low hematocrit values in patients with cardiac disease who are receiving hemodialysis and epoetin. *N Engl J Med* 339: 584–590, 1998

Published online ahead of print. Publication date available at www.cjasn.org.

See related article, “Habitual Physical Activity Measured by Accelerometer and Survival in Maintenance Hemodialysis Patients” on pages 2010–2016.