

Dialysis and Mortality: Does It Matter Where You Live?

Rudolph A. Rodriguez

Clin J Am Soc Nephrol 7: 1055–1057, 2012. doi: 10.2215/CJN.05410512

A 70-year-old man with diabetic nephropathy and a history of multiple admissions for congestive heart failure is approaching the need for renal replacement therapy. He is struggling with his decision regarding the optimal dialysis modality. He lives alone in rural Washington, 60 miles from the nearest dialysis unit. He is a man of modest means and cannot afford the transportation costs associated with thrice weekly in-center hemodialysis. Given his less than optimal prior compliance with his complex medical regimen, he is a risky candidate for peritoneal or home hemodialysis. Due to transportation issues, he decides that peritoneal dialysis is his only viable option. Does geography impact on patient survival? Clinicians usually do not consider a patient's residence when making clinical decisions, but epidemiologic evidence has shown that patient characteristics, health services, and patient outcomes vary significantly according to rural versus urban residence. These geographic differences also exist within urban areas depending on residential characteristics. Rural America represents approximately 20% of the US population and is comprised of diverse cultures, landscapes, and economic and social characteristics. In 2005 the Institute of Medicine report, *Quality through Collaboration: The Future of Rural Health Care*, was released with the goal of providing an independent assessment of health and health care in rural America and formulating an action plan for quality-focused rural community health systems (1). The report highlighted a number of important issues in rural America including reduced access to quality health services, poor access to quality emergency services, and adverse patient characteristics such as higher rates of obesity, suicide, alcohol and tobacco use, poverty, and inferior treatment for acute myocardial infarction. These factors are of importance to rural patients with CKD and could represent major barriers to quality renal and general medical care.

In this issue of the *CJASN*, Maripuri *et al.* (2) present a retrospective cohort study showing that rural residence is associated with higher mortality in those rural patients receiving peritoneal dialysis. This study is a nice extension to our previous work, which had not stratified survival results by dialysis modality and therefore had not shown an increase in mortality in rural patients on peritoneal dialysis. Our study did hint at a problem by showing that facilities in rural

areas were less likely to offer peritoneal dialysis than those units in urban areas (3). Other studies in the United States and Canada have also shown an association between poor patient outcomes and rural residence (4,5).

When reading this study, it is important to note the difficulty in comparing survival between peritoneal and hemodialysis patients. Not only do the therapies differ tremendously, but the patients themselves are quite different, with peritoneal dialysis patients tending to be younger with less comorbidities. The findings of numerous studies comparing survival between the modalities have not been consistent (6). In an attempt to better match patients by important baseline characteristics, a recent study compared survival between the modalities by using a matched-pair cohort with the matching based on propensity of initial peritoneal dialysis (7). The study found that survival was modestly longer for patients initiating renal replacement therapy with peritoneal dialysis compared with matched hemodialysis patients. However, similar to other studies, they found this modality survival effect is modified by age and the presence of diabetes. In general, peritoneal dialysis patients >65 years of age and those with diabetes have a higher risk of death than those on hemodialysis. Adding to the complexity of comparing dialysis modalities, the rural population tends to be older with more comorbidities, and this is true in this study by Maripuri *et al.*, in which the rural and micropolitan population is older and all medical comorbid conditions with the exception for alcohol and drug use are more prevalent in the non-urban population. Therefore, despite controlling for the baseline characteristics, the results of the study could be confounded by the unmeasured differences and/or the differences in severity of the comorbid conditions in the rural population. Despite this possibility, the results of the study are important and should not be ignored by clinicians. The results of this study are consistent with other studies and call attention to the fact that many rural patients are high-risk candidates for peritoneal dialysis on the basis of age and comorbid conditions.

It is also important to note that there is no accepted definition of rural and that the level of isolation varies dramatically among the rural population in the United States. As clinicians, we worry about patients who live

University of Washington, Seattle and VA Puget Sound Healthcare System, Seattle, Washington

Correspondence: Dr. Rudolph A. Rodriguez, Puget Sound VA Medical Center, Nephrology and Renal Dialysis Unit, Building 100, Room 5B113, 1660 S. Columbian Way, Seattle, WA 98108. Email: rudyrod@u.washington.edu

require a multidisciplinary team approach involving local primary care physicians and local medical facilities, which are likely to be unfamiliar with the management of peritoneal dialysis patients and the transplant referral process.

Returning to the case, this gentleman said he would rather die than move from his home located in an idyllic rural area. He accepted the risks of peritoneal dialysis and is currently thriving on the therapy. However, one day he will likely start to fail on peritoneal dialysis and he will be faced with some difficult decisions. From a medical standpoint, he may be best served by relocating closer to a hemodialysis unit, but moving from his community would leave him isolated from his friends and support structure and possibly drive him into poverty. A frank discussion would explore relocation along with other options including optimizing his peritoneal dialysis prescription, providing more peritoneal dialysis support, switching to home hemodialysis, and if appropriate, kidney transplantation. Geography does matter, and the renal community should start adapting to the needs of the rural patients requiring renal replacement therapy.

Disclosures

None.

References

1. Institute of Medicine (U.S.). Committee on the Future of Rural Health Care: *Quality Through Collaboration: The Future of Rural Health*, Washington, DC, National Academies Press, 2005
2. Maripuri S, Arbogast P, Ikizler TA, Cavanaugh KL: Rural and micropolitan residence and mortality in patients on dialysis. *Clin J Am Soc Nephrol* 7: 1121–1129, 2012
3. O'Hare AM, Johansen KL, Rodriguez RA: Dialysis and kidney transplantation among patients living in rural areas of the United States. *Kidney Int* 69: 343–349, 2006
4. Mehrotra R, et al: Neighborhood location, rurality, geography, and outcomes of peritoneal dialysis patients in the United States. *Perit Dial Int* 32: 322–331, 2011
5. Tonelli M, Hemmelgarn B, Culleton B, Klarenbach S, Gill JS, Wiebe N, Manns B; Alberta Kidney Disease Network: Mortality of Canadians treated by peritoneal dialysis in remote locations. *Kidney Int* 72: 1023–1028, 2007
6. Vonesh EF, Snyder JJ, Foley RN, Collins AJ: Mortality studies comparing peritoneal dialysis and hemodialysis: what do they tell us? *Kidney Int Suppl* 103: S3–S11, 2006
7. Weinhandl ED, Foley RN, Gilbertson DT, Arneson TJ, Snyder JJ, Collins AJ: Propensity-matched mortality comparison of incident hemodialysis and peritoneal dialysis patients. *J Am Soc Nephrol* 21: 499–506, 2010
8. National Center for Frontier Communities: National Center for Frontier Communities. Available at <http://www.frontierus.org/2000update.php>. Accessed May 28, 2012
9. Northwest Renal Network: Annual report. Available at <http://www.nwrenalnetwork.org/S/dfr10/dfr3.htm>. Accessed May 28, 2012
10. Network 8, Inc.: Annual report. Available at <http://www.esrdnetwork8.org/dialysis-transplant-providers/annual-reports-data-tables/Assets/2010-AR-data-tables.pdf>. Accessed May 28, 2012
11. Jiang Z, Yu X: Advancing the use and quality of peritoneal dialysis by developing a peritoneal dialysis satellite center program. *Perit Dial Int* 31: 121–126, 2011
12. Nakamoto H: Telemedicine system for patients on continuous ambulatory peritoneal dialysis. *Perit Dial Int* 27[Suppl 2]: S21–S26, 2007
13. Tonelli M, Klarenbach S, Rose C, Wiebe N, Gill J: Access to kidney transplantation among remote- and rural-dwelling patients with kidney failure in the United States. *JAMA* 301: 1681–1690, 2009

Published online ahead of print. Publication date available at www.cjasn.org.

See related article, “Rural and Micropolitan Residence and Mortality in Patients on Dialysis,” on pages 1121–1129.