

# CJASN

Clinical Journal of the American Society of Nephrology

August 2012 • Vol. 7 • No. 8

## Editorials

---

### 1201 Hyperkalemia in Predialysis Patients

*Biff F. Palmer*

*See related article on page 1234.*

### 1203 Is Long-Term Prediction in Membranous Nephropathy (MGN) Better Than the Weatherman's Forecast Capacity?

*Daniel C. Cattran and Joseph Kim*

*See related article on page 1242.*

### 1206 Getting Excited about Exit Sites in Peritoneal Dialysis?

*Peter G. Blake*

*See related article on page 1266.*

### 1209 Sequestration and Suppression of Anti-HLA Antibodies by a Failed Kidney Allograft

*Vinay Nair and Peter S. Heeger*

*See related article on page 1310.*

## Original Articles

---

### Acid/Base and Electrolyte Disorders

#### 1211 Does the Exposure of Urine Samples to Air Affect Diagnostic Tests for Urine Acidification?

*Joo-Hark Yi, Hyun-Jong Shin, Sun-Moon Kim, Sang-Woong Han, Ho-Jung Kim, and Man-Seok Oh*

### Acute Kidney Injury /Acute Renal Failure

#### 1217 FGF-23 Levels in Patients with AKI and Risk of Adverse Outcomes

*David E. Leaf, Myles Wolf, Sushrut S. Waikar, Herbert Chase, Marta Christov, Serge Cremers, and Leonard Stern*

#### 1224 Association between Peritransplant Kidney Injury Biomarkers and 1-Year Allograft Outcomes

*Isaac E. Hall, Mona D. Doshi, Peter P. Reese, Richard J. Marcus, Heather Thiessen-Philbrook, and Chirag R. Parikh*

### Clinical Nephrology

#### 1234 Prevalence and Factors Associated with Hyperkalemia in Predialysis Patients Followed in a Low-Clearance Clinic

*Pantelis A. Sarafidis, Rochelle Blacklock, Eleri Wood, Adam Rumjon, Shanique Simmonds, Jessica Fletcher-Rogers, Rachel Ariyanayagam, Aziza Al-Yassin, Claire Sharpe, and Katie Vinen*  
*See related editorial on page 1201.*

#### 1242 Prognostic Value of Risk Score and Urinary Markers in Idiopathic Membranous Nephropathy

*Jan A.J.G. van den Brand, Julia M. Hofstra, and Jack F.M. Wetzels*  
*See related editorial on page 1203.*

### Clinical Pharmacology

#### 1249 Pharmacokinetics of Intraperitoneal Gentamicin in Peritoneal Dialysis Patients with Peritonitis (GIPD Study)

*Julie M. Varghese, Jason A. Roberts, Steven C. Wallis, Robert J. Boots, Helen Healy, Robert G. Fassett, Jeffrey Lipman, and Dwarakanathan Ranganathan*

ESRD and Chronic Dialysis

**1257 Decoy Receptor 3, a Novel Inflammatory Marker, and Mortality in Hemodialysis Patients**

*Szu-Chun Hung, Ta-Wei Hsu, Yao-Ping Lin, and Der-Cherng Tarng*

**1266 The Association between Exit Site Infection and Subsequent Peritonitis among Peritoneal Dialysis Patients**

*Anouk T.N. van Diepen, George A. Tomlinson, and Sarbjit V. Jassal*  
See related editorial on page 1206.

**1272 Serum Potassium and Cause-Specific Mortality in a Large Peritoneal Dialysis Cohort**

*Klara Torlén, Kamyar Kalantar-Zadeh, Miklos Z. Molnar, Tania Vashistha, and Rajnish Mehrotra*

**1285 Troponin T for the Detection of Dialysis-Induced Myocardial Stunning in Hemodialysis Patients**

*Tobias Breidthardt, James O. Burton, Aghogho Odudu, Mohamed Tarek Eldehni, Helen J. Jefferies, and Christopher W. McIntyre*

Genetics

**1293 Association of HLA Genes with Clinical Outcomes of ANCA-Associated Vasculitis**

*Dong-yuan Chang, Huan Luo, Xu-jie Zhou, Min Chen, and Ming-Hui Zhao*

Hypertension

**1300 Probing the Mechanisms of Intradialytic Hypertension: A Pilot Study Targeting Endothelial Cell Dysfunction**

*Jula K. Inrig, Peter Van Buren, Catherine Kim, Wanpen Vongpatanasin, Thomas J. Povsic, and Robert Toto*

Renal Transplantation

**1310 Donor-Specific Antibodies after Ceasing Immunosuppressive Therapy, with or without an Allograft Nephrectomy**

*Arnaud Del Bello, Nicolas Congy-Jolivet, Federico Sallusto, Celine Guilbeau-Frugier, Isabelle Cardeau-Desangles, Marylise Fort, Laure Esposito, Joelle Guitard, Olivier Cointault, Laurence Lavayssière, Marie Béatrice Nogier, Antoine Blancher, Lionel Rostaing, and Nassim Kamar*  
See related editorial on page 1209.

**1320 Incidence and Outcomes of BK Virus Allograft Nephropathy among ABO- and HLA-Incompatible Kidney Transplant Recipients**

*Adnan Sharif, Nada Alachkar, Serena Bagnasco, Duvuru Geetha, Gaurav Gupta, Karl Womer, Lois Arend, Lorraine Racusen, Robert Montgomery, and Edward Kraus*

---

In-Depth Review

**1328 Renal Replacement Therapy in Critically Ill Patients Receiving Extracorporeal Membrane Oxygenation**

*David J. Askenazi, David T. Selewski, Matthew L. Paden, David S. Cooper, Brian C. Bridges, Michael Zappitelli, and Geoffrey M. Fleming*

---

Mini-Review

**1337 Transthyretin Amyloidosis and the Kidney**

*Luísa Lobato and Ana Rocha*

---

Special Features

**1347 Nephrology Quiz and Questionnaire: Renal Replacement Therapy**

*Thomas A. Golper (Discussant), Richard J. Glassock (Co-Moderator), and Anthony J. Bleyer (Co-Moderator)*

**1353 A Tribute to George E. Schreiner**

*Robert J. Rubin*  
*Paul D. Doolan*

## Special Features (Continued)

---

### **1355 New Metrics for Assessing Diagnostic Potential of Candidate Biomarkers**

*John W. Pickering and Zoltan H. Endre*

### **1365 A Position Paper on Standardizing the Nonneoplastic Kidney Biopsy Report**

*Anthony Chang, Ian W. Gibson, Arthur H. Cohen, Jan J. Weening, J. Charles Jennette, and Agnes B. Fogo*

## Errata

---

**1369 Correction**

**1370 Correction**

**1372 Correction**

### **On the Cover**

*What's the diagnosis?* Nodular glomerulopathy is a lesion classically seen in patients with diabetic renal disease. Similar pattern of glomerular injury may also be seen in heavy smokers and patients with light and/or heavy chain deposition disease. The mechanism of injury in these circumstances is probably similar, where glucose, cigarette smoke components, or paraproteins cause toxic injury of mesangial cells, and they in turn overproduce the mesangial matrix with formation of nodules. Long-standing membranoproliferative glomerulonephritis (MPGN) can also result in prominent mesangial expansion and nodular glomerulopathy; PAS or silver stains usually show well-developed double contours of the peripheral capillary loops in such cases. Healed mesangiolytic lesions in the context of thrombotic microangiopathy can also result in segmental nodular glomerulosclerosis. Immunofluorescence and electron microscopy studies, as well as detailed clinical history, are essential in differentiating these conditions and diseases. (Image and text provided by Vanesa Bijol, MD, Brigham and Women's Hospital)