

# Ionizing Radiation During Pretransplant Evaluation: Time to Reconsider the Evaluation Process

Deborah B. Adey

*Clin J Am Soc Nephrol* 8: 711–713, 2013. doi: 10.2215/CJN.03120313

Patients with CKD presenting for evaluation for renal transplant often undergo an extensive pretransplant assessment. The purpose of the evaluation is to detect underlying medical conditions that pose potential perioperative risk to the patient. Although kidney transplant prolongs life expectancy compared with remaining on dialysis, it is an elective surgery. As such, centers often leave no stone unturned when it comes to assessment of risk. Cardiovascular evaluation of potential renal transplant candidates is one of the most controversial topics dominating selection committees. The leading cause of death after transplant is cardiovascular. Diabetes and hypertension are the leading causes of kidney disease, both of which are significant risk factors for cardiovascular disease. In addition, CKD is felt to be an independent risk factor for cardiovascular disease.

In this issue of *CJASN*, Nguyen *et al.* (1) report on their retrospective analysis of radiation exposure to potential renal transplant recipients undergoing evaluation at a single transplant center. Their study included only imaging examinations performed for evaluation for transplant and therefore may underreport the full radiation exposure to patients. Imaging studies ordered by primary care physicians, nephrologists, or other medical subspecialists were not captured. Nuclear medicine studies were the predominant source of radiation exposure, 89.2% of which were nuclear medicine cardiac stress tests. Nearly 30% of patients were exposed to radiation doses considered high (>50–100 mSv) or very high (>100 mSv). An average nuclear medicine perfusion study results in exposure to 41 mSv, whereas coronary arteriography is associated with exposure of 2–30 mSv (2), depending on such factors as procedural complexity, imaging equipment, and operator experience. Routine posteroanterior chest radiography provides an estimated 0.02 mSv of radiation exposure compared with 2–16 mSv for computed tomography, 7 mSv for cardiac catheterization, and 15 mSv for percutaneous intervention (3,4). One must take pause when the average background exposure is 3 mSv per year and the recommended maximum occupational exposure is 20 mSv per year, or 100 mSv over 5 years.

In Nguyen and colleagues' study, factors associated with more exposure to radiation included black race, with an odds ratio (OR) of 3.39; age >40 years, with an

OR of 6.97; and diabetes, with an OR of 3.35. In a study by De Mauri and associates, this same pretransplant population was at increased risk for excessive exposure to ionizing radiation during pretransplant evaluation, although in the form of computed tomography (5). The risk factors associated with increased exposure to radiation reflect the population at greater risk for cardiovascular disease.

The real question to address is whether the studies being performed during pretransplant evaluation are providing the information needed to make a decision about the need for intervention. Imaging to ascertain the presence of underlying malignancy is reasonable and not under question. An occult malignancy could become aggressive and more difficult to treat with exposure to immunosuppression. However, the evaluation for asymptomatic coronary artery disease must be reconsidered. Nuclear medicine myocardial perfusion studies are routinely performed to assess ischemic heart disease. The reported sensitivity and specificity of nuclear medicine stress testing for detecting ischemia are 85% and 91%, respectively (6). The ironic point is that the clinical scenario in which nuclear medical myocardial perfusion studies are weakest involves balanced ischemia, where there is no differential perfusion of the myocardium due to diffuse disease.

The transplant community needs to take a step back and seriously consider what the right study is for the given situation. What is the goal when imaging studies are ordered? The DIAD study (7) randomly assigned 1123 patients with type 2 diabetes and no symptoms of ischemia to nuclear medicine myocardial perfusion screening versus no screening and found no difference in outcomes. The Coronary Artery Prophylaxis (CARP) study found no benefit to cardiac revascularization in the absence of three-vessel disease or left main disease (8). The CARP investigators randomly assigned 510 patients to revascularization versus no revascularization before elective major vascular surgery. Three-vessel coronary artery disease, left main disease, or unstable angina warrants revascularization. A study by Friedman *et al.* compared the disparate guidelines put forth by the Kidney Disease Outcomes Quality Initiative and by the American Society of Transplantation, the Lisbon guidelines, and the guidelines of the American College of Cardiology as they apply to renal transplant cardiac risk assessment (9). The

Kidney and Pancreas Transplant Service, University of California-Davis, Sacramento, California

**Correspondence:**  
Dr. Deborah B. Adey, Transplant Nephrology, 2233 Stockton Boulevard, Room 2015, University of California-Davis, Sacramento, CA 95817. Email: deborah.adey@ucdmc.ucdavis.edu

American College of Cardiology guidelines may underestimate cardiovascular risk in this population, whereas the American Society of Transplantation guidelines may subject patients to unnecessary testing.

Options for noninvasive cardiac evaluation without ionizing radiation should be considered. Echocardiographic assessment of ejection fraction correlated well with post-transplant outcomes. de Mattos and colleagues (10) demonstrated a correlation with outcomes. Patients with an ejection fraction of <40% had a worse prognosis after transplant, even in the absence of ischemia. There was a graded reverse-stepped correlation between ejection fraction and mortality. This finding supports the value of noninvasive studies without the use of ionizing radiation. Dobutamine stress echocardiography has been reported to have the best sensitivity and specificity among noninvasive studies in assessing underlying coronary artery disease (11). Positive results on dobutamine stress echocardiography demonstrated an 88% sensitivity and 94% specificity for predicting severe coronary artery disease in renal transplant recipients, values similar if not superior to those seen with nuclear medicine myocardial perfusion studies.

Perioperative cardiovascular risk may be adequately assessed from noninvasive evaluation of a patient's physical functioning status. A recent study by Rosas *et al.* (12) demonstrated that assessment of physical functioning correlates with post-transplant cardiovascular events. A Physical Activity Scale for the Elderly (PASE) questionnaire was used to assess physical activity within 2 weeks of transplantation. Although the PASE was designed for the elderly, this questionnaire has been validated in the chronically ill and non-elderly, the sedentary, and patients with CKD. Lower scores for physical activity were associated with all-cause mortality, as well as death with a functioning graft. These data are consistent with data from other studies that demonstrate a relationship between physical functioning and outcomes after transplant (13,14). However, there is no consensus on what constitutes adequate physical functioning. The 6-minute walk test, sit-stand studies, ability to climb stairs, and maximal oxygen consumption with ambulation have been used to assess physical functioning in patients with CKD. Those with low physical functioning may benefit from an intervention, such as referral to physical therapy or cardiac rehabilitation program.

The concern regarding exposure to ionizing radiation is the potential to increase the risk of malignancy. Patients presenting for evaluation for transplant typically need to wait a specified amount of time if they have had a malignancy before transplant, to assure there is no evidence of recurrence. The risk of malignancy is certainly increased in the renal transplant population; it is estimated to be 2.5–5 times higher than that in the general population. Immunosuppression impairs immune surveillance and increases malignancy risk. The amount of radiation exposure during the pretransplant evaluation may be contributing to post-transplant malignancy risk. In 2010, the U.S. Food and Drug Administration issued an initiative to reduce unnecessary radiation exposure from medical imaging (15). This was borne from the recognition that radiation exposure among the United States population has nearly doubled over two decades. The initiative recommends that the healthcare community develop criteria for use of procedures that

expose patients to ionizing radiation in the form of computed tomography, nuclear medicine studies, or fluoroscopy.

In conclusion, the study by Nguyen *et al.* brings attention to one of the most controversial topics in transplantation: How should patients be evaluated for cardiovascular risk? The authors conclude that a less invasive approach to the evaluation of the transplant patient be considered. This echoes the "Call to Action" paper by Friedman and colleagues (9), which highlights the need to reconsider the pretransplant cardiac evaluation of transplant candidates. The transplant community might consider the use of a combination of assessments that involve evaluation of physical functioning and dobutamine stress echocardiography to predict patients at higher perioperative cardiovascular risk. Those patients deemed at high cardiovascular risk on the basis of risk factors, physical functioning, and dobutamine stress echocardiography could then be further evaluated with coronary angiography to assess underlying coronary artery disease. Such an approach may limit unnecessary exposure to ionizing radiation in a population already at an increased risk of malignancy after transplant.

#### Disclosures

D.B.A. reports no disclosures directly related to this editorial topic. She is a coinvestigator on a Genentech-sponsored study of desensitization.

#### References

1. Nguyen KN, Patel AM, Weng FL: Ionizing radiation exposure among kidney transplant recipients due to medical imaging during the pretransplant evaluation. *Clin J Am Soc Nephrol* 8: 833–839, 2013
2. Einstein AJ, Moser KW, Thompson RC, Cerqueira MD, Henzlova MJ: Radiation dose to patients from cardiac diagnostic imaging. *Circulation* 116: 1290–1305, 2007
3. Chen J, Einstein AJ, Fazel R, Krumholz HM, Wang Y, Ross JS, Ting HH, Shah ND, Nasir K, Nallamothu BK: Cumulative exposure to ionizing radiation from diagnostic and therapeutic cardiac imaging procedures: A population-based analysis. *J Am Coll Cardiol* 56: 702–711, 2010
4. Gerber TC, Carr JJ, Arai AE, Dixon RL, Ferrari VA, Gomes AS, Heller GV, McCollough CH, McNitt-Gray MF, Mettler FA, Mieres JH, Morin RL, Yester MV: Ionizing radiation in cardiac imaging: a science advisory from the American Heart Association Committee on Cardiac Imaging of the Council on Clinical Cardiology and Committee on Cardiovascular Imaging and Intervention of the Council on Cardiovascular Radiology and Intervention. *Circulation* 119: 1056–1065, 2009
5. De Mauri A, Brambilla M, Izzo C, Matheoud R, Chiarinotti D, Carriero A, Stratta P, De Leo M: Cumulative radiation dose from medical imaging in kidney transplant patients. *Nephrol Dial Transplant* 27: 3645–3651, 2012
6. O'Rourke RA, Brundage BH, Froelicher VF, Greenland P, Grundy SM, Hachamovitch R, Pohost GM, Shaw LJ, Weintraub WS, Winters WL Jr: American College of Cardiology/American Heart Association Expert Consensus Document on electron-beam computed tomography for the diagnosis and prognosis of coronary artery disease. *J Am Coll Cardiol* 36: 326–340, 2000
7. Young LH, Wackers FJ, Chyun DA, Davey JA, Barrett EJ, Taillefer R, Heller GV, Iskandrian AE, Wittlin SD, Filipchuk N, Ratner RE, Inzucchi SE: DIAD Investigators: Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: A randomized controlled trial. *JAMA* 301: 1547–1555, 2009
8. McFall EO, Ward HB, Moritz TE, Goldman S, Krupski WC, Littooy F, Pierpont G, Santilli S, Rapp J, Hattler B, Shunk K, Jaenicke C, Thottapurathu L, Ellis N, Reda DJ, Henderson WG: Coronary-artery revascularization before elective major vascular surgery. *N Engl J Med* 351: 2795–2804, 2004

9. Friedman SE, Palac RT, Zlotnick DM, Chobanian MC, Costa SP: A call to action: Variability in guidelines for cardiac evaluation before renal transplantation. *Clin J Am Soc Nephrol* 6: 1185–1191, 2011
10. de Mattos AM, Siedlecki A, Gaston RS, Perry GJ, Julian BA, Kew CE, Deierhoi MH, Young C, Curtis JJ, Iskandrian AE: Systolic dysfunction portends increased mortality among those waiting for renal transplant. *J Am Soc Nephrol* 19: 1191–1196, 2008
11. Sharma R, Pellerin D, Gaze DC, Gregson H, Streather CP, Collinson PO, Brecker SJ: Dobutamine stress echocardiography and the resting but not exercise electrocardiograph predict severe coronary artery disease in renal transplant candidates. *Nephrol Dial Transplant* 20: 2207–2214, 2005
12. Rosas SE, Reese PP, Huan Y, Doria C, Cochetti PT, Doyle A: Pretransplant physical activity predicts all-cause mortality in kidney transplant recipients. *Am J Nephrol* 35: 17–23, 2012
13. Zelle DM, Corpeleijn E, Stolk RP, de Greef MHG, Gans ROB, van der Heide JJH, Navis G, Bakker SJ: Low physical activity and risk of cardiovascular and all-cause mortality in renal transplant recipients. *Clin J Am Soc Nephrol* 6: 898–905, 2011
14. Kutner NG, Zhang R, Bowles T, Painter P: Pretransplant physical functioning and kidney patients' risk for posttransplantation hospitalization/death: Evidence from a national cohort. *Clin J Am Soc Nephrol* 1: 837–843, 2006
15. U.S. Food and Drug Administration: Initiative to reduce unnecessary radiation exposure from medical imaging. 2010. Available at: <http://www.fda.gov/Radiation-EmittingProducts/RadiationSafety/RadiationDoseReduction/ucm199994.htm/>. Accessed February 22, 2013

Published online ahead of print. Publication date available at [www.cjasn.org](http://www.cjasn.org).

See related article, "Ionizing Radiation Exposure among Kidney Transplant Recipients Due to Medical Imaging during the Pretransplant Evaluation," on pages 833–839.