

# Progressive Inflammation and Wasting in Patients with ESRD

George A. Kayesen

*Clin J Am Soc Nephrol* 9: 225–226, 2014. doi: 10.2215/CJN.12541213

Body mass and, more importantly, body cell mass decline in patients with increasing time on RRT (1–3). Malnutrition, defined by low levels of albumin, low levels of prealbumin, low body mass index, and reduced physical function, is predictive of mortality (4). Although albumin concentration may increase in the first year after hemodialysis initiation, as a consequence of either nutritional improvement or the reduction in proteinuria that accompanies loss of residual renal function (5), after that year, the trajectory for albumin and prealbumin is a decrease with vintage (1,3). Albumin concentration as well as creatinine, which is a surrogate for muscle mass, are determined by several factors, but inflammation and nutrition are major determinants (6), with inflammation having greater predictive power (7). Importantly, inflammation is also strongly associated with mortality, and therefore, recognition, prevention, or reversal of it has potential to improve outcomes.

Dialysis patients are inflamed compared with the healthy population, which is assessed by C-reactive protein (CRP) or IL-6. The median value of CRP in most dialysis populations is considerably above the upper tertile in the general population (8). There is an incremental increase in inflammation among patients with CKD stage 5 after initiation of dialysis (9). The average already increased levels of acute phase proteins found in dialysis patients are interrupted by episodes of increased inflammation, which are accompanied by decreases in negative acute phase proteins albumin, prealbumin, and transferrin (5,7,10) (also nutritional surrogate biomarkers) and decreases in creatinine (7), suggesting decline in muscle mass (11). It is unclear whether effects on muscle mass reverse when acute inflammatory insults remit. Over the short term, however, albumin values return to baseline after inflammatory events (5,7), which does not explain the downward trajectory for both postdialysis weight and serum albumin concentration that was observed (3). Decreases in nutritional intake over time have been documented (3), providing one potential target for therapeutic intervention and one mechanism for nutritional decline. den Hoedt *et al.* (12) report a longitudinal decline in albumin similar to the decline reported in the Hemodialysis (HEMO) study (3), but in contrast to previous reports, den Hoedt *et al.* (12) describe a time-dependent increase in inflammation in patients in

both treatment arms of the Convective Transport Study (CONTRAST). The greater than 80% prevalence of arteriovenous fistulas at baseline was substantially greater than in the HEMO study (27%), and CRP levels were significantly less than in the HEMO study as well as incident and prevalent hemodialysis patients in the United States (5,7). Serum albumin concentration reflects both inflammation, which is measured as CRP, and nutrition, which is estimated as normalized protein catabolic rate (7) and health-related quality of life (13). However, the dominant driver for albumin concentration as well as creatinine, which is associated with muscle mass (7), is inflammation. Muscle mass has been observed to decrease with years on dialysis (14).

It is rational to associate inflammatory processes with a dialysis population acquiring vascular access other than arteriovenous fistulas. It is unclear within the CONTRAST population whether vascular access changed significantly over time. If there was a significant increase in the prevalence of arteriovenous fistulas, it would offer a potential explanation for the increased inflammatory status described. Nevertheless, the basis for increased inflammation in CKD and dialysis patients overall is not well understood. Alterations in enzyme function, oxidative stress, lipoprotein structure, and function may all play a role, and how these processes progress over time as well as the underlying mechanisms for their derangement may also affect alterations. Improving outcomes in dialysis patients will require more in-depth understanding of these processes. Although it is useful to target vascular access as a driver of inflammation in dialysis patients, underlying processes, such as alterations in the microbiome (15,16), gut permeability to endotoxin (17), and gene expression, also occur with aging (also associated with loss of muscle mass and increased prevalence of inflammation) and in CKD by mechanisms, such as the reduction in nuclear factor erythroid 2–related factor 2 (18,19), and increase in myostatin expression (20). The relative abundance of patients having arteriovenous fistulas likely provides the relatively low level of inflammation in this population, allowing for appreciation that inflammation is a progressive process in dialysis patients. Nutritional deterioration accompanied by increased inflammation over time may indicate intermediate outcomes that require resolution as well as act as canaries in the coal mine, marking processes that increase disability

Department of Medicine, Division of Nephrology and Department of Biochemistry and Molecular Medicine, University of California, Davis, California

#### Correspondence:

Dr. George A. Kayesen, Department of Medicine, Division of Nephrology and Department of Biochemistry and Molecular Medicine, University of California, One Shields Avenue, GBSF 451 East Health Sciences Drive, Room 6301, Davis, CA 95616. Email: gakaysen@ucdavis.edu

and the risk of death in dialysis patients. Recognition that inflammation, like the nutritional variables with which it is associated, progressively changes over time offers a potential area for basic scientific exploration to clarify underlying mechanisms and serves as a potential target for therapeutic intervention.

#### Disclosures

G.A.K. is a consultant for the Renal Research Institute and has been a speaker for Merck. G.A.K. is also a member of a data safety monitoring board providing oversight for the National Institutes of Health on clinical trials exploring treatment of inflammation, alteration in the gut microbiome, and other interventions.

#### References

- Chertow GM, Johansen KL, Lew N, Lazarus JM, Lowrie EG: Vintage, nutritional status, and survival in hemodialysis patients. *Kidney Int* 57: 1176–1181, 2000
- Nakao T, Kanazawa Y, Nagaoka Y, Iwasawa H, Uchinaga A, Matsumoto H, Okada T, Yoshino M: Body protein index based on bioelectrical impedance analysis is a useful new marker assessing nutritional status: Applications to patients with chronic renal failure on maintenance dialysis. *Contrib Nephrol* 155: 18–28, 2007
- Rocco MV, Dwyer JT, Larive B, Greene T, Cockram DB, Chumlea WC, Kusek JW, Leung J, Burrowes JD, McLeroy SL, Poole D, Uhlin L; HEMO Study Group: The effect of dialysis dose and membrane flux on nutritional parameters in hemodialysis patients: Results of the HEMO Study. *Kidney Int* 65: 2321–2334, 2004
- Vashistha T, Mehrotra R, Park J, Streja E, Dukkipati R, Nissenson AR, Ma JZ, Kovesdy CP, Kalantar-Zadeh K: Effect of age and dialysis vintage on obesity paradox in long-term hemodialysis patients [published online ahead of print October 9, 2013]. *Am J Kidney Dis* doi:10.1053/j.ajkd.2013.07.021
- Dalrymple LS, Johansen KL, Chertow GM, Grimes B, Anand S, McCulloch CE, Kaysen GA: Longitudinal measures of serum albumin and prealbumin concentrations in incident dialysis patients: The comprehensive dialysis study. *J Ren Nutr* 23: 91–97, 2013
- Kaysen GA, Greene T, Daugirdas JT, Kimmel PL, Schulman GW, Toto RD, Levin NW, Yan G; HEMO Study Group: Longitudinal and cross-sectional effects of C-reactive protein, equilibrated normalized protein catabolic rate, and serum bicarbonate on creatinine and albumin levels in dialysis patients. *Am J Kidney Dis* 42: 1200–1211, 2003
- Kaysen GA, Chertow GM, Adhikarla R, Young B, Ronco C, Levin NW: Inflammation and dietary protein intake exert competing effects on serum albumin and creatinine in hemodialysis patients. *Kidney Int* 60: 333–340, 2001
- Ridker PM, Buring JE, Shih J, Matias M, Hennekens CH: Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. *Circulation* 98: 731–733, 1998
- McIntyre CW, Selby NM, Sigrist M, Pearce LE, Mercer TH, Naish PF: Patients receiving maintenance dialysis have more severe functionally significant skeletal muscle wasting than patients with dialysis-independent chronic kidney disease. *Nephrol Dial Transplant* 21: 2210–2216, 2006
- Kaysen GA, Dubin JA, Müller HG, Rosales LM, Levin NW; The HEMO Study Group: The acute-phase response varies with time and predicts serum albumin levels in hemodialysis patients. *Kidney Int* 58: 346–352, 2000
- Workeneh BT, Mitch WE: Review of muscle wasting associated with chronic kidney disease. *Am J Clin Nutr* 91: 1128S–1132S, 2010
- den Hoedt CH, Bots ML, Grooteman MPC, van der Weerd NC, Penne EL, Mazairac AHA, Levesque R, Blankestijn PJ, Nubé MJ, ter Wee PM, van den Dorpel MA; CONTRAST Investigators: Clinical predictors of decline in nutritional parameters over time in ESRD. *Clin J Am Soc Nephrol* 9: 318–325, 2014
- Mazairac AH, de Wit GA, Penne EL, van der Weerd NC, Grooteman MP, van den Dorpel MA, Nubé MJ, Buskens E, Lévesque R, Ter Wee PM, Bots ML, Blankestijn PJ; CONTRAST Investigators: Protein-energy nutritional status and kidney disease-specific quality of life in hemodialysis patients. *J Ren Nutr* 21: 376–386, 2011
- Kaynar K, Songul Tat T, Ulusoy S, Cansiz M, Ozkan G, Gul S, Bektas O: Evaluation of nutritional parameters of hemodialysis patients. *Hippokratia* 16: 236–240, 2012
- Poesen R, Meijers B, Evenepoel P: The colon: An overlooked site for therapeutics in dialysis patients. *Semin Dial* 26: 323–332, 2013
- Wang IK, Lai HC, Yu CJ, Liang CC, Chang CT, Kuo HL, Yang YF, Lin CC, Lin HH, Liu YL, Chang YC, Wu YY, Chen CH, Li CY, Chuang FR, Huang CC, Lin CH, Lin HC: Real-time PCR analysis of the intestinal microbiotas in peritoneal dialysis patients. *Appl Environ Microbiol* 78: 1107–1112, 2012
- McIntyre CW, Harrison LE, Eldehni MT, Jefferies HJ, Szeto CC, John SG, Sigrist MK, Burton JO, Hothi D, Korshheed S, Owen PJ, Lai KB, Li PK: Circulating endotoxemia: A novel factor in systemic inflammation and cardiovascular disease in chronic kidney disease. *Clin J Am Soc Nephrol* 6: 133–141, 2011
- Kim HJ, Vaziri ND: Contribution of impaired Nrf2-Keap1 pathway to oxidative stress and inflammation in chronic renal failure. *Am J Physiol Renal Physiol* 298: F662–F671, 2010
- El Assar M, Angulo J, Rodríguez-Mañas L: Oxidative stress and vascular inflammation in aging. *Free Radic Biol Med* 65: 380–401, 2013
- Zhang L, Rajan V, Lin E, Hu Z, Han HQ, Zhou X, Song Y, Min H, Wang X, Du J, Mitch WE: Pharmacological inhibition of myostatin suppresses systemic inflammation and muscle atrophy in mice with chronic kidney disease. *FASEB J* 25: 1653–1663, 2011

Published online ahead of print. Publication date available at [www.cjasn.org](http://www.cjasn.org).

See related article, “Clinical Predictors of Decline in Nutritional Parameters over Time in ESRD,” on pages 318–325.