

Factors Influencing Treatment of Atypical Hemolytic Uremic Syndrome

Carla M. Nester and Richard J.H. Smith

Clin J Am Soc Nephrol 9: 1516–1518, 2014. doi: 10.2215/CJN.07540714

On the Verge of a Revolution

Over the past 2 decades, tremendous strides have been made in our understanding of the ultra-rare disease atypical hemolytic uremic syndrome (aHUS). At the center of this progress has been the discovery that the majority of patients with aHUS (>50%) carry causative DNA variants in alternate complement pathway (AP) genes (1–4). In another approximately 10% of patients, the disease is driven by antibodies to factor H, the major regulator of the AP, triggering acquired complement-mediated hemolytic uremic syndrome (HUS) (2,5,6). The clinical consequence of these discoveries has been the successful implementation of focused efforts to block the terminal complement cascade as a practical approach to targeted therapy for patients with aHUS. With the adoption of targeted therapy has come the recognition of new challenges. In this editorial, we address factors that may influence treatment of aHUS, a number of which are highlighted by Chinchilla *et al.* in this issue of *CJASN* (7).

Chinchilla *et al.* present a comprehensive genetic assessment of a cohort of patients with aHUS (83 of 275) aged <2 years and negative for autoimmune-mediated HUS (7). The authors aimed to determine the contribution of diacylglycerol kinase ϵ (DGK ϵ) mutations to aHUS in this Spanish population. DGK ϵ is the most recent gene causally related to aHUS (8–10). The novelty of this discovery resides in the presumed mechanism by which DGK ϵ causes aHUS. As an intracellular protein without an overt association with the complement cascade, DGK ϵ is hypothesized to cause aHUS by promoting a prothrombotic state, which has implications for treatment.

An Imprecise Nomenclature

At the heart of defining factors that influence treatment of aHUS is the nonspecific nature of the term aHUS. Created as a foil to “typical” HUS (HUS associated with shiga-toxin carrying bacteria), the adjective “atypical” was never meant to imply a cause. At best, this term encompasses a group of diseases that share in common the clinical features of a microangiopathic hemolytic anemia associated with thrombocytopenia and renal failure. At its worst, however, aHUS lumps together a group of diseases with very different underlying pathologies. In practice, there is little agreement

on what defines or limits classifying someone as an aHUS patient.

The disparate definitions of aHUS are illustrated by the reports of two of the larger aHUS cohorts. Fremeaux-Bacchi *et al.* use the term aHUS to designate HUS after all secondary forms had been excluded (1). By contrast, Noris *et al.* adopted a broader definition (2) and included all patients with HUS, excluding only those with thrombotic thrombocytopenic purpura or those associated with shiga toxin-carrying bacteria. In this issue of *CJASN*, Chinchilla *et al.* define their aHUS cohort as patients with microangiopathic hemolytic anemia with thrombocytopenia and renal failure in whom autoimmunity has been excluded (7). The finding that 5% of their aHUS cohort carries recessive mutations in DGK ϵ is considerably lower than the 27% reported in the French cohort (1). Given the differences in inclusion criteria, it remains possible that these results reflect enrollment criteria and not ethnic-specific genetic differences.

Strict definitions of aHUS based on a mechanistic understanding of disease are essential for well designed clinical trials. An optimal therapeutic approach should target the primary pathology and if the primary pathology is appropriately targeted, a disease response should be seen. At the simplest level, if the term aHUS is applied too broadly, it becomes difficult to determine whether an apparent lack of response relates to an ineffective treatment or an inappropriate treatment. Similarly, lumping different pathologies together may hide treatment successes.

Matching Pathology to Treatment

An example within the aHUS group demonstrates how unique, pathology-related factors influence the treatment of aHUS in these relative settings. As mentioned previously, at the genetic level, a *CFH* mutation may be transcribed into a nonfunctional protein leading to aHUS. However, loss of factor H function can also be the consequence of autoantibodies to this protein. Although the phenotypic consequence is dysregulation of the alternate pathway of complement and aHUS, different targeted treatments are possible because the pathophysiology is fundamentally different. Certainly, anticomplement therapy appears to work for both types of aHUS. However, in patients with autoimmune-mediated HUS, a more direct therapy

Molecular Otolaryngology and Renal Research Laboratories, Division of Nephrology, Department of Internal Medicine, and Stead Family Department of Pediatrics, Carver College of Medicine, University of Iowa, Iowa City, Iowa

Correspondence:

Dr. Carla M. Nester, Rare Renal Disease Clinic, University of Iowa, 200 Hawkins Drive, 4036BT, Iowa City, IA, 52242. Email: carla-nester@uiowa.edu

would target the antibody and not the terminal complement pathway. This tenet is the basis for considering anticellular immune suppression to reduce antibody titer and limit risk for exacerbation without compromising complement activity in this select patient cohort.

The study by Chinchilla *et al.* highlights other factors that may influence the treatment of aHUS (7). In their cohort, Chinchilla *et al.* identified four patients that carried mutations in DGK ϵ . Of particular interest is that three of these four patients also carried complement mutations, the significance of which is 3-fold. First, it mandates screening of the complement genes in patients with DGK ϵ mutations. Second, their findings suggest that segregating both complement mutations and DGK ϵ mutations alters the disease phenotype. Third, response to anticomplement therapy is affected by mutational analyses.

Lemaire *et al.* first reported DGK ϵ -mediated HUS, noting that this type of HUS did not respond to plasma therapy or eculizumab, a reasonable expectation given the presumption that DGK ϵ mediates HUS through a different mechanism (8). Importantly, neither of their patients had complement mutations. By contrast, Chinchilla *et al.* observed a clinical response with either plasma therapy or eculizumab in the three patients carrying both DGK ϵ mutations and complement mutations (7). The overriding message is that response to therapy depends on the mechanism of disease. As such, a complete genetic workup may influence treatment choices.

Treatment When the Pathophysiology Is Uncertain

Although a comprehensive understanding of the primary pathophysiology underlying aHUS is critical to informing treatment, it is also possible that by defining secondary pathology, alternative therapeutic targets may be identified. Over 30% of patients with aHUS do not have an identifiable mutation in the complement genes currently screened in most clinical laboratories. Yet many of these patients are treated empirically and appear to respond to either plasma exchange or eculizumab. Multiple research centers are studying these cases to identify disease drivers that include pathways that interact with the complement cascade (1–6,8,10–39). It is now understood, for example, that there is crosstalk between the complement system and the coagulation pathway, but the precise relationship between the two pathways and how that relationship could affect a disease like aHUS is unclear. In the absence of a more complete pathologic understanding, as an interim step, it may be reasonable to use biomarkers as indices of disease activity. For example, most patients with aHUS do not have a low C3. Perhaps a low C3 during disease exacerbations, even in the absence of explanatory genetic or acquired factors, should be enough to label the process as “complement associated” and initiate anticomplement therapy.

Chinchilla *et al.* teach us an important lesson: A comprehensive understanding of the genetic and acquired drivers of aHUS is necessary to understand the predisposing pathology and to guide appropriate diagnosis and effective treatment. Patients respond to therapies that are appropriate to the cause of their disease. Until we define the underlying pathology, treatment regimens remain empirical. As currently used, the term aHUS includes diseases of

differing causes. It remains a major task to identify the additional pathologic drivers in this group of diseases and to refine our treatment protocols on that basis. Perhaps the generic term aHUS has outlived its usefulness and a more informative nomenclature should be adopted?

Disclosures

C.M.N. has received fees from Alexion for Advisory Board Activity. Both R.J.H.S. and C.M.N. have received grant support from Celldex.

References

1. Fremeaux-Bacchi V, Fakhouri F, Garnier A, Bienaimé F, Dragon-Durey MA, Ngo S, Moulin B, Servais A, Provot F, Rostaing L, Burtsey S, Niaudet P, Deschênes G, Lebranchu Y, Zuber J, Loirat C: Genetics and outcome of atypical hemolytic uremic syndrome: A nationwide French series comparing children and adults. *Clin J Am Soc Nephrol* 8: 554–562, 2013
2. Noris M, Caprioli J, Bresin E, Mossali C, Pianetti G, Gamba S, Daina E, Fenili C, Castelletti F, Sorosina A, Piras R, Donadelli R, Maranta R, van der Meer I, Conway EM, Zipfel PF, Goodship TH, Remuzzi G: Relative role of genetic complement abnormalities in sporadic and familial aHUS and their impact on clinical phenotype. *Clin J Am Soc Nephrol* 5: 1844–1859, 2010
3. Bresin E, Rurali E, Caprioli J, Sanchez-Corral P, Fremeaux-Bacchi V, Rodriguez de Cordoba S, Pinto S, Goodship TH, Alberti M, Ribes D, Valoti E, Remuzzi G, Noris M; European Working Party on Complement Genetics in Renal Diseases: Combined complement gene mutations in atypical hemolytic uremic syndrome influence clinical phenotype. *J Am Soc Nephrol* 24: 475–486, 2013
4. Maga TK, Nishimura CJ, Weaver AE, Frees KL, Smith RJ: Mutations in alternative pathway complement proteins in American patients with atypical hemolytic uremic syndrome. *Hum Mutat* 31: E1445–E1460, 2010
5. Hofer J, Janecke AR, Zimmerhackl LB, Riedl M, Rosales A, Giner T, Cortina G, Haindl CJ, Petzelberger B, Pawlik M, Jeller V, Vester U, Gadner B, van Husen M, Moritz ML, Würzner R, Jungraithmayr T; German-Austrian HUS Study Group: Complement factor H-related protein 1 deficiency and factor H antibodies in pediatric patients with atypical hemolytic uremic syndrome. *Clin J Am Soc Nephrol* 8: 407–415, 2013
6. Sinha A, Gulati A, Saini S, Blanc C, Gupta A, Gurjar BS, Saini H, Kotresh ST, Ali U, Bhatia D, Ohri A, Kumar M, Agarwal I, Gulati S, Anand K, Vijayakumar M, Sinha R, Sethi S, Salmona M, George A, Bal V, Singh G, Dinda AK, Hari P, Rath S, Dragon-Durey MA, Bagga A; Indian HUS Registry: Prompt plasma exchanges and immunosuppressive treatment improves the outcomes of anti-factor H autoantibody-associated hemolytic uremic syndrome in children. *Kidney Int* 85: 1151–1160, 2014
7. Chinchilla DS, Pinto S, Hoppe B, Adragna M, Lopez L, Roldan MLJ, Peña A, Trascasa ML, Sánchez-Corral P, de Córdoba SR: Complement mutations in DGKE-associated atypical hemolytic uremic syndrome. *Clin J Am Soc Nephrol* 9: 1611–1619, 2014
8. Lemaire M, Frémeaux-Bacchi V, Schaefer F, Choi M, Tang WH, Le Quintec M, Fakhouri F, Taque S, Nobili F, Martinez F, Ji W, Overton JD, Mane SM, Nürnberg G, Altmüller J, Thiele H, Morin D, Deschenes G, Baudouin V, Llanas B, Collard L, Majid MA, Simkova E, Nürnberg P, Rioux-Leclerc N, Moeckel GW, Gubler MC, Hwa J, Loirat C, Lifton RP: Recessive mutations in DGKE cause atypical hemolytic-uremic syndrome. *Nat Genet* 45: 531–536, 2013
9. Ozaltin F, Li B, Rauhauser A, An SW, Soylemezoglu O, Gonul II, Taskiran EZ, Ibsirlioglu T, Korkmaz E, Bilginer Y, Duzova A, Ozen S, Topaloglu R, Besbas N, Ashraf S, Du Y, Liang C, Chen P, Lu D, Vadnagara K, Arbuckle S, Lewis D, Wakeland B, Quigg RJ, Ransom RF, Wakeland EK, Topham MK, Bazan NG, Mohan C, Hildebrandt F, Bakkaloglu A, Huang CL, Attanasio M: DGKE variants cause a glomerular microangiopathy that mimics membranoproliferative GN. *J Am Soc Nephrol* 24: 377–384, 2013

10. Westland R, Bodria M, Carrea A, Lata S, Scolari F, Fremeaux-Bacchi V, D'Agati VD, Lifton RP, Gharavi AG, Ghiggeri GM, Sanna-Cherchi S: Phenotypic expansion of DGKE-associated diseases. *J Am Soc Nephrol* 25: 1408–1414, 2014
11. Bento D, Mapril J, Rocha C, Marchbank KJ, Kavanagh D, Barge D, Strain L, Goodship TH, Meneses-Oliveira C: Triggering of atypical hemolytic uremic syndrome by influenza A (H1N1). *Ren Fail* 32: 753–756, 2010
12. Berner R, Krause MF, Gordjani N, Zipfel PF, Boehm N, Krueger M, Brandis M, Zimmerhackl LB: Hemolytic uremic syndrome due to an altered factor H triggered by neonatal pertussis. *Pediatr Nephrol* 17: 190–192, 2002
13. Brackman D, Sartz L, Leh S, Kristoffersson AC, Bjerre A, Tati R, Fremeaux-Bacchi V, Karpman D: Thrombotic microangiopathy mimicking membranoproliferative glomerulonephritis. *Nephrol Dial Transplant* 26: 3399–3403, 2011
14. Bu F, Maga T, Meyer NC, Wang K, Thomas CP, Nester CM, Smith RJ: Comprehensive genetic analysis of complement and coagulation genes in atypical hemolytic uremic syndrome. *J Am Soc Nephrol* 25: 55–64, 2014
15. Chapin J, Eyler S, Smith R, Tsai HM, Laurence J: Complement factor H mutations are present in ADAMTS13-deficient, ticlopidine-associated thrombotic microangiopathies. *Blood* 121: 4012–4013, 2013
16. Delvaeye M, Noris M, De Vriese A, Esmon CT, Esmon NL, Ferrell G, Del-Favero J, Plaisance S, Claes B, Lambrechts D, Zoja C, Remuzzi G, Conway EM: Thrombomodulin mutations in atypical hemolytic-uremic syndrome. *N Engl J Med* 361: 345–357, 2009
17. Edey M, Strain L, Ward R, Ahmed S, Thomas T, Goodship TH: Is complement factor H a susceptibility factor for IgA nephropathy? *Mol Immunol* 46: 1405–1408, 2009
18. El Karoui K, Hill GS, Karras A, Jacquot C, Moulounguet L, Kourilsky O, Frémeaux-Bacchi V, Delahousse M, Duong Van Huyen JP, Loupy A, Bruneval P, Nochy D: A clinicopathologic study of thrombotic microangiopathy in IgA nephropathy. *J Am Soc Nephrol* 23: 137–148, 2012
19. Fakhouri F, Roumenina L, Provot F, Sallée M, Caillard S, Couzi L, Essig M, Ribes D, Dragon-Durey MA, Bridoux F, Rondeau E, Frémeaux-Bacchi V: Pregnancy-associated hemolytic uremic syndrome revisited in the era of complement gene mutations. *J Am Soc Nephrol* 21: 859–867, 2010
20. Feng S, Eyler SJ, Zhang Y, Maga T, Nester CM, Kroll MH, Smith RJ, Afshar-Kharghan V: Partial ADAMTS13 deficiency in atypical hemolytic uremic syndrome. *Blood* 122: 1487–1493, 2013
21. Gilbert RD, Nagra A, Haq MR: Does dysregulated complement activation contribute to haemolytic uraemic syndrome secondary to *Streptococcus pneumoniae*? *Med Hypotheses* 81: 400–403, 2013
22. Gilbert RD, Stanley LK, Fowles DJ, Angus EM, Hardy SA, Goodship TH: Cisplatin-induced haemolytic uraemic syndrome associated with a novel intronic mutation of treated with eculizumab. *Clin Kidney J* 6: 421–425, 2013
23. Guignon V, Fremeaux-Bacchi V, Giraudier S, Favier R, Borderie D, Massy Z, Mougnot B, Rosenblatt DS, Deschenes G: Late-onset thrombotic microangiopathy caused by cblC disease: Association with a factor H mutation. *Am J Kidney Dis* 45: 588–595, 2005
24. Hale GA, Bowman LC, Rochester RJ, Benaim E, Heslop HE, Krance RA, Horwitz EM, Cunningham JM, Tong X, Srivastava DK, Handgretinger R, Jones DP: Hemolytic uremic syndrome after bone marrow transplantation: Clinical characteristics and outcome in children. *Biol Blood Marrow Transplant* 11: 912–920, 2005
25. Jodele S, Fukuda T, Vinks A, Mizuno K, Laskin BL, Goebel J, Dixon BP, Teusink A, Pluthero FG, Lu L, Licht C, Davies SM: Eculizumab therapy in children with severe hematopoietic stem cell transplantation-associated thrombotic microangiopathy. *Biol Blood Marrow Transplant* 20: 518–525, 2014
26. Jodele S, Licht C, Goebel J, Dixon BP, Zhang K, Sivakumaran TA, Davies SM, Pluthero FG, Lu L, Laskin BL: Abnormalities in the alternative pathway of complement in children with hematopoietic stem cell transplant-associated thrombotic microangiopathy. *Blood* 122: 2003–2007, 2013
27. Kömhoff M, Roofthoof MT, Westra D, Teertstra TK, Losito A, van de Kar NC, Berger RM: Combined pulmonary hypertension and renal thrombotic microangiopathy in cobalamin C deficiency. *Pediatrics* 132: e540–e544, 2013
28. Kwon T, Belot A, Ranchin B, Baudouin V, Fremeaux-Bacchi V, Dragon-Durey MA, Cochat P, Loirat C: Varicella as a trigger of atypical haemolytic uraemic syndrome associated with complement dysfunction: Two cases. *Nephrol Dial Transplant* 24: 2752–2754, 2009
29. Le Quintrec M, Lionet A, Kamar N, Karras A, Barbier S, Buchler M, Fakhouri F, Provost F, Fridman WH, Thervet E, Legendre C, Zuber J, Fremeaux-Bacchi V: Complement mutation-associated de novo thrombotic microangiopathy following kidney transplantation. *Am J Transplant* 8: 1694–1701, 2008
30. Lorcy N, Rioux-Leclercq N, Lombard ML, Le Pogamp P, Vigneau C: Three kidneys, two diseases, one antibody? *Nephrol Dial Transplant* 26: 3811–3813, 2011
31. Noone D, Al-Matrafi J, Tinckam K, Zipfel PF, Herzenberg AM, Thorner PS, Pluthero FG, Kahr WH, Filler G, Hebert D, Harvey E, Licht C: Antibody mediated rejection associated with complement factor H-related protein 3/1 deficiency successfully treated with eculizumab. *Am J Transplant* 12: 2546–2553, 2012
32. Obando I, Camacho MS, Falcon-Neyra D, Hurtado-Mingo A, Neth O: Atypical hemolytic uremic syndrome associated with *Bordetella pertussis* infection. *Pediatr Infect Dis J* 31: 1210, 2012
33. Réti M, Farkas P, Csuka D, Rázsó K, Schlammadinger Á, Udvardy ML, Madách K, Domján G, Bereczki C, Reusz GS, Szabó AJ, Prohászka Z: Complement activation in thrombotic thrombocytopenic purpura. *J Thromb Haemost* 10: 791–798, 2012
34. Schmitt R, Krmar RT, Kristoffersson A, Söderberg M, Karpman D: IgA nephropathy associated with a novel N-terminal mutation in factor H. *Eur J Pediatr* 170: 107–110, 2011
35. Szilagyí A, Kiss N, Bereczki C, Talosi G, Racz K, Turi S, Gyorko Z, Simon E, Horvath E, Kelen K, Reusz GS, Szabo AJ, Tulassay T, Prohászka Z: The role of complement in *Streptococcus pneumoniae*-associated haemolytic uraemic syndrome. *Nephrol Dial Transplant* 28: 2237–2245, 2013
36. Thurman JM, Marians R, Emlen W, Wood S, Smith C, Akana H, Holers VM, Lesser M, Kline M, Hoffman C, Christen E, Trachtman H: Alternative pathway of complement in children with diarrhea-associated hemolytic uremic syndrome. *Clin J Am Soc Nephrol* 4: 1920–1924, 2009
37. Totina A, Iorember F, El-Dahr SS, Yosypiv IV: Atypical hemolytic-uremic syndrome in a child presenting with malignant hypertension. *Clin Pediatr (Phila)* 52: 183–186, 2013
38. Westwood JP, Langley K, Heelas E, Machin SJ, Scully M: Complement and cytokine response in acute thrombotic thrombocytopenic purpura. *Br J Haematol* 164: 858–866, 2014
39. Wu TC, Yang S, Haven S, Holers VM, Lundberg AS, Wu H, Cataland SR: Complement activation and mortality during an acute episode of thrombotic thrombocytopenic purpura. *J Thromb Haemost* 11: 1925–1927, 2013

Published online ahead of print. Publication date available at www.cjasn.org.

See related article, “Complement Mutations in Diacylglycerol Kinase-ε-Associated Atypical Hemolytic Uremic Syndrome,” on pages 1611–1619.