

Mirela Dobre¹, Sarah A. Gaussoin², Jeffrey T. Bates^{3,4}, Michel B. Chonchol⁵, Debbie L. Cohen⁶, Thomas H. Hostetter¹, Kalani L. Raphael⁷, Addison A. Taylor^{3,4}, Alan J. Lerner¹, Jackson T. Wright Jr¹, Mahboob Rahman^{1,8} for the SPRINT Research Group

¹Division of Nephrology and Hypertension, Department of Medicine, University Hospitals Cleveland Medical Center, Case Western Reserve University, Cleveland, Ohio;

²Division of Public Health Sciences, Department of Biostatistical Sciences, Wake Forest School of Medicine, Winston–Salem, North Carolina;

³Department of Medicine, Baylor College of Medicine, Houston, Texas;

⁴Department of Medicine, Michael E. DeBakey VA Medical Center, Houston, Texas;

⁵Medicine-Renal Medical Diseases/Hypertension, University of Colorado Denver Anschutz Medical Campus, Aurora, Colorado;

⁶Department of Medicine, Hospital at the University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania;

⁷Division of Nephrology and Hypertension, Department of Medicine, Salt Lake City Veterans Affairs Healthcare System, University of Utah, Salt Lake City, Utah; and

⁸Division of Nephrology, Department of Medicine, Louis Stokes Cleveland VA Medical Center, Cleveland, Ohio